

Portland State University

PDXScholar

Mathematics and Statistics Faculty
Publications and Presentations

Fariborz Maseeh Department of Mathematics
and Statistics

10-1-2000

Dependence Among Spacings

Baha-Eldin Khaledi

Subhash C. Kochar
Portland State University

Follow this and additional works at: https://pdxscholar.library.pdx.edu/mth_fac



Part of the [Mathematics Commons](#)

Let us know how access to this document benefits you.

Citation Details

Khaledi, E. and Kochar, S. (2000). Dependence among spacings. *Probability in the Engineering and Informational Sciences*, 14, pp 461-472.

This Article is brought to you for free and open access. It has been accepted for inclusion in Mathematics and Statistics Faculty Publications and Presentations by an authorized administrator of PDXScholar. Please contact us if we can make this document more accessible: pdxscholar@pdx.edu.

DEPENDENCE AMONG SPACINGS

BAHA-ELDIN KHALEDI AND SUBHASH KOCHAR

*Indian Statistical Institute
New Delhi-110016, India
E-mail: kochar@isid.ac.in*

In this paper, we study the dependence properties of spacings. It is proved that if X_1, \dots, X_n are exchangeable random variables which are TP_2 in pairs and their joint density is log-convex in each argument, then the spacings are MTP_2 dependent. Next, we consider the case of independent but nonhomogeneous exponential random variables. It is shown that in this case, in general, the spacings are not MTP_2 dependent. However, in the case of a single outlier when all except one parameters are equal, the spacings are shown to be MTP_2 dependent and, hence, they are associated. A consequence of this result is that in this case, the variances of the order statistics are increasing. It is also proved that in the case of the multiple-outliers model, all consecutive pairs of spacings are TP_2 dependent.

1. INTRODUCTION

Let X_1, \dots, X_n be n random variables. We shall denote by $X_{i:n}$ the i th-order statistic of X_1, \dots, X_n . Let $D_{i:n} = X_{i:n} - X_{i-1:n}$ denote the i th spacing, $i = 1, \dots, n$, with $X_{0:n} \equiv 0$. It is well known that if X_1, \dots, X_n is a random sample from an exponential distribution, then $D_{1:n}, \dots, D_{n:n}$ are independent. In this paper, we study the dependence properties of spacings when X_i 's are not necessarily independent and identically distributed as exponentials. The related problem of stochastic orderings among spacings has been extensively studied in the literature. For details, the reader is referred to a recent review paper on this topic by Kochar [9]. Throughout this paper, *increasing* means nondecreasing and *decreasing* means nonincreasing.

There are several notions of positive and negative dependences among random variables with varying degrees of strength. There is a vast literature on this topic, with important contributions by Lehmann [11], Esary and Proschan [3], Barlow and Proschan [1], Block and Ting [2], and Karlin and Rinott [6,7], among others. Per-

haps the strongest notion of positive dependence between random variables S and T is that of TP_2 dependence (also known as *likelihood* dependence). S and T are TP_2 dependent if their joint density $f(s, t)$ is totally positive of order 2 in s and t , or, more precisely, if

$$\begin{vmatrix} f(s_1, t_1) & f(s_1, t_2) \\ f(s_2, t_1) & f(s_2, t_2) \end{vmatrix} \geq 0 \tag{1.1}$$

whenever $s_1 < s_2$ and $t_1 < t_2$.

We say that T is *right-tail increasing* in S if $P[T > t | S > s]$ is increasing in s for all t , and we denote this relationship by $RTI(T|S)$. Finally, random variables S and T are *associated* [written $A(S, T)$] if $\text{cov}[\Gamma(S, T), \Delta(S, T)] \geq 0$ for all pairs of increasing binary functions Γ and Δ . As shown in Barlow and Proschan [1, p. 143], the following chain of implications holds among the above notions of positive dependence:

$$TP_2(S, T) \Rightarrow RTI(S|T) \Rightarrow A(S, T). \tag{1.2}$$

There are many other notions of dependence, but we will not discuss them here.

These concepts of bivariate dependence can be easily extended to the multivariate case. A function $\psi : R^n \rightarrow [0, \infty)$ is said to be *multivariate total positivity of order 2* (denoted by MTP_2) if

$$\psi(\mathbf{x})\psi(\mathbf{y}) \leq \psi(\mathbf{x} \wedge \mathbf{y})\psi(\mathbf{x} \vee \mathbf{y}) \quad \text{for every } \mathbf{x} \text{ and } \mathbf{y} \text{ in } \mathcal{R}^n,$$

where $\mathbf{x} \wedge \mathbf{y} = (\min(x_1, y_1), \dots, \min(x_n, y_n))$ and $\mathbf{x} \vee \mathbf{y} = (\max(x_1, y_1), \dots, \max(x_n, y_n))$. Random variables X_1, \dots, X_n are said to be MTP_2 dependent if their joint density function is MTP_2 . It is shown in Kemperman [8] (see also Block and Ting [2]) that if the support of a random vector $\mathbf{X} = (X_1, \dots, X_n)$ is a lattice (i.e., if \mathbf{x} and \mathbf{y} are in the support of \mathbf{X} , then so are $\mathbf{x} \wedge \mathbf{y}$ and $\mathbf{x} \vee \mathbf{y}$), then X is MTP_2 if and only if its density function f is TP_2 in each pair of its variables when the other $n - 2$ variables are held fixed. See Karlin and Rinott [6] for more details on the properties of MTP_2 functions. Random variables X_1, \dots, X_n are *conditionally increasing in sequence* if $P[X_i > x | X_1 = x_1, \dots, X_{i-1} = x_{i-1}]$ is increasing in x_1, \dots, x_{i-1} for $i = 2, \dots, n$. Finally, a set of random variables $\mathbf{X} = (X_1, \dots, X_n)$ are associated if $\text{cov}(u(\mathbf{X}), v(\mathbf{X})) \geq 0$ for all increasing binary functions u and v . Karlin and Rinott [6] proved that if a set of random variables are MTP_2 dependent, then they are conditionally increasing in sequence, which, in turn, implies that they are associated (cf. Barlow and Proschan [1, p. 146]), a result which extends (1.2) to the multivariate case.

It is known that spacings of a random sample from a DFR (decreasing failure rate) distribution are conditionally increasing in sequence (cf. Barlow and Proschan [1, p. 151]). Karlin and Rinott [6] have pointed out that if the DFR assumption is strengthened to assume that the parent distribution has a log-convex density, then the spacings have the corresponding stronger property of being MTP_2 dependent. In Section 2, we extend this result to the case when the random variables X_1, \dots, X_n are

dependent. It is proved that if the joint probability density function (p.d.f.) of X_i 's is permutation symmetric, TP_2 in pairs, and log-convex in each argument, then their spacings are MTP_2 dependent (Theorem 2.1). In Section 3, we study the dependence properties of spacings of independent but nonidentically distributed exponential random variables. We show with the help of a counterexample that in this case, the spacings may not be MTP_2 dependent. In fact, for $n = 3$, even $RTI(D_{3:3}|D_{2:3})$ does not hold for some values of the parameters (Example 3.1). However, it is shown that $cov(D_{2:3}, D_{3:3})$ is nonnegative (Corollary 3.1) due to its Schur convexity (Theorem 3.2). It is also proved that in the case of a single outlier when all except one of the parameters are equal, the spacings are MTP_2 dependent (Theorem 3.4). A consequence (Corollary 3.2) of this result is that in this case, $var(X_{1:n}) \leq var(X_{2:n}) \leq \dots \leq var(X_{n:n})$. We also prove that in the case of the multiple-outliers model (Theorem 3.5), any pair of consecutive spacings $D_{i:n}$ and $D_{i+1:n}$ are TP_2 dependent for $i = 1, \dots, n - 1$.

2. DEPENDENCE AMONG SPACINGS OF EXCHANGEABLE RANDOM VARIABLES

As pointed out in Karlin and Rinott [6, p. 483], the spacings of a random sample from a distribution with log-convex density are MTP_2 dependent. In Theorem 2.1, we extend this result to the case when random variables are exchangeable and TP_2 in pairs.

THEOREM 2.1: *Let X_1, \dots, X_n be exchangeable random variables with absolutely continuous joint p.d.f. $f_{\mathbf{X}}(x_1, \dots, x_n)$, which is positive on $\Pi_{i=1}^n \Omega_i^n$, $\Omega_i \subset \mathcal{R}^1$, $i = 1, \dots, n$, and satisfies the following conditions:*

- (a) $f_{\mathbf{X}}$ is TP_2 in pairs.
- (b) $f_{\mathbf{X}}$ is log-convex in each argument when remaining arguments are held fixed.
- (c) The first partial derivative of $f_{\mathbf{X}}(\mathbf{x})$ with respect to x_i exists for $i = 1, \dots, n$.

Then, $D_{1:n}, \dots, D_{n:n}$ are MTP_2 dependent.

PROOF: The joint p.d.f. of $D_{1:n}, \dots, D_{n:n}$ is

$$f_{\mathbf{D}}(d_1, \dots, d_n) = n! f_{\mathbf{X}}\left(d_1, \sum_{j=1}^2 d_j, \dots, \sum_{j=1}^i d_j, \dots, \sum_{j=1}^n d_j\right).$$

By Theorem 1.5 of Karlin [5], $f_{\mathbf{D}}(d_1, \dots, d_n)$ will be TP_2 in pairs of d_1, \dots, d_n if and only if for any $i \neq j$, $1 \leq i, j \leq n$, $(\partial/\partial d_i) \log f_{\mathbf{D}}(d_1, \dots, d_n)$ is increasing in d_j . Let $i < j$. By the chain rule of differentiation,

$$\left(\frac{\partial}{\partial d_i}\right) \log f_{\mathbf{D}}(d_1, \dots, d_n) = \sum_{k=i}^n \left(\frac{\partial}{\partial x_k}\right) \log f_{\mathbf{X}}\left(d_1, \sum_{j=1}^2 d_j, \dots, \sum_{j=1}^i d_j, \dots, \sum_{j=1}^n d_j\right), \tag{2.1}$$

where $x_k = \sum_{l=1}^k d_l$ for $k \in \{1, \dots, n\}$. The term $(\partial/\partial x_k) \log f_{\mathbf{X}}(\mathbf{x})$ is increasing in x_k for $k \in \{1, \dots, n\}$, as $f_{\mathbf{X}}$ is log-convex in x_k for each k . It is increasing in $x_m, m \neq k, m \in \{1, \dots, n\}$ because $f_{\mathbf{X}}$ is TP₂ in pairs. Now, x_m and x_k are both increasing functions of d_j . This implies that $(\partial/\partial d_i) \log f_{\mathbf{D}}(d_1, \dots, d_n)$ is an increasing function of d_j . Hence, $f_{\mathbf{D}}(d_1, \dots, d_n)$ is TP₂ in pairs. Clearly, the support of spacings is a lattice under the given conditions. Combining these facts, we get the required result. ■

Remark: In Theorem 2.1, if instead of conditions (a) and (b), we assume that $f_{\mathbf{X}}$ is RR₂ (reverse regular of order 2) [two random variables S and T are RR₂ dependent if the inequality in (1.1) is reversed] in pairs and $f_{\mathbf{X}}$ is log-concave in each argument, then one can prove that the joint p.d.f. of spacings is RR₂ in pairs.

LEMMA 2.1: For a bivariate random vector (X, Y) ,

$$\text{cov}(Y - X, X) \geq 0 \Rightarrow \text{var}(X) \leq \text{var}(Y). \tag{2.2}$$

PROOF: The inequality $\text{cov}(Y - X, X) \geq 0$ implies $\text{cov}(X, Y) \geq \text{var}(X)$, which, in turn, implies that

$$\left\{ \frac{\text{var}(X)}{\text{var}(Y)} \right\} \leq \rho^2(X, Y) \leq 1,$$

where $\rho(X, Y)$ is the correlation coefficient between X and Y . The required result follows from this. ■

This lemma and Theorem 2.1 lead to the following interesting corollary.

COROLLARY 2.1: Under the assumptions of Theorem 2.1,

$$\text{var}(X_{1:n}) \leq \text{var}(X_{2:n}) \leq \dots \leq \text{var}(X_{n:n}).$$

PROOF: Because under the given conditions $D_{i:n}$'s are MTP₂ dependent, they are associated. This implies that for $n = 1, \dots, j - 1$,

$$\text{cov}(X_{j:n} - X_{j-1:n}, X_{j-1:n}) \equiv \text{cov}\left(D_{j:n}, \sum_{i=1}^{j-1} D_{i:n}\right) \geq 0, \tag{2.3}$$

as $\sum_{i=1}^{j-1} D_{i:n}$ and $D_{j:n}$ are increasing functions of $(D_{1:n}, \dots, D_{n:n})$. The required result follows from Lemma 2.1. ■

Example 2.1 (Inverted Dirichlet Distribution): Let $X_i, i = 0, \dots, n$, be independent gamma random variables each with scale parameter 1 such that X_0 has shape parameter β and X_i has shape parameter α , for $i \in \{1, \dots, n\}$. Then, the joint p.d.f. of $Y_i = X_i/X_0, i = 1, \dots, n$, is

$$f_{Y_1, \dots, Y_n}(y_1, \dots, y_n) = \frac{\Gamma(n\alpha + \beta)}{(\Gamma(\alpha))^n \Gamma(\beta)} \frac{\left(\prod_{i=1}^n y_i\right)^{\alpha-1}}{\left(1 + \sum_{i=1}^n y_i\right)^{n\alpha+\beta}} \quad \text{for } y_i \geq 0.$$

It is easy to see that $f_{Y_1, \dots, Y_n}(y_1, \dots, y_n)$ is exchangeable, TP_2 in pairs, and log-convex in each argument when $0 < \alpha < 1$ and $n\alpha + \beta \geq 1$. Thus, the conditions of Theorem 2.1 are satisfied, and as a result, the spacings of Y_1, \dots, Y_n are MTP_2 dependent. By Corollary 2.1, the variances of the successive order statistics increase as i goes from 1 to n .

3. THE CASE OF HETEROGENEOUS EXPONENTIALS

The exponential distribution plays a central role in reliability theory. In this section, we study the dependence properties of spacings when the observations X_1, \dots, X_n are independent with X_i having exponential distribution with parameter $\lambda_i, i = 1, \dots, n$. Their joint density is given by (cf. Kochar and Korwar [10])

$$f_{D_{1:n}, \dots, D_{n:n}}(x_1, \dots, x_n) = \sum_{(\mathbf{r})} \frac{\prod_{i=1}^n \lambda_i}{\prod_{i=1}^n \sum_{j=i}^n \lambda(r_j)} \prod_{i=1}^n \left(\sum_{j=i}^n \lambda(r_j) \right) \exp \left\{ -x_i \sum_{j=i}^n \lambda(r_j) \right\} \tag{3.1}$$

for $x_i \geq 0, i = 1, \dots, n$, where $(\mathbf{r}) = (r_1, \dots, r_n)$ is a permutation of $(1, \dots, n)$ and $\lambda(i) = \lambda_i$. It is a mixture of products of exponential distributions. From (3.1), it is easy to find that the joint p.d.f. of $(D_{i:n}, D_{j:n})$ for $1 \leq i < j \leq n$ is

$$f_{D_{i:n}, D_{j:n}}(x, y) = \sum_{(\mathbf{r})} \frac{\prod_{i=1}^n \lambda_i}{\prod_{i=1}^n \sum_{j=i}^n \lambda(r_j)} \times \left(\sum_{m=i}^n \lambda(r_m) \right) \exp \left\{ -x \sum_{m=i}^n \lambda(r_m) \right\} \left(\sum_{m=j}^n \lambda(r_m) \right) \exp \left\{ -y \sum_{m=j}^n \lambda(r_m) \right\} \tag{3.2}$$

for $x, y \geq 0$.

The next example shows that the spacings may not be MTP_2 dependent if the λ_i 's are all different.

Example 3.1: Let X_1, X_2 , and X_3 be independent exponential random variables with respective hazard rates 5, 4, and 1. Using (3.2), we find, after some simplifications, that

$$h(y) = P(D_{3:3} > 2 | D_{2:3} > y) = \frac{20e^{20y} \left(\left(\frac{1}{9e^{9y}} + \frac{1}{6e^{6y}} \right) \left(\frac{1}{5} \right) e^{-10} + \left(\frac{1}{9e^{9y}} + \frac{1}{5e^{5y}} \right) \left(\frac{1}{4} \right) e^{-8} + \left(\frac{1}{6e^{6y}} + \frac{1}{5e^{5y}} \right) e^{-2} \right)}{e^{11y} + 4e^{14y} + 5e^{15y}}.$$

It is clear from Figure 1 that the function $h(y)$ is not monotonically increasing, proving thereby that $D_{3:3}$ is even not RTI in $D_{2:3}$. Hence, $D_{2:3}$ and $D_{3:3}$ are not TP_2 dependent.

The covariance between $D_{i:n}$ and $D_{j:n}$ for $i < j$ is

$$\begin{aligned} \text{cov}(D_{i:n}, D_{j:n}) = & \sum_{(r)} \frac{\prod_{i=1}^n \lambda_i}{\prod_{i=1}^n \sum_{j=i}^n \lambda(r_j)} \left\{ \sum_{m=i}^n \lambda(r_m) \right\}^{-1} \left\{ \sum_{m=j}^n \lambda(r_m) \right\}^{-1} \\ & - \left[\sum_{(r)} \frac{\prod_{i=1}^n \lambda_i}{\prod_{i=1}^n \sum_{j=i}^n \lambda(r_j)} \left\{ \sum_{m=i}^n \lambda(r_m) \right\}^{-1} \right] \left[\sum_{(r)} \frac{\prod_{i=1}^n \lambda_i}{\prod_{i=1}^n \sum_{j=i}^n \lambda(r_j)} \left\{ \sum_{m=j}^n \lambda(r_m) \right\}^{-1} \right]. \end{aligned}$$

We conjecture that, in general, the covariance between $D_{i:n}$ and $D_{j:n}$ for $i < j$ is nonnegative. We prove this conjecture for $n = 3$ in Corollary 3.1. In fact, we prove in Theorem 3.2 that the covariance between $D_{2:3}$ and $D_{3:3}$ is Schur convex in λ_i 's.

Let $\{x_{(1)} \leq x_{(2)} \leq \dots \leq x_{(n)}\}$ denote the increasing arrangement of the components of the vector $\mathbf{x} = (x_1, x_2, \dots, x_n)$. The vector \mathbf{y} is said to majorize the vector \mathbf{x} (written $\mathbf{x} \preceq \mathbf{y}$) if $\sum_{i=1}^j y_{(i)} \leq \sum_{i=1}^j x_{(i)}$ for $j = 1, \dots, n - 1$ and $\sum_{i=1}^n y_{(i)} = \sum_{i=1}^n x_{(i)}$. Functions that preserve the ordering of majorization are said to be Schur convex; that is, a real function ϕ defined on a set $\mathcal{A} \subset R^n$ is said to be Schur convex on \mathcal{A} if $\mathbf{x} \preceq \mathbf{y} \Rightarrow \phi(\mathbf{x}) \leq \phi(\mathbf{y})$. See Marshall and Olkin [12, Chap. 3] for properties and more




FIGURE 1. Graph of $h(y)$.

details of such functions. The following characterization of Schur-convex functions will be used to prove Theorem 3.2.

THEOREM 3.1. (cf. Marshall and Olkin [12, p. 57]: *Let $I \subset R$ be an open interval and let $\phi : I^n \rightarrow R$ be continuously differentiable. Necessary and sufficient conditions for ϕ to be Schur convex on I^n are ϕ is symmetric on I^n and, for all $i \neq j$,*

$$(z_i - z_j)[\phi_{(i)}(z_i) - \phi_{(j)}(z_j)] \geq 0 \quad \text{for all } z \in I^n,$$

where $\phi_{(i)}(z)$ denotes the partial derivative of ϕ with respect to its i th argument.

THEOREM 3.2: *Let $X_1, X_2,$ and X_3 be independent exponential random variables having hazard rates $\lambda_1, \lambda_2,$ and $\lambda_3,$ respectively. Then, $\text{cov}(D_{2:3}, D_{3:3})$ is Schur convex in λ_i 's.*

PROOF: The covariance between $D_{2:3}$ and $D_{3:3}$ is

$$\begin{aligned} \phi(\lambda_1, \lambda_2, \lambda_3) &= \text{cov}(D_{2:3}, D_{3:3}) \\ &= (\lambda_1 \lambda_2 \lambda_3)(\lambda_1 + \lambda_2 + \lambda_3)^{-1} \\ &\quad \times [(\lambda_1^{-2} + \lambda_2^{-2})(\lambda_1 + \lambda_2)^{-2} + (\lambda_1^{-2} + \lambda_3^{-2})(\lambda_1 + \lambda_3)^{-2} \\ &\quad + (\lambda_2^{-2} + \lambda_3^{-2})(\lambda_2 + \lambda_3)^{-2}] \\ &\quad - \{(\lambda_1 + \lambda_2 + \lambda_3)^{-1}(\lambda_3/(\lambda_1 + \lambda_2) + \lambda_2/(\lambda_1 + \lambda_3) \\ &\quad \quad + \lambda_1/(\lambda_2 + \lambda_3))\} \\ &\quad \times \{(\lambda_1 \lambda_2 \lambda_3)(\lambda_1 + \lambda_2 + \lambda_3)^{-1} \\ &\quad \quad \times \{(\lambda_2^{-2} + \lambda_3^{-2})(\lambda_2 + \lambda_3)^{-1} \\ &\quad \quad + (\lambda_1^{-2} + \lambda_3^{-2})(\lambda_1 + \lambda_3)^{-1} \\ &\quad \quad + (\lambda_1^{-2} + \lambda_2^{-2})(\lambda_1 + \lambda_2)^{-1}\}\}. \end{aligned} \tag{3.3}$$

After some simplifications, we find that $(\lambda_1 - \lambda_2)\{\phi_{(1)}(\lambda_1, \lambda_2, \lambda_3) - \phi_{(2)}(\lambda_1, \lambda_2, \lambda_3)\}$ is equal to

$$\frac{8(\lambda_1 - \lambda_2)^2 \lambda_3^2}{(\lambda_1 + \lambda_2)(\lambda_1 + \lambda_3)(\lambda_2 + \lambda_3)(\lambda_1 + \lambda_2 + \lambda_3)},$$

which is nonnegative for all $\lambda_1, \lambda_2, \lambda_3 > 0$. Because the function ϕ is symmetric in $(\lambda_1, \lambda_2, \lambda_3)$, the required result follows from Theorem 3.1. ■

COROLLARY 3.1: *Under the assumptions of Theorem 3.2, $\text{cov}(D_{2:3}, D_{3:3}) \geq 0$ and $\text{var}(X_{1:3}) \leq \text{var}(X_{2:3}) \leq \text{var}(X_{3:3})$.*

PROOF: Let $\bar{\lambda}$ be the average of λ_i 's. It is easy to see that $(\bar{\lambda}, \bar{\lambda}, \bar{\lambda}) \preceq^m (\lambda_1, \lambda_2, \lambda_3)$. From Theorem 3.2, we get

$$\phi(\bar{\lambda}, \bar{\lambda}, \bar{\lambda}) \leq \phi(\lambda_1, \lambda_2, \lambda_3), \tag{3.4}$$

where the function ϕ is given by (3.3). The left-hand side of (3.4) is zero, since spacings of a random sample from an exponential distribution are independent. This proves that $\text{cov}(D_{2:3}, D_{3:3}) \geq 0$. Since $D_{1:3}$ is independent of $D_{2:3}$ and $D_{3:3}$, it follows that $\text{cov}(X_{3:3} - X_{2:3}, X_{2:3}) \geq 0$. The required result follows from Lemma 2.1. ■

Gross, Hunt, and Odeh [4] considered the single-outlier model in which all except one of the λ_i 's are equal; that is, $\lambda_1 = \lambda$ and $\lambda_2 = \dots = \lambda_n = \lambda^*$, $\lambda \neq \lambda^*$. They incorrectly pointed out that in this case, the spacings $D_{i:n}$ and $D_{j:n}$ are independent for $j - i \geq 2$. Although it is true that $D_{1:n}$ is independent of $(D_{2:n}, \dots, D_{n:n})$, the other D_i 's are not independent. In fact, for $n = 4$,

$$\text{cov}(D_{2:4}, D_{4:4}) = \frac{2\lambda^*(\lambda^* - \lambda)^2}{(\lambda^* + \lambda)(2\lambda^* + \lambda)^2(3\lambda^* + \lambda)^2},$$

which is positive unless $\lambda^* = \lambda$. Theorem 3.4, which follows, replaces the incorrect result of Gross et al. [4] for the single-outlier model.

To prove the remaining results of this section, we shall repeatedly use the following known result.

THEOREM 3.3 (Shaked and Spizzichino [13]): *Let the joint distribution function of $\mathbf{X} = (X_1, \dots, X_n)$ be*

$$F(x_1, \dots, x_n) = \int_{-\infty}^{+\infty} \prod_{i=1}^n F_i(x_i | \theta) dG(\theta),$$

where $F_i(\cdot | \theta)$ is an absolutely continuous distribution function with respect to Lebesgue measure on R for each θ in the support of Θ with density function $f_i(\cdot | \theta)$ for $i = 1, \dots, n$. Suppose that the support of (X_1, \dots, X_n) is a lattice. If $f_i(x | \theta)$ is TP_2 (RR_2) in (x, θ) for all $i \in \{1, \dots, n\}$, then (X_1, \dots, X_n) is MTP_2 .

In the next theorem, we prove that in the case of a single-outlier exponential model, the spacings are MTP_2 dependent.

THEOREM 3.4: *Let X_i , $i = 1, \dots, n$, be independent exponential random variables such that X_1 has hazard rate λ and X_i has hazard rate λ^* for $i \in \{2, \dots, n\}$. Then, $(D_{1:n}, \dots, D_{n:n})$ is MTP_2 dependent.*

PROOF: Using (3.1), we find that the joint p.d.f. of $(D_{1:n}, \dots, D_{n:n})$ in this case is

$$\begin{aligned} f_{D_{1:n}, \dots, D_{n:n}}(x_1, \dots, x_n) &= \sum_{\theta=1}^n \frac{(n-1)! \lambda (\lambda^*)^{n-1}}{\prod_{i=1}^{\theta} ((n-i)\lambda^* + \lambda) \prod_{i=\theta+1}^n (n-i+1)\lambda^*} \\ &\times \prod_{i=1}^{\theta} ((n-i)\lambda^* + \lambda) e^{-((n-i)\lambda^* + \lambda)x_i} \\ &\times \prod_{i=\theta+1}^n (n-i+1)\lambda^* e^{-(n-i+1)\lambda^* x_i}, \end{aligned}$$

which can be expressed as

$$f_{D_{1:n}, \dots, D_{n:n}}(x_1, \dots, x_n) = \int_{-\infty}^{+\infty} \prod_{i=1}^n f_{D_{i:n}}(x_i | \theta) dP_{\Theta}(\theta),$$

where Θ is a discrete random variable with the probability mass function,

$$p_{\Theta}(\theta) = \frac{(n-1)! \lambda (\lambda^*)^{n-1}}{\prod_{i=1}^{\theta} ((n-i)\lambda^* + \lambda) \prod_{i=\theta+1}^n (n-i+1)\lambda^*} \quad \text{for } \theta = 1, \dots, n,$$

and

$$f_{D_{i:n}}(x | \theta) = \begin{cases} ((n-i)\lambda^* + \lambda) e^{-((n-i)\lambda^* + \lambda)x}, & i \leq \theta \\ (n-i+1)\lambda^* e^{-(n-i+1)\lambda^*x}, & i \geq \theta + 1. \end{cases} \tag{3.5}$$

We show that the conditional densities as given by (3.5) are all TP₂ if $\lambda < \lambda^*$ and are all RR₂ if $\lambda > \lambda^*$. Suppose $\theta_1 < \theta_2$ and $\theta_1, \theta_2 \in \{1, \dots, n\}$. Then, the ratio

$$\frac{f_{D_{i:n}}(x | \theta_2)}{f_{D_{i:n}}(x | \theta_1)} = \begin{cases} 1, & i \leq \theta_1 \\ \frac{((n-i)\lambda^* + \lambda) e^{-((n-i)\lambda^* + \lambda)x}}{(n-i+1)\lambda^* e^{-(n-i+1)\lambda^*x}}, & \theta_1 < i \leq \theta_2 \\ 1, & \theta_2 < i \end{cases}$$

is increasing (decreasing) in x if $\lambda < (>) \lambda^*$ for $i = 1, \dots, n$; that is, $f_{D_{i:n}}(x | \theta)$ is TP₂ (RR₂) in (x, θ) . The required result follows from Theorem 3.1. ■

COROLLARY 3.2: *Under the assumptions of Theorem 3.4,*

$$\text{var}(X_{1:n}) \leq \text{var}(X_{2:n}) \leq \dots \leq \text{var}(X_{n:n}).$$

In the next theorem, we consider the multiple-outliers model. According to this model, X_1, \dots, X_k are i.i.d. exponentials with hazard rate λ and X_{k+1}, \dots, X_n are i.i.d. exponentials with hazard rate λ^* , where $k \in \{2, \dots, n-2\}$. We prove that in this case, $D_{i:n}$ and $D_{i+1:n}$ are TP₂ dependent for $i = 1, \dots, n-1$. It is not known whether the spacings are MTP₂ dependent in this case.

THEOREM 3.5: *Let $X_i, i = 1, \dots, n$, be independent exponential random variables such that X_i has hazard rate λ for $i \in \{1, \dots, k\}$ and hazard rate λ^* for $i \in \{k+1, \dots, n\}$, $k \in \{2, \dots, n-2\}$. Then, $D_{i:n}$ and $D_{i+1:n}$ are TP₂ dependent.*

PROOF: Without loss of generality, we assume that $k \leq n-k$.

Case (i): Let $k < i \leq n-k$. From (3.2), the joint p.d.f. of $(D_{i:n}, D_{i+1:n})$ for this set of λ_i 's can be expressed as

$$f_{D_{i:n}, D_{i+1:n}}(x, y) = \int_{-\infty}^{+\infty} f_{D_{i:n}}(x | \theta) f_{D_{i+1:n}}(y | \theta) dP_{\Theta}(\theta),$$

where Θ is a discrete random variable taking values $0, 1, 2, \dots, 2k$ with the following probability mass function. For $\theta = 0, 2, 4, \dots, 2k$,

$$p_{\Theta}(\theta) = \lambda^k (\lambda^*)^{n-k} k!(n-k)! \sum_{(\mathbf{r}_{\theta})} \frac{1}{\prod_{i=1}^n \sum_{j=i}^n \lambda(r_j)},$$

where the summation is taken over all permutations of

$$(\mathbf{r}_{\theta}) = (\underbrace{\lambda, \dots, \lambda}_{k-\theta/2}, \underbrace{\lambda^*, \dots, \lambda^*}_{i-1-k+\theta/2}, \underbrace{\lambda^*}_{1}, \underbrace{\lambda, \dots, \lambda}_{\theta/2}, \underbrace{\lambda^*, \dots, \lambda^*}_{n-i-\theta/2}) \tag{3.6}$$

for which the i th component of (\mathbf{r}_{θ}) is λ^* and its last $n - i$ components consist of $(\theta/2)$ λ 's and $(n - i - \theta/2)$ λ^* 's.

For $\theta = 1, 3, 5, \dots, 2k - 1$,

$$p_{\Theta}(\theta) = \lambda^k (\lambda^*)^{n-k} k!(n-k)! \sum_{(\mathbf{r}'_{\theta})} \frac{1}{\prod_{i=1}^n \sum_{j=i}^n \lambda(r'_j)},$$

where the summation is taken over all permutations of

$$(\mathbf{r}'_{\theta}) = (\underbrace{\lambda, \dots, \lambda}_{k-(\theta+1)/2}, \underbrace{\lambda^*, \dots, \lambda^*}_{i-1-k+(\theta+1)/2}, \underbrace{\lambda}_{1}, \underbrace{\lambda, \dots, \lambda}_{(\theta+1)/2-1}, \underbrace{\lambda^*, \dots, \lambda^*}_{n-i-(\theta+1)/2+1}) \tag{3.7}$$

for which the i th component of (\mathbf{r}'_{θ}) is λ and the last $n - i$ components of (\mathbf{r}'_{θ}) consist of $((\theta + 1)/2 - 1)$ λ 's and $(n - i - (\theta + 1)/2 + 1)$ λ^* 's.

For $\theta \in \{0, \dots, 2k\}$,

$$f_{D_{i:n}}(x|\theta) = \{(n - i - [(\theta + 1)/2] + 1)\lambda^* + [(\theta + 1)/2]\lambda\} \times e^{-\{(n-i-[(\theta+1)/2]+1)\lambda^* + [(\theta+1)/2]\lambda\}x}, \tag{3.8}$$

and

$$f_{D_{i+1:n}}(x|\theta) = \{(n - i - [\theta/2])\lambda^* + [\theta/2]\lambda\} e^{-\{(n-i-[\theta/2])\lambda^* + [\theta/2]\lambda\}x}, \tag{3.9}$$

where $[x]$ denotes the greatest integer less than or equal to x .

To prove the required result, we show that $f_{D_{i:n}}(x|\theta)$ and $f_{D_{i+1:n}}(x|\theta)$ are all TP₂ if $\lambda < \lambda^*$ and are all RR₂ if $\lambda > \lambda^*$.

$$\frac{f_{D_{i:n}}(x|\theta + 1)}{f_{D_{i:n}}(x|\theta)} = \frac{\{(n - i - [(\theta + 2)/2] + 1)\lambda^* + [(\theta + 2)/2]\lambda\} e^{-\{(n-i-[(\theta+2)/2]+1)\lambda^* + [(\theta+2)/2]\lambda\}x}}{\{(n - i - [(\theta + 1)/2] + 1)\lambda^* + [(\theta + 1)/2]\lambda\} e^{-\{(n-i-[(\theta+1)/2]+1)\lambda^* + [(\theta+1)/2]\lambda\}x}}$$

$$= \begin{cases} 1, & \text{if } \theta = 1, 3, 5, \dots, 2k - 1 \\ \frac{\{(n - i - \theta/2)\lambda^* + (\theta/2 + 1)\lambda\}}{\{(n - i - \theta/2 + 1)\lambda^* + (\theta/2)\lambda\}} e^{-(\lambda - \lambda^*)x}, & \text{if } \theta = 0, 2, 4, \dots, 2k - 2. \end{cases} \tag{3.10}$$

From (3.10), we conclude that if $\lambda < \lambda^*$ ($\lambda > \lambda^*$), then $f_{D_{i:n}}(x|\theta)$ is $TP_2(RR_2)$ for $i = 1, \dots, n$. Similarly, for $f_{D_{i+1:n}}(x|\theta)$, we have

$$\frac{f_{D_{i+1:n}}(x|\theta + 1)}{f_{D_{i+1:n}}(x|\theta)} = \frac{\{(n - i - [(\theta + 1)/2])\lambda^* + [(\theta + 1)/2]\lambda\}e^{-\{(n - i - [(\theta + 1)/2])\lambda^* + [(\theta + 1)/2]\lambda\}x}}{\{(n - i - [\theta/2])\lambda^* + [\theta/2]\lambda\}e^{-\{(n - i - [\theta/2])\lambda^* + [\theta/2]\lambda\}x}}$$

$$= \begin{cases} 1, & \text{if } \theta = 0, 2, 4, \dots, 2k - 2 \\ \frac{\{(n - i - (\theta + 1)/2)\lambda^* + ((\theta + 1)/2)\lambda\}}{\{(n - i - (\theta - 1)/2)\lambda^* + ((\theta - 1)/2)\lambda\}} e^{-(\lambda - \lambda^*)x}, & \text{if } \theta = 1, 3, 5, \dots, 2k - 1. \end{cases}$$

(3.11)

Again, from (3.11), it follows that $f_{D_{i+1:n}}(x|\theta)$ is $TP_2(RR_2)$ if $\lambda < \lambda^*$ ($\lambda > \lambda^*$). Using these observations, the required result follows from Theorem 3.1.

Case (ii): $i > n - k$. In this case for $\theta \in \{0, 2, \dots, 2(n - i)\}$, (\mathbf{r}_θ) is given by (3.6), and for $\theta \in \{1, 3, \dots, 2(n - i) + 1\}$, (\mathbf{r}'_θ) is given by (3.7). Hence, for $\theta \in \{0, 1, 2, \dots, 2(n - i + 1) - 1\}$, $f_{D_{i:n}}(x|\theta)$ and $f_{D_{i+1:n}}(x|\theta)$ are the same as given by (3.8) and (3.9), respectively. The required result follows from the same kind of arguments as in case (i).

Case (iii): $i \leq k$. The proof is similar to the previous case. The vectors (\mathbf{r}_θ) and (\mathbf{r}'_θ) corresponding to (3.6) and (3.7) are as follows. For $\theta = 0, 2, \dots, 2i - 2$,

$$(\mathbf{r}_\theta) = (\underbrace{\lambda, \dots, \lambda}_{i-1-\theta/2}, \underbrace{\lambda^*, \dots, \lambda^*}_{\theta/2}, \underbrace{\lambda}_{1}, \underbrace{\lambda, \dots, \lambda}_{k-i+\theta/2}, \underbrace{\lambda^*, \dots, \lambda^*}_{n-k-\theta/2}),$$

for which the i th component of (\mathbf{r}_θ) is λ and the last $n - i$ components of (\mathbf{r}_θ) consist of $(k - i + \theta/2)$ λ 's and $(n - k - \theta/2)$ λ^* 's.

For $\theta = 1, 3, \dots, 2i - 1$,

$$(\mathbf{r}'_\theta) = (\underbrace{\lambda, \dots, \lambda}_{i-((\theta+1)/2)}, \underbrace{\lambda^*, \dots, \lambda^*}_{((\theta+1)/2)-1}, \underbrace{\lambda^*}_{1}, \underbrace{\lambda, \dots, \lambda}_{k-i+((\theta+1)/2)}, \underbrace{\lambda^*, \dots, \lambda^*}_{n-k-((\theta+1)/2)}),$$

for which the i th component of (\mathbf{r}'_θ) is λ^* and the last $n - i$ components of (\mathbf{r}'_θ) consist of $(k - i + (\theta + 1)/2)$ λ 's and $(n - k - (\theta + 1)/2)$ λ^* 's.

Therefore, for $\theta \in \{0, \dots, 2i - 1\}$,

$$f_{D_{i:n}}(x|\theta) = \{(k - i + 1 + [\theta/2])\lambda + (n - k - [\theta/2])\lambda^*\} \\ \times e^{-\{(k-i+1+[\theta/2])\lambda+(n-k-[\theta/2])\lambda^*\}x}$$

and

$$f_{D_{i+1:n}}(x|\theta) = \{(k - i - [(\theta + 1)/2])\lambda + (n - k - [(\theta + 1)/2])\lambda^*\} \\ \times e^{-\{(k-i-[(\theta+1)/2])\lambda+(n-k-[(\theta+1)/2])\lambda^*\}x}.$$

The required result follows from the same kind of arguments as in case (i). ■

4. CONCLUDING REMARKS

In this paper, we have obtained some new results on dependence among spacings of heterogeneous independent exponential random variables. Whereas in the case of a single-outlier exponential model, the spacings are shown to be MTP_2 dependent, it is not known whether the same result holds for the multiple-outliers model. In the latter case, we are only able to establish TP_2 dependence between consecutive spacings. Another unsettled question is to examine whether in the case of independent exponential random variables, in general, the spacings are positively correlated. We have given a proof of this conjecture for $n = 3$.

Acknowledgments

One of the authors (B.-E.K.) thanks the Indian Council for Cultural Relations, New Delhi, India, Razi University, Kermanshah, Iran and the Ministry of Culture and Higher Education of the Islamic Republic of Iran, Tehran, Iran for arranging scholarships which enabled him to study at the Indian Statistical Institute, New Delhi, India.

The authors are grateful to the referee for helpful comments and suggestions which greatly improved the presentation of the results.

References

1. Barlow, R.E. & Proschan, F. (1981). *Statistical theory of reliability and life testing*. Silver Spring, MD: To Begin With.
2. Block, H.W. & Ting, M.L. (1981). Some concepts of multivariate dependence. *Communications in Statistics: Theory and Methods* 10: 749–762.
3. Esary, J.D. & Proschan, F. (1972). Relationships among some notions of bivariate dependence. *Annals of Mathematical Statistics* 43: 651–655.
4. Gross, A.J., Hunt, H.H., & Odeh, R.E. (1986). The correlation coefficient between the smallest and largest observations when $(n - 1)$ of the n observations are i.i.d. exponentially distributed. *Communications in Statistics: Theory and Methods* 15: 1113–1123.
5. Karlin, S. (1968). *Total positivity*. Palo Alto, CA: Stanford University Press.
6. Karlin, S. & Rinott, Y. (1980). Classes of orderings of measures and related correlation inequalities. I. Multivariate totally positive distributions. *Journal of Multivariate Analysis* 10: 467–498.
7. Karlin, S. & Rinott, Y. (1980). Classes of orderings of measures and related correlation inequalities. II. Multivariate reverse rule distributions. *Journal of Multivariate Analysis* 10: 499–516.
8. Kemperman, J.H.B. (1977). On the FKG inequality for measures on a partially ordered space. *Indagationes Mathematicae* 39: 313–331.
9. Kochar, S.C. (1998). Stochastic comparisons of spacings and order statistics. In A.P. Basu, S.K. Basu, & S. Mukhopadhyay (eds.), *Frontiers in reliability*. Singapore: World Scientific, pp. 201–216.
10. Kochar, S.C. & Korwar, R. (1996). Stochastic orders for spacings of heterogeneous exponential random variables. *Journal of Multivariate Analysis* 57: 69–83.
11. Lehmann, E.L. (1966). Some concepts of dependence. *Annals of Mathematical Statistics* 37: 1137–1153.
12. Marshall, A.W. & Olkin, I. (1979). *Inequalities: Theory of majorization and its applications*. New York: Academic Press.
13. Shaked, M. & Spizzinchino, F. (1998). Positive dependence properties of conditionally independent random life times. *Mathematics of Operations Research* 23: 944–959.