

Portland State University

PDXScholar

Institute for Sustainable Solutions Publications
and Presentations

Institute for Sustainable Solutions

2-1-2012

Artificial modifications of the coast in response to the Deepwater Horizon oil spill: quick solutions or long-term liabilities?

M. Luisa Martinez

Rusty A. Feagin
Texas A&M University

Kevin M. Yeager
University of Southern Mississippi

John W. Day

Robert Costanza
Portland State University

See next page for additional authors

Follow this and additional works at: https://pdxscholar.library.pdx.edu/iss_pub



Part of the [Sustainability Commons](#)

Let us know how access to this document benefits you.

Citation Details

Artificial modifications of the coast in response to the Deepwater Horizon oil spill: quick solutions or long-term liabilities? M Luisa Martínez, Rusty A Feagin, Kevin M Yeager, John Day, Robert Costanza, et al. *Frontiers in Ecology and the Environment* 2012 10:1, 44-49. doi: 10.1890/100151

This Article is brought to you for free and open access. It has been accepted for inclusion in Institute for Sustainable Solutions Publications and Presentations by an authorized administrator of PDXScholar. Please contact us if we can make this document more accessible: pdxscholar@pdx.edu.

Authors

M. Luisa Martinez, Rusty A. Feagin, Kevin M. Yeager, John W. Day, Robert Costanza, Jim A. Harris, Richard J. Hobbs, Jorge López-Portillo, Ian J. Walker, Eric Higgs, Patricia Moreno-Casasola, Julio Sheinbaum, and Alejandro Yáñez-Arancibia

Artificial modifications of the coast in response to the *Deepwater Horizon* oil spill: quick solutions or long-term liabilities?

M Luisa Martínez^{1*}, Rusty A Feagin², Kevin M Yeager³, John Day⁴, Robert Costanza⁵, Jim A Harris⁶, Richard J Hobbs⁷, Jorge López-Portillo¹, Ian J Walker⁸, Eric Higgs⁹, Patricia Moreno-Casasola¹, Julio Sheinbaum¹⁰, and Alejandro Yáñez-Arancibia¹¹

The *Deepwater Horizon* oil spill threatened many coastal ecosystems in the Gulf of Mexico during the spring and summer of 2010. Mitigation strategies included the construction of barrier sand berms, the restriction or blocking of inlets, and the diversion of freshwater from rivers to the coastal marshes and into the ocean, in order to flush away the oil, on the premise that these measures could reduce the quantity of oil reaching sensitive coastal environments such as wetlands or estuaries. These projects result in changes to the ecosystems that they were intended to protect. Long-term effects include alterations of the hydrological and ecological characteristics of estuaries, changes in sediment transport along the coastal barrier islands, the loss of sand resources, and adverse impacts to benthic and pelagic organisms. Although there are no easy solutions for minimizing the impacts of the *Deepwater Horizon* disaster on coastal ecosystems, we recommend that federal, state, and local agencies return to the strategic use of long-term restoration plans for this region.

Front Ecol Environ 2012; 10(1): 44–49, doi:10.1890/100151 (published online 1 Aug 2011)

On 20 April 2010, the drilling rig *Deepwater Horizon* exploded over the Macondo Prospect oil field and subsequently sank, initiating the release of huge amounts of petroleum and natural gas into the Gulf of Mexico. Almost 3 months later, on 15 July 2010, the corporation responsible for the accident reported that the leak had been stopped by placing a cap on the wellhead.

■ Dimensions of the problem

Considerable uncertainty remains regarding the actual oil-flow rate and the total amount of oil spilled because

of the difficulty of installing accurate measurement devices at the depth at which the well was situated (~1500 m). The best estimate of the spill rate was 60 thousand barrels (~2.5 million gallons) of crude oil per day, which represents a total of about 5.2 million barrels (~216 million gallons) during the 86 days that the well was leaking (Kerr 2010). This was the second largest oceanic oil spill in history, topped only by the combined output of the Arabian Gulf spills that occurred in 1991 (~336 million gallons; NOAA 2010). By comparison, the *Exxon Valdez* oil spill in Alaska in 1989, which generated considerable national and international attention, was much smaller (10.8 million gallons; NOAA 2010).

Because the *Deepwater Horizon* spill originated deep underwater, its effects were “three-dimensional”, contaminating a large volume of the ocean water column, millions of hectares of the ocean surface, and probably large areas of the seafloor as well. Residual oil plumes and tar balls are still impacting more than 100 km of the southern US coastline, and a full cleanup is expected to take years. Estimates vary considerably regarding how much of the oil was removed by the initial attempts at skimming and burning. It has been estimated that perhaps 23% of the oil released from the well was dissolved or evaporated, approximately 13% was dispersed naturally and, as a result of the widespread application of the chemical dispersant COREXIT® 9500, around 16% remains as microscopic droplets suspended in Gulf waters (Lehr *et al.* 2010). The residual amount (~38%) was thought to have been either washed ashore or buried in seafloor sediments. These estimates should be considered with

In a nutshell:

- The *Deepwater Horizon* oil spill seriously impacted coastal ecosystems
- Sand berms were built, inlets were restricted or blocked, and freshwater was diverted from upstream to protect coastal ecosystems from oil exposure
- These projects are likely to compound rather than limit ecological damage
- Financial and other resources would be better used by following established, long-term restoration plans

¹Red de Ecología Funcional, Instituto de Ecología, Xalapa, Veracruz, México (*marisa.martinez@inecol.edu.mx); ²Department of Ecosystem Science and Management, Texas A&M University, College Station, TX; ³Department of Marine Science, University of Southern Mississippi, Stennis Space Center, MS; (continued on p49)



Beyond the Frontier: Listen to Marisa Martínez discussing this research on *Frontiers'* monthly podcast, at www.frontiersinecology.org.

caution, however, because accurate measurements were difficult to make.

■ Ecosystems at risk

Many ecosystems – including deep-sea benthic environments, the pelagic ocean, beaches and dunes, estuaries, and tidal brackish and freshwater wetlands – and numerous species were, and remain, at risk of long-term detrimental effects as a result of the oil spill. These ecosystems provide valuable ecosystem services to society, including fisheries, recreation, habitat refugia, water purification, climate regulation, carbon sequestration, hurricane protection, and nutrient cycling (Batker *et al.* 2010).

The movement of surface waters is unpredictable beyond a few days; ocean currents in the Gulf of Mexico are highly variable and are dominated by the presence of an intense eddy field in the open Gulf, seasonal changes in continental shelf currents, tides, winds, hurricanes, and waves. The lingering effects of the oil spill and oil-dispersant mixtures residing in the ocean are currently unknown and subject to intensive research. However, the risk of damage to neighboring states and other countries is by no means over. The potential impacts on coastal communities and the geopolitical consequences of the spill could be extensive; more than 15 million people from three nations (11 states in the US and Mexico, combined, plus Cuba) live in the coastal areas bordering the Gulf of Mexico. Protection of these coastal ecosystems is therefore an international necessity. In particular, the marshes on the Alabama, Florida, Louisiana, and Mississippi coasts are in special need of attention because their proximity to the oil well increased their exposure and because large and economically important fisheries are located in these ecosystems.

■ Analysis of oil-spill mitigation efforts

The *Gulf Coast Restoration Plan*, pledged by President Obama on 16 June 2010 and subsequently expanded (The White House 2010), was aimed at mitigating the ecological, economic, and social damage due to the oil-spill's impacts, as well as recovering the thousands of square kilometers of wetlands that have vanished over the past century due to poor watershed management, stream diversion, levee construction, hydrological alteration, and diminished sediment supply (Day *et al.* 2007). However, the details of how, or even if, this restoration is to be carried out are incomplete, and it is unclear how this effort relates to the long-term restoration plan already in place for Louisiana (ie the Coastal Protection and Restoration Authority's master plan; CPRA 2010).




Figure 1. Location of the proposed sand berms along the Louisiana coast. Adapted from Lavoie *et al.* (2010).

On 11 May 2010, in direct response to threats to coastal ecosystems, the State of Louisiana requested emergency authorization from the US Army Corps of Engineers to perform spill mitigation work on all barrier islands, from Grand Isle and areas west, and from Grand Terre Island eastward to Sandy Point, as well as around the Chandeleur Islands. This request had no overt links to the Coastal Protection and Restoration Authority's master plan, nor was it connected with President Obama's *Gulf Coast Restoration Plan*. Furthermore, the request was made without consulting the regional scientific community. Its stated purpose was to enhance the capability of the barrier islands to reduce the amount of oil reaching coastal marshes. The proposal included the following actions: (1) building a barrier sand berm on the seaward side of existing barrier islands, and (2) closing some inlets between barrier islands and reducing the cross section of some inlets by partially filling them with large rocks (Lavoie *et al.* 2010).

The premise was essentially twofold: (1) sand berms can function as a geomorphic obstruction to, and an ecological filter for, oil spills, thereby protecting coastal marsh and estuarine ecosystems; and (2) spill cleanup is easier on sandy (dredged) substrate than in marshes and wetlands. In spite of intense controversy surrounding this project, it was partially authorized on 27 May 2010 (*Deepwater Horizon Response* 2010). The US Army Corps of Engineers approved a permit for 46 miles of sand berms to be constructed on the Chandeleur Islands (on the eastern side of the Mississippi River's outlet to the Gulf) and from Scofield Island to Timbalier Island (on the western side) (Figure 1; USACE 2010). As of 22 November 2010, 12.5 miles of berms had been completed (Figure 2; USACE 2010). Subsequently, authorities from the state and coastal counties presented a plan to use rock



Figure 2. Building sand berms (with dredged sand) along the Louisiana coast.

barriers to close some inlets into Barataria Bay to prevent oil intrusion into the estuary. This permit request was denied (USACE 2010).

A similar situation occurred on Dauphin Island, Alabama, where berms were constructed in May 2010. During Hurricane Katrina, a large gap was created as the island was cut in half by the storm surge (Feagin and Williams 2008) and the “Katrina Cut” project was subsequently initiated to block a major inlet (Figure 3). The aim of the approximately \$13 million Katrina Cut project was to fill and armor the gap with a combination of geo-textile tubes and riprap (a barrier consisting of large rocks or lumps of concrete). Initially, the plan was to re-open the inlet once the oil risk abated, although this has since changed; the intention now is to leave the barrier in place.

Freshwater diversions were also used in an effort to mitigate the impacts of the spilled oil. In late April 2010, the State of Louisiana opened floodgates into the Barataria Bay and Breton Sound basins, the premise being that by increasing the quantity of water flowing down the basins, oil could be prevented from approaching coastal marshes and penetrating inland. The initiated flow rate was much greater than that recommended in long-term planning documents describing the purpose of these diversion structures, which was to maintain salinities appropriate for fish and oyster production and to allow sedimentation into submerging wetlands (CPRA 2010).

Decisions to construct sand berms, fill inlets, or divert freshwater as protective measures against encroaching oil should be based on scientific knowledge and experience, and should take into account long-term restoration and management plans. To the best of our knowledge, there are no documented examples in the peer-reviewed literature of the effectiveness of these methods in mitigating oil-spill impacts on coastal ecosystems. The role

played by natural barrier islands in reducing erosion rates of inland coastal margins has been extensively investigated in this region (eg Stone and McBride 1998); however, there is little evidence of the successful use of constructed sand berms for oil protection in the literature (Froede 2007). The absence of evidence to support or refute the effectiveness of these methods in mitigating coastal ecosystem oil impacts underscores the importance of involving the scientific community in decision making and the integration of any coastal oil-spill mitigation strategies into existing coastal restoration plans. Below, we outline scientific and technical considerations that bear directly on the *Deepwater Horizon* accident and the subsequent remediation efforts.

■ Mitigation strategies

Sand berms

The amount of sediment needed to build the proposed sand berms in Louisiana alone is substantial. Estimates show that building a 2-m-high, 130-km-long barrier requires 2 million cubic meters of sand (Bahr 2010). The Northern Gulf region no longer features a continuous chain of islands for a variety of reasons, including human perturbations in sediment supply, sea-level rise, subsidence, and periodic hurricane damage (eg Otvos and Carter 2008). In this context, any sedimentary modification will be relatively short-lived. There is also the question of where the sediment for the berms comes from. Offshore sandy sediment resources are limited in extent and volume (Kulp *et al.* 2005), and the transport of suitable sediment from farther afield (Finkl *et al.* 2006) can be prohibitively expensive in terms of both time and resources. Sediment is precious in this region and should be used strategically (Kulp *et al.* 2005; Finkl *et al.* 2006). Moreover, when planning any large-scale dredging operation (Figure 2), serious consideration must be given to the potential impacts on in situ benthic organisms and habitat, and the potential for increasing turbidity and its effects on pelagic flora and fauna (Wilber and Clarke 2001; Erftemeijer and Lewis 2006).

Inlet restrictions

Inlets, particularly their geographical positions, are critical to the hydrological and ecological function of bays and estuaries. Geomorphic and ecological changes are likely to occur in locations where sand berms or other oil-spill mitigation operations block or restrict inlets. Profound alterations to sediment and biological transport

routes are possible whenever local hydrology is manipulated (Chaibi and Sedrati 2009). The reduction of inlet volume decreases the exchange of water and sediments during tides (Kraus and Militello 1999). This, in turn, affects flow velocities, typically resulting in scouring along bank margins, while near-shore wave dynamics often change as well (Komar 1996). The closing of inlets can modify the salinity, oxygen level, and turbidity of back bays. Associated ecosystems often experience large fish kills or other radical shifts toward new ecological states, as physical and chemical parameters adjust to a decrease in mixing throughout the water column (Goodwin 1996). If inlet restriction is absolutely necessary, it should aim to minimize the impacts on local hydrology and ecosystem function.



Figure 3. Crews blockade the inlet with riprap at Katrina Cut on Dauphin Island, Alabama. This view is from the uninhabited west end of the island, looking across the inlet toward the heavily developed east end.

Long-term barrier dynamics

Barrier islands, inlets, and sand resources are dynamic features of regressing coastlines. They change shape and move in response to ocean currents, winds, hurricanes, and storm surges (Grasso *et al.* 2009). All barrier islands are subject to, and impacted by, sand supply and mean sea level, but the details of how they form differ. Barrier islands along the southeastern US coastline are formed by natural wave, tidal, and aeolian processes. In contrast, the barrier islands of the Mississippi Delta were formed by the deposition of silt-laden sands arriving from upstream, and have a relatively short life cycle as compared with coastal barrier islands elsewhere. Additionally, the other Gulf barrier islands differ from those in North Carolina, for example, by being more sand starved and by being exposed to higher rates of relative sea-level rise due to high local subsidence rates.

Construction of sand berms and inlet restrictions should be based on knowledge of consolidation loading to the underlying sediment (eg compaction of sediments relative to sea level), overwash processes, and other features associated with long-term barrier island migration (Rosati and Stone 2009) or broad-spatial-scale littoral processes (Kraus and Militello 1999). In this sense, berms will be easily eroded, overtopped, and assimilated into the littoral budget of the islands. For example, Sallenger *et al.* (2009) observed exceptionally large geomorphic changes in the Chandeleur Islands as a result of several storms. In the broader context of the loss of these islands, relatively small waves from Hurricane Alex (2010) breached and eroded 2-m-high berms in southwest Louisiana and the Chandeleur Islands. Although provi-

sion of sand to the adjacent islands could be a positive outcome of rapid erosion, this would still represent a relatively expensive, non-strategic use of sediment resources.

In contrast, wherever inlets are blocked or restricted, the sedimentary and biological responses will be long-lasting and far-reaching. Inlet restriction projects, while initially driven by a sense of urgency to prevent oil from reaching estuaries, may turn out to have the most detrimental effects on estuarine ecosystems. The current trends in coastal science and engineering research lead in the opposite direction, toward a measured and strategic use of human engineering in the context of overwhelmingly large, powerful, and long-term natural processes (Feagin *et al.* 2010).

Freshwater diversions

Earlier studies that tested the effectiveness of river diversions in removing nutrients and organic loads draining into adjacent streams and rivers concluded that such strategies are useful. Mitsch and Day (2006) found that the use of these large freshwater diversions (maximum discharge 150–200 m³ s⁻¹) was sufficient to remove from rivers and streams a substantial percentage (>60%) of the nitrate–nitrogen supplied to these fluvial systems by farmland runoff. At the same time, the diversions increased the supply of sediments and nutrients to coastal areas, where they are needed to rebuild eroding marshes.

Because these studies have shown that freshwater diversions can flush contaminants downstream and potentially away from the coast, the State of Louisiana opened diversion structures in the Barataria Bay and Breton Sound basins, in an attempt to prevent oil from

flowing into coastal marshes. The general consensus among researchers and fishermen has been that this action devastated oyster recruitment, and is projected to wreck the oyster harvest for approximately 3 years (Greater New Orleans Inc Regional Economic Alliance 2010). In terms of reducing oiling, there has been no scientific evidence to support or refute that the use of freshwater diversions was effective in the Gulf. As with any environmental manipulation, there is a need for formal and rigorous research to reduce uncertainties and to ensure that the benefits are likely to be greater than potentially negative side effects.

■ Conclusions

Coastal ecosystems are dynamic, which means there are inherent risks for their inhabitants; hurricanes, storm surges, tsunamis, and rainfall-induced flooding are natural hazards, some of which may increase in frequency and/or magnitude with climate change. Continued pressure to increase oil extraction from coastal zones may lead to further stress on coastal ecosystems and recurring contaminant spills, resulting in large-scale and temporally extended social and economic impacts. The coasts of the Gulf of Mexico have been exposed to two of the five largest oil spills in history, many minor oil spills, and ongoing natural leaks (Jernelöv 2010). This necessitates a carefully considered approach to any restoration efforts for the Gulf's ecosystems.

Sand-berm construction, inlet modification, and the use of freshwater diversions to reduce oil exposure are high-risk strategies with dubious short-term effectiveness and potentially long-term negative implications. They involve the expenditure of large amounts of finite material and fiscal resources. Responses to the *Deepwater Horizon* accident appear to have been largely politically motivated, with little input from the regional, national, or international scientific community. Why did this happen? Why did risky programs, based on little or no technical or scientific evidence, continue to move forward? Are we prepared to deal with the lasting side effects? Did these projects make any difference to oil-spill remediation? The answers to these questions are largely unknown.

Comprehensive coastal restoration plans, including the CPRA (2010), have been extensively developed, and there is broad support for them from scientists and stakeholders. The consideration and development of these restoration plans is in direct contrast to the poorly planned schemes aimed at limiting the impacts of the *Deepwater Horizon* oil spill. The scientific community is regularly portrayed as being internally conflicted on many environmental issues and consistently calling for more research; in this particular case, however, the existence of a comprehensive plan should have at least provided a secure platform for developing an effective, efficient, and engaged approach to ecological restoration. In this instance, the scientific community had delivered

clear, comprehensive recommendations, but these were not adequately taken into account.

Were there other alternatives? Another option may have been to simply do nothing in direct response to the spill, and instead to redouble commitment to the CPRA. Some oil-spill damage may have been inevitable, yet the responses to it (eg artificial sand berms) may, in the end, cause even greater long-term harm to coastal ecosystems. In trying to reduce harm, we may have introduced more and longer-range impacts.

The challenges involved in designing mitigation and restoration plans are extensive and complex, and need to balance environmental and socioeconomic benefits. Ultimately, to prevent future disasters like the *Deepwater Horizon* spill, we need to shift the burden of risk (prior to the approval of drilling permits) more directly to the oil companies by improving and enforcing regulations, and requiring adequate insurance or assurance bonds to indemnify the public (Costanza *et al.* 2010). It is of particular concern that a Mexican oil company has begun the exploration and drilling of deep-water oil/gas wells, even though there is no regulation of these activities and no research regarding their safety and/or possible impacts in case of an accident. In addition to exploring for new wells, it is time to invest in research, development of safety procedures, accident prevention, long-term monitoring, and environmental conservation and restoration. We need to ensure that the shock of the oil spill does not distract us from the development of a viable, long-term plan for the restoration of the Gulf Coast. Elements of such a plan are already in place and judicious implementation needs to begin.

Finally, a year after the oil spill began, the results of studies to assess whether these artificial modifications of the coast actually worked, or if their impacts were worse than the spill itself, are beginning to trickle in. It is time to develop methods, policies, and institutions to act on that knowledge.

■ Acknowledgements

This paper was generated from a workshop (26–29 May 2010) of the Coastal Barrier Island Network, a multidisciplinary group concerned with the preservation of barrier islands (www.coastalbarrierisland.org/), and the Restoration Institute at the University of Victoria, British Columbia, Canada. Support was provided by the National Science Foundation, Research Coordination Networks in Biological Sciences (NSF-RCN, grant numbers IOB 0607921 and DBI 0741928). All authors have contributed ideas and text to each of the sections, and have been involved in formulation of the manuscript, editing, and approval of the final draft.

■ References

Bahr L. 2010. Like sand through the hour glass, so are the dunes of our lies. <http://lacoastpost.com/blog/?p=24268>. Viewed 15 Jul 2010.

- Batker D, de la Torre I, Costanza R, *et al.* 2010. Gaining ground. Wetlands, hurricanes and the economy: the value of restoring the Mississippi River delta. Tacoma, WA: Earth Economics. www.earthconomics.org/FileLibrary/file/Reports/Louisiana/Earth_Economics_Report_on_the_Mississippi_River_Delta_compressed.pdf. Viewed 23 May 2011.
- Chaibi M and Sedrati M. 2009. Coastal erosion induced by human activities: the case of two embayed beaches on the Moroccan coast. *J Coastal Res* **56**: 1184–88.
- Costanza R, Batker D, Day J, *et al.* 2010. The perfect spill: solutions for averting the next *Deepwater Horizon*. *Solutions* **1**: 17–20. www.thesolutionsjournal.com/node/629. Viewed 23 May 2011.
- CPRA (Coastal Protection and Restoration Authority). 2010. Integrated ecosystem restoration and hurricane protection: Louisiana's comprehensive master plan for a sustainable coast. Baton Rouge, LA: CPRA.
- Day J, Boesch D, Clairain E, *et al.* 2007. Restoration of the Mississippi Delta: lessons from Hurricanes Katrina and Rita. *Science* **315**: 1679–84.
- Deepwater Horizon* Response. 2010. Admiral Allen approves one section of Louisiana barrier island project proposal as part of federal oil spill response. www.restorethegulf.gov/release/2010/05/27/admiral-allen-approves-one-section-louisiana-barrier-island-project-proposal-part. Viewed 24 Jun 2011.
- Erfteimeijer PLA and Lewis RRR. 2006. Environmental impacts of dredging on seagrasses: a review. *Mar Pollut Bull* **52**: 1553–72.
- Feagin RA and Williams AM. 2008. Sediment spatial patterns in a Hurricane Katrina overwash fan on Dauphin Island, Alabama, USA. *J Coastal Res* **24**: 1063–70.
- Feagin RA, Smith WK, Psuty NP, *et al.* 2010. Barrier islands: coupling anthropogenic stability with ecological sustainability. *J Coastal Res* **26**: 987–92.
- Finkl CW, Khalil SM, Andrews J, *et al.* 2006. Fluvial sand sources for barrier island restoration in Louisiana: geotechnical investigations in the Mississippi River. *J Coastal Res* **22**: 773–87.
- Froede CR. 2007. Elevated waves erode the western end of the recently completed sand berm on Dauphin Island, Alabama. *J Coastal Res* **23**: 1602–04.
- Goodwin P. 1996. Predicting the stability of tidal inlets for wetland and estuary management. *J Coastal Res* **23**: 83–101.
- Grasso F, Michallet H, Certain R, and Barthelemy E. 2009. Experimental flume simulation of sandbar dynamics. *J Coastal Res* **56**: 54–58.
- Greater New Orleans Inc Regional Economic Alliance. 2010. A study on the economic impact of the *Deepwater Horizon* oil spill. Part one – fisheries. www.uflib.ufl.edu/docs/Economic%20Impact%20Study,%20Part%201%20-%20Full%20Report.pdf. Viewed 24 May 2011.
- Jernelöv A. 2010. How to defend against future oil spills. *Nature* **466**: 182–83.
- Kerr RA. 2010. Gulf oil flow as fast as 60 000 barrels per day. <http://news.sciencemag.org/scienceinsider/2010/06/gulf-oil-flow-as-fast-as-60000-b.html>. Viewed 15 Jun 2010.
- Komar PD. 1996. Tidal-inlet processes and morphology related to the transport of sediments. *J Coastal Res* **23**: 23–45.
- Kraus NC and Militello A. 1999. Hydraulic study of multiple inlet system: East Matagorda Bay, Texas. *J Hydrol Eng* **125**: 224–32.
- Kulp M, Penland S, Williams J, *et al.* 2005. Geologic framework, evolution, and sediment resources for restoration of the Louisiana coastal zone. *J Coastal Res* **44**: 56–71.
- Lavoie D, Flocks JG, Kindinger JL, *et al.* 2010. Effects of building a sand barrier berm to mitigate the effects of the *Deepwater Horizon* oil spill on Louisiana marshes. Stennis Space Center, MS: US Geological Survey. Open-file report 2010–1108.
- Lehr B, Bristol S, and Possolo A. 2010. Oil budget calculator: *Deepwater Horizon*. A report to the National Incident Command. Durham, NH: Coastal Response Research Center, University of New Hampshire.
- Mitsch WJ and Day Jr JW. 2006. Restoration of wetlands in the Mississippi–Ohio–Missouri (MOM) river basin: experience and needed research. *Ecol Eng* **26**: 55–69.
- NOAA (National Oceanic and Atmospheric Administration). 2010. Incident news: Arabian Gulf spills. Silver Spring, MD: NOAA Office of Response and Restoration. www.incident-news.gov/incident/6786. Viewed 24 May 2011.
- Otvos EG and Carter GA. 2008. Hurricane degradation–barrier development cycles, northeastern Gulf of Mexico: landform evolution and island chain history. *J Coastal Res* **24**: 463–78.
- Rosati JD and Stone GW. 2009. Geomorphologic evolution of barrier islands along the northern US Gulf of Mexico and implications for engineering design in barrier restoration. *J Coastal Res* **25**: 8–22.
- Sallenger Jr AH, Wright CW, Howd P, *et al.* 2009. Extreme coastal changes on the Chandeleur Islands, Louisiana, during and after Hurricane Katrina. In: Lavoie D (Ed). Sand resources, regional geology, and coastal processes of the Chandeleur Islands coastal system – an evaluation of the Breton National Wildlife Refuge. Reston, VA: US Geological Survey. <http://pubs.usgs.gov/sir/2009/5252/downloads/SIR2009-5252.pdf>. Viewed 24 May 2011.
- Stone GW and McBride RA. 1998. Louisiana barrier islands and their importance in wetland protection: forecasting shoreline change and subsequent response to wave climate. *J Coastal Res* **14**: 900–15.
- The White House. 2010. Memorandum from the President on the long-term Gulf Coast restoration support plan. Washington, DC: The White House. www.whitehouse.gov/the-press-office/memorandum-president-long-term-gulf-coast-restoration-support-plan. Viewed 24 May 2011.
- USACE (US Army Corps of Engineers). 2010. Team New Orleans. *Deepwater Horizon* emergency permit request. Project MVN-2010-01066-ETT and Project MVN-2010-01271-EOO. www.mvn.usace.army.mil/pao/mvnoilspill.asp.
- Wilber DH and Clarke DG. 2001. Biological effects of suspended sediments: a review of suspended sediment impacts on fish and shellfish with relation to dredging activities in estuaries. *N Am J Fish Manage* **21**: 855–75.

⁴Department of Oceanography and Coastal Sciences, School of the Coast and Environment, Louisiana State University, Baton Rouge, LA; ⁵Institute for Sustainable Solutions, Portland State University, Portland, OR; ⁶Department of Natural Resources, School of Applied Sciences, Cranfield University, Bedfordshire, Cranfield, UK; ⁷School of Plant Biology, University of Western Australia, Crawley, Australia; ⁸Department of Geography, University of Victoria, Victoria, British Columbia, Canada; ⁹University of Victoria, Victoria, British Columbia, Canada; ¹⁰Oceanografía Física, Ensenada, Baja California, Mexico; ¹¹Red de Ambiente y Sustentabilidad, Instituto de Ecología, El Haya, Xalapa, Veracruz, México