

6-5-2012

A Genetic Study of SSV1, the Prototypical Fusellovirus

Eric Iverson

Portland State University

Kenneth M. Stedman

Portland State University

Let us know how access to this document benefits you.

Follow this and additional works at: https://pdxscholar.library.pdx.edu/bio_fac



Part of the [Biology Commons](#), and the [Genomics Commons](#)

Citation Details

Iverson E and Stedman K (2012) A genetic study of SSV1, the prototypical fusellovirus. *Front. Microbio.* 3:200. doi: 10.3389/fmicb.2012.00200

This Article is brought to you for free and open access. It has been accepted for inclusion in Biology Faculty Publications and Presentations by an authorized administrator of PDXScholar. For more information, please contact pdxscholar@pdx.edu.



A genetic study of SSV1, the prototypical fusellovirus

Eric Iverson and Kenneth Stedman*

Biology Department, Center for Life in Extreme Environments, Portland State University, Portland, OR, USA

Edited by:

Frank T. Robb, University of Maryland, USA

Reviewed by:

Brian P. Hedlund, University of Nevada Las Vegas, USA

Matthias Hess, Washington State University, USA

*Correspondence:

Kenneth Stedman, Biology Department, Center for Life in Extreme Environments, Portland State University, P.O. Box 751, Portland, OR 97207-0751, USA.
e-mail: kstedman@pdx.edu

Viruses of thermophilic Archaea are unique in both their structures and genomic sequences. The most widespread and arguably best studied are the lemon-shaped fuselloviruses. The spindle-shaped virus morphology is unique to Archaea but widespread therein. The best studied fusellovirus is SSV1 from Beppu, Japan, which infects *Sulfolobus solfataricus*. Very little is known about the function of the genes in the SSV1 genome. Recently we have developed genetic tools to analyze these genes. In this study, we have deleted three SSV1 open reading frames (ORFs) ranging from completely conserved to poorly conserved: VP2, d244, and b129. Deletion of the universally conserved ORF b129, which encodes a predicted transcriptional regulator, results in loss of infectivity. Deletion of the poorly conserved predicted DNA-binding protein gene VP2 yields viable virus that is indistinguishable from wild-type. Deletion of the well-conserved ORF d244 that encodes a predicted nuclease yields viable virus. However, infection of *S. solfataricus* with virus lacking ORF d244 dramatically retards host growth, compared to the wild-type virus.

Keywords: DNA binding, nuclease, transcription factor

INTRODUCTION

Viruses of Archaea are very poorly understood with only about 50 known archaeal viruses relative to the ca. 5000 characterized viruses of bacteria, plants, and animals (Pina et al., 2011). The best studied of archaeal viruses are those infecting the thermoacidophiles, with an unprecedented new seven virus families introduced in the last few years to accommodate the astonishing morphological and sequence diversity present in these viruses (Pina et al., 2011).

The *Sulfolobus* spindle-shaped viruses (SSVs) of the family Fuselloviridae were the first discovered and probably the best studied family of archaeal viruses. SSVs are found throughout the world in high temperature (>70°C) and acidic (pH < 4) environments where their hosts, *Sulfolobus solfataricus* and its close relatives thrive (Wiedenheft et al., 2004; Held and Whitaker, 2009). The type virus, SSV1, encodes a positively supercoiled, 15.5 kbp circular dsDNA genome (NC_001338.1) that is enclosed within a lemon or spindle-shaped capsid (Yeats et al., 1982; Martin et al., 1984; Nadal et al., 1986). The genome encodes 34 open reading frames (ORFs; Palm et al., 1991), most of which have no recognizable homologs apart from other Fuselloviridae. The only SSV1 gene with clear homology to proteins outside the Fuselloviridae is the viral integrase, encoded by ORF d355. The main structural proteins purified from virus particles are the major and minor capsid proteins VP1 and VP3 and the putative DNA packaging protein VP2 (Reiter et al., 1987a). More recently, mass spectrometric analysis of SSV1 virions revealed two additional proteins, the products of ORFs c792 and d244 (Menon et al., 2008; Figure 1).


In the absence of homologous sequences, three complementary approaches have been used to try and determine the function of the proteins encoded in the SSV1 genome; structural genomics, comparative genomics, and genetics. Atomic resolution structures have been obtained by C. Martin Lawrence and his group for

proteins encoded by SSV1 ORFs b129, f112, d63, e96, f93, and d244 or their homologs from other fuselloviruses. The products of ORFs b129, f112, and f93 resemble transcriptional regulators and d244 a novel nuclease (Lawrence et al., 2009; Menon et al., 2010). However, the function of these proteins in virus replication remains to be determined. Two of these ORFs, b129 and d244, are the targets of the current study.

In parallel, we and others have undertaken comparative genomic studies. Fifteen ORFs are completely conserved in 12 canonical SSV genomes (Stedman et al., 2003; Wiedenheft et al., 2004; Held and Whitaker, 2009; Redder et al., 2009; Stedman, unpublished; Figure 1). Most of the universally conserved genes are clustered in half of the genome with the notable exception of the VP2 gene, a target of this study. Conservation in the rest of the genome is lower. Nonetheless, there are very few completely unique genes in the SSV1 genome (Figure 1). It is highly probable that the conserved genes are required for virus function, but again this has not been confirmed.

We developed methods for gene disruption in order to determine the requirements for genes in the virus genome directly. About 10 years ago, we showed that four SSV1 ORFs did not tolerate insertion of the 3.2 kbp pBluescript plasmid and allow virus function. Twelve other SSV1 ORFs appeared, indirectly, to not tolerate insertion. However, two ORFs, e178 and e51, were able to tolerate insertion of the entire pBluescript plasmid (Stedman et al., 1999). This result allowed the development of viral shuttle vectors and the beginnings of *Sulfolobus* genetics (Jonuscheit et al., 2003). Insertion of the pBluescript plasmid and up to ca. 5 kbp of exogenous DNA in these ORFs does not appear to have a noticeable effect on virus function (Stedman et al., 1999; Jonuscheit et al., 2003; Clore and Stedman, 2006; Albers et al., 2006).

However, insertion of large DNA fragments into the SSV1 genome is not straightforward and the possible insertion locations



are limited. Therefore, Long Inverse PCR (LIPCR) using high-fidelity highly processive DNA polymerases (e.g., Phusion[®]) was developed to specifically change the SSV1 genome at single nucleotide resolution. LIPCR was used to delete precisely the SSV1 viral integrase gene. Surprisingly, this “integrase-less” SSV1 was functional (Clore and Stedman, 2006). However, consistent with its conservation, the virus lacking the integrase gene is at a competitive disadvantage relative to integrase-containing viruses (Clore and Stedman, 2006). All of the SSV1 ORFs that can be deleted or tolerate insertion without abrogating virus function are in the “early” transcript, T5, that is induced soon after UV-irradiation of SSV-infected cultures (Reiter et al., 1987b; Fröls et al., 2007).

Three ORFs in the SSV1 genome were targeted for gene disruption in this study. The VP2 gene ([NP_039802.1](#)) was chosen for disruption because it is only present in SSV1 and the very distantly related SSV6 (Held and Whitaker, 2009; Redder et al., 2009), and is in the middle of the most highly conserved part of fusellovirus genomes (**Figure 1**). VP2 has DNA-binding activity (Reiter et al., 1987a; Iverson and Stedman, unpublished) that is presumably required for DNA packaging. ORF b129 ([NP_039795.1](#)) was chosen because it is intolerant of insertional mutagenesis (Stedman et al., 1999), a high resolution structure is known (Lawrence et al., 2009) and the gene is completely conserved in all SSVs (**Figure 1**). Finally, ORF d244 ([NP_039781.1](#)) was chosen for gene disruption because a high-resolution structure of its homolog from SSVRH

is known (Menon et al., 2008) and it is conserved in most SSV genomes with the exception of SSVK1.

MATERIALS AND METHODS

CULTURE CONDITIONS

Sulfolobus solfataricus strains, **Table 1**, were grown aerobically at 76°C on plates or in liquid media containing yeast extract and sucrose as carbon and energy sources (YS Media), both as in Jonuscheit et al. (2003). *Escherichia coli* strains were grown in LB medium at 37°C as suggested by the manufacturer (Novagen).

PURIFICATION OF DNA

Plasmid DNA used for LIPCR was purified from *E. coli* using the alkaline lysis method of Birnboim and Doly (1979). Plasmid DNA used to transform *Sulfolobus* was purified using the GeneJet Plasmid Purification Kit (Fermentas) following the manufacturer's protocols. Total genomic DNA was isolated from *S. solfataricus* in late log phase growth (OD600 ~0.6) as in Stedman et al. (1999). Plasmid DNA was purified from a 50 mL culture of *S. solfataricus* transformed with SSV-Δd244 (late log, OD600 ~0.6) using the GeneJet plasmid purification kit (Fermentas) following the manufacturer's protocols. This DNA was retransformed into *E. coli* (Novagen), purified therefrom and analyzed by restriction endonuclease digestion with *EcoRI* (Fermentas).

CONSTRUCTION OF SSV1 DELETION MUTANTS

Deletion mutants were constructed from the pAJC97 shuttle vector using LIPCR (Clare and Stedman, 2006). Primers were designed to overlap with the start and stop codon of the ORF to be deleted to keep the deletion in frame. Initially primers were designed using the archaea genome browser¹. Primer melting temperatures were matched and then checked for potential primer dimer and secondary structure formation using online tools from IDT². **Table 2** contains a list of oligonucleotide sequences used. LIPCR was performed using Phusion[®] High-Fidelity DNA Polymerase (NEB/Finnzymes) at a final concentration of 0.005 U/μL. LIPCR cycling conditions as follows: initial denaturation at 98°C for 3 min; 35 cycles of 98°C for 15 s, annealing for 15 s, 72°C for 6 min, and a final extension at 72°C for 6 min. The annealing temperatures for deletion of VP2, ORF d244, and ORF b129 were 59, 53, and 66°C, respectively. DNA was precipitated directly from LIPCR reactions using sodium acetate at a final concentration of 0.3 M and 95% EtOH. This DNA was phosphorylated using T4 polynucleotide kinase according to the manufacturer's protocols (Fermentas). DNA was ligated overnight (~20 h) at 16°C using 5 Weiss units of T4 DNA ligase (Fermentas). Ligated DNA was transformed into NovaBlue Singles chemically competent *E. coli* following the manufacturer's protocol (Novagen). Plasmids were purified from single colonies and deletion constructs were identified by restriction endonuclease digestions. The deletion borders were confirmed by sequencing of the plasmids.

ELECTROPORATION OF SULFOLOBUS

Purified plasmid DNA was electroporated into *Sulfolobus* strain GΘ as in Schleper et al. (1992). Following electroporation (400Ω,

1.5 kV, 25 μF), cells were immediately resuspended in 1 mL of YS media at 75°C and incubated for 1 h at 75°C. The cells were then added to 50 mL of prewarmed YS media (75°C) and grown in liquid media as outlined below.

SCREEN FOR FUNCTIONAL INFECTIOUS VIRUS/HALO ASSAY

To confirm the presence of infectious virus, halo assays were performed in duplicate 48 and 72 h post-electroporation (Stedman et al., 2003). Uninfected *Sulfolobus* GΘ cells were diluted to an OD600 nm = ~0.3 and allowed to grow until the OD600 nm reached ~0.35 (about 2.5 h). Half of a milliliter of this uninfected culture was added to 5 mL YS media containing 0.2% wt/vol Gelrite[®] as a softlayer and poured onto prewarmed YS plates. Two microliters of supernatant from electroporated cultures was spotted onto the lawns and plates were incubated at 75°C for up to 3 days. A halo of host growth inhibition, typically observed 48–72 h after incubation, indicated the presence of an infectious virus (**Figure 2**).

GROWTH CURVES

Portions of halos of growth inhibition from infected *S. solfataricus* GΘ cells were removed from plates with a sterile pipette tip and inoculated into liquid YS media. The culture was grown to an OD600 nm of ~0.6. One milliliter of this culture was diluted in 100 mL YS media to an OD600 ~0.050. Cultures were placed in a shaking incubator at 75°C and the OD600 nm was measured




FIGURE 2 | Typical growth inhibition of *S. solfataricus* on plates due to infectious virus. Lawns of *S. solfataricus* strain GΘ were prepared as in Stedman et al. (2003). Two microliters of supernatant from cultures transformed with either (A) SSV-ΔVP2 or (B) SSV-Δd244 were placed on the lawns where indicated. Δ indicates where SSV-ΔVP2 was spotted, ΔD indicates where SSV-Δd244 was spotted. P indicates SSV-WT spotted as a positive control. T or Tx indicates 2 μL of 0.01% Triton X-100 spotted as a control for lawn growth.

¹<http://archaea.ucsc.edu>

²<http://www.idtdna.com>

every 24 h. After 96 h, 1 mL of culture was diluted into 100 mL fresh YS media and returned to 75°C. One milliliter of culture was removed 72 h after each dilution, cells removed by centrifugation (14000 rpm for 5 min in a microcentrifuge) and the supernatant was screened for virus using the halo assay above.

TRANSMISSION ELECTRON MICROSCOPY

Supernatant from infected cultures was collected by centrifugation at 14,000 rpm for 5 min in a microcentrifuge. Five microliters of supernatant was absorbed onto a 400 mesh carbon/formvar grid (Ted Pella) for 2 min and negatively stained with 2% uranyl acetate for 20 sec. Grids were viewed on a JEOL 100CX TEM operated at 100 keV and images captured with a Gatan imager.

RESULTS

SSV1 IS INFECTIOUS WITHOUT THE VP2 GENE

The VP2 protein was purified from SSV1 virus particles and reported to be a DNA-binding protein (Reiter et al., 1987a). Surprisingly, a gene for VP2 was not found in SSV2 (Stedman et al., 2003) or SSVRH or SSVK1 (Wiedenheft et al., 2004). Moreover, a homolog is not present in the *S. solfataricus* or *S. islandicus* genomes (She et al., 2001; Reno et al., 2009; Guo et al., 2011). However, a very distant relative of SSV1, SSV6, which also contains an atypical putative tail fiber protein, has a VP2 gene (Redder et al., 2009). Thus, it is not clear whether SSV1 can function without a VP2 gene.

Therefore, we made an in-frame deletion of the majority of the VP2 gene by LIPCR in the context of the pAJC97 SSV1 shuttle vector (Clare and Stedman, 2006), leaving the first four codons and the last four codons (including the stop codon) of the ORF intact (see **Table 1**). The putative promoter for the T9 “early” transcript was also left intact. The construct containing the deletion, pAJC97- Δ VP2, is hereafter referred to as SSV- Δ VP2.

To determine if the SSV- Δ VP2 was able to make infectious virus, the shuttle vector was electroporated into *S. solfataricus* strain G \ominus . Two days after electroporation, the supernatant from the transformed strains caused inhibition of growth of uninfected *S. solfataricus* strain G \ominus on plates (**Figure 2**) that was indistinguishable from growth inhibition caused by the virus containing the VP2 gene. Similar growth inhibition was also observed on lawns of uninfected *S. solfataricus* strain S443, a new *S. solfataricus* isolate from Lassen Volcanic National Park that is a host

for all tested SSVs (Ceballos et al., in preparation). Moreover, the supernatant contained SSV-like particles when observed by transmission electron microscopy (**Figure 3**).

Infection by wild-type SSV1 and shuttle vectors does not drastically slow growth of cells in liquid culture for unknown reasons (Martin et al., 1984; Schleper et al., 1992; Stedman et al., 1999). The same is true of SSV- Δ VP2 (**Figure 4**). Infection with SSV- Δ VP2 was confirmed via PCR amplification (data not shown)

SSV1 CONSTRUCTS LACKING THE CONSERVED ORF b129 DO NOT APPEAR TO MAKE INFECTIVE VIRUSES

The b129 ORF in SSV1 is universally conserved in all fuselloviruses (Redder et al., 2009). Moreover shuttle vectors with pBluescript inserted into ORF b129 did not produce infective virus when electroporated into *Sulfolobus* (Stedman et al., 1999). However, a similar insertion mutant in the equally conserved SSV1 viral integrase appears to be non-functional (Stedman et al., 1999), but an in-frame deletion was functional (Clare and Stedman, 2006). A structure for the b129 ORF is also known (Lawrence et al., 2009) and it contains two Zn-finger putative DNA-binding motifs.

The b129 ORF was deleted with LIPCR. The deletion of the b129 ORF left the first four and last two codons of the ORF intact and maintained the predicted T3 promoter (Reiter et al., 1987b). This construct is referred to as SSV- Δ b129. Unlike the SSV- Δ VP2 construct, supernatants from *Sulfolobus* cells electroporated with SSV1- Δ b129 did not cause zones of growth inhibition when spotted on lawns of uninfected *S. solfataricus* strain G \ominus . A total of nine independent transformations were performed in which the wild-type virus consistently caused growth inhibition but SSV- Δ b129 did not. Moreover, no halos of growth inhibition were formed on lawns of *S. solfataricus* strain S443. It is not currently known at which step of virus replication the SSV- Δ b129 is deficient.

SSV1 LACKING ORF d244 IS INFECTIOUS BUT HAS A NOVEL PHENOTYPE

SSV1 ORF d244 is in the UV-inducible transcript T5, upstream of the viral integrase gene (**Figure 1**). The entire pBluescript plasmid can be inserted into the ORF directly upstream of ORF d244 without abrogating SSV1 function (Stedman et al., 1999). ORF d244 is well conserved in other Fusellovirus genomes with the exception of SSVK1 (Wiedenheft et al., 2004; Redder et al., 2009). The X-ray crystal structure of the homolog of SSV1 ORF d244, SSVRH ORF d212 has been solved and it is predicted to be a nuclease (Menon et al., 2010). Moreover, the product of ORF d244 has been reported to be in purified SSV1 particles (Menon et al., 2008).

The SSV1 d244 ORF was deleted with LIPCR. The deletion of the d244 ORF left the first two and last three codons of the ORF intact as well as maintained the ORF to avoid polar effects. This construct is referred to as SSV- Δ d244.

To determine if SSV- Δ d244 was able to make infectious virus, the shuttle vector was electroporated into *S. solfataricus* strain G \ominus . Two days after electroporation, the supernatant from the transformed strains caused inhibition of growth of uninfected *S. solfataricus* strain G \ominus on plates (**Figure 2**) and also inhibited growth of *S. solfataricus* strain S443 (data not shown).

Table 1 | Strains and plasmid vectors used in this work.

Strain/vector	Description	Reference
<i>S. solfataricus</i> G \ominus	MT4 Derivative	Cannio et al. (1998)
<i>S. solfataricus</i> S443	Novel <i>Sulfolobus</i> isolate	Unpublished data
<i>E. coli</i> NovaBlue [®]	Expression strain	Novagen, Inc.
pAJC97	SSV1 with TOPO PCR Blunt II	Clare and Stedman (2006)
pAJC97- Δ VP2	pAJC97 lacking VP2 gene	This Work
pAJC97- Δ d244	pAJC97 lacking ORF d244	This Work
pAJC97- Δ b129	pAJC97 lacking ORF b129	This Work

Table 2 | Oligonucleotides used in this work.

Name	Sequence	Description
VP2 LIPCR F	5'-CAC CGC AAG TAG GCC-3'	Flanks VP2 gene for deletion
VP2 LIPCR R	5'-CAC CCA CTT CAT ATC ACT CC-3'	Flanks VP2 gene for deletion
d244 LIPCR F	5'-ATC CAT TTA CCA TAA TCC ACC-3'	Flanks ORF d244 for deletion
d244 LIPCR R	5'-GGA AAA TGA TAT TCA ACT CAG AGG-3'	Flanks ORF d244 for deletion
b129 LIPCR F	5'-AGT TAG GCT CTT TTT AAA GTC TAC C-3'	Flanks ORF b129 for deletion
b129 LIPCR R	5'-TGA CTC CGT CAT CCT CTA AC-3'	Flanks ORF b129 for deletion
VP2 Check F	5'-ATT CAG ATT CTG WAT WCA GAA C-3'	Amplifies VP2 gene and flanking sequences
VP2 Check R	5'-TCS CCT AAC GCA CTC ATC-3'	Amplifies VP2 gene and flanking sequences
d244 Check F	5'-GGA ACT CCT CTC ATT AAC C-3'	Amplifies ORF d244 and flanking sequences
d244 Check R	5'-GAT CAT CAA CGA GTA TAT TGA CC-3'	Amplifies ORF d244 and flanking sequences
b129 Check F	5'-ATG AAG GCT GAG GAA ACA ATC GTG-3'	Amplifies ORF b129 and flanking sequences
b129 Check R	5'-TTA ATA TAG CTG CGA TGC AGT ATA GTT TAT TTG TGC-3'	Amplifies ORF b129 and flanking sequences

*Underlined sequence indicates ORF.




FIGURE 3 | Transmission electron micrographs of SSV particles. Supernatants from cultures of *S. solfataricus* strain G \ominus transformed with (A) pAJC97, (B) SSV- Δ VP2, (C) SSV- Δ d244, were negatively

stained with uranyl acetate and observed with a JEOL 100CX transmission electron microscope. Bar represents 0.2 μ m (B) or 0.5 μ m (A,C).

The supernatant contained SSV-like particles when observed by transmission electron microscopy (Figure 3).

Infection by wild-type SSV1, shuttle vectors and SSV- Δ VP2 does not slow growth of cells in liquid culture (Martin et al., 1984; Schleper et al., 1992; Stedman et al., 1999; see above). However, infection by SSV- Δ d244 drastically slows growth of *S. solfataricus* strains G \ominus and S443 in liquid culture (Figure 4). Infection with SSV- Δ d244 was confirmed via PCR. Moreover, restriction endonuclease digestion of viral DNA recovered from transformed *S. solfataricus* cells and retransformed into *E. coli* revealed no obvious alterations of the SSV- Δ d244 construct (data not shown).

DISCUSSION

THE PUTATIVE DNA PACKAGING PROTEIN VP2 IS NOT REQUIRED FOR SSV1 FUNCTION

The deletion of VP2 from SSV1 results in a functional virus that is indistinguishable from the wild-type virus (Figures 2–4). Based on the lack of conservation of VP2 this result is not completely

unexpected. However, almost all viruses contain a genome packaging protein. There is no clear sequence homolog of VP2 in the host genome, but there are a number of small DNA-binding proteins, such as Sso7d or Cren7 that may be able to functionally substitute for VP2 in SSV1 genome packaging (Choli et al., 1988; Guo et al., 2008). This will be tested with mass spectrometry of SSV- Δ VP2 particles. Alternatively, the VP2 protein may be involved in maintenance of the positive supercoiling of the SSV1 viral genome (Nadal et al., 1986). It would be interesting to know if the topology of the viral DNA is affected by the absence of VP2. It is predicted that positive supercoiling should increase the thermal stability of the DNA, so SSV- Δ VP2 may be less thermally stable than the wild-type virus.

The VP2 gene may be more prevalent than previously thought. VP2-like sequences have been reported from metagenomic studies, one in an acid mine drainage metagenome (Andersson and Banfield, 2008) and the other from Boiling Springs Lake in California (Diemer and Stedman, unpublished). These VP2 genes may be in the context of a SSV6 or ASV-like genome (Redder et al., 2009).




FIGURE 4 | Typical growth inhibition in liquid culture of virus constructs. Cultures of *S. solfataricus* G Θ infected with wild-type SSV1, diamonds, SSV- Δ VP2, triangles and SSV- Δ d244, squares, were diluted in YS media to equal starting OD600 nm and incubated at 75°C. At the indicated times, samples were removed and the OD600 nm was determined and the presence of virus was confirmed in each culture via halo assay. After 96 h, 1 mL of cells were diluted 1:100 in fresh YS media and returned to 75°C.

THE PRODUCT OF ORF b129 APPEARS TO BE ESSENTIAL FOR SSV1 INFECTIVITY

Homologs of SSV1 ORF b129 are present in all known SSVs (Redder et al., 2009). The b129 ORF also does not tolerate insertion of the pBluescript plasmid (Stedman et al., 1999). Thus, it is not surprising that deletion of ORF b129 leads to an incompletely replicating virus. However, the SSV1 integrase, a gene also conserved in all fuselloviruses, did not appear to tolerate insertion of pBluescript (Stedman et al., 1999), but could be deleted with LIPCR without abrogating virus function (Clare and Stedman, 2006). This indicates that either polar effects are important, which seems unlikely since the SSV1 integrase is at the end of the T5 transcript, or that insufficient replicate transformations were performed in the earlier study.

Nine replicate transformations of *S. solfataricus* with SSV- Δ b129 did not generate functional virus. However, we cannot absolutely determine that SSV1 ORF b129 is essential for virus function without complementation experiments, which are underway. The reasons for the apparent necessity of SSV1 ORF b129 are unclear, but the structure of the b129 ORF product, a predicted transcriptional regulator (Lawrence et al., 2009) and induction of the T6 transcript containing ORF b129 after UV-irradiation (Reiter et al., 1987b; Fröls et al., 2007) provides clues to its function.

The assay used herein for virus infection, ability to cause a zone of growth inhibition on a lawn of uninfected cells, is for virus spread and infectivity. There are many other aspects of virus replication that could be affected by disruption of ORF b129. An attractive hypothesis is that the b129 protein activates transcription of virus structural genes encoded by the “late” transcripts T7/8/9, T1, and T2 (Reiter et al., 1987b; Fröls et al., 2007; see **Figure 1**). This would be one of very few archaeal transcriptional activators characterized to date and the only the second archaeal viral transcriptional activator (Kessler et al., 2006). Thus, the SSV- Δ b129 construct may be able to replicate its genome, integrate into the host,

and have genome replication induced by UV-irradiation or some subset of these activities. Experiments to test these hypotheses are underway.

TRANSFECTION WITH SSV- Δ d244 PRODUCES VIRUS AND RETARDS HOST CELL GROWTH

The SSV1 d244 ORF is well-conserved in fuselloviruses with the exception of SSVK1 (Wiedenheft et al., 2004; Redder et al., 2009). However, SSV1 lacking ORF d244 clearly makes infectious virus particles (**Figures 2 and 3**). Moreover, the zones of clearing produced by supernatants of cells transfected with SSV- Δ d244 are clearer than those produced by either the wild-type or SSV- Δ VP2 viruses (**Figure 3**; unpublished data). They are reminiscent of zones of clearing produced by SSVK1 (data not shown). Unlike wild-type virus and SSV- Δ VP2, transfection by SSV- Δ d244 leads to drastically reduced host growth (**Figure 4**). The reasons for this growth inhibition are unclear. Similar growth phenotypes have been observed in SSVK1 infections (Stedman et al., in preparation). SSVK1 consistently produces more virus than similar cultures of the wild-type virus, so this may account for the growth defect (unpublished data). Whether SSV- Δ d244 consistently produces more virus than the wild-type or SSV- Δ VP2 is currently unknown.

The structure of the product of SSV1 ORF d244 is a predicted nuclease (Menon et al., 2010), similar to Holiday junction resolvase enzymes. Why the lack of a resolvase leads to slower host growth is unclear. Possibly SSV1 ORF d244 is involved in the specificity of SSV1 integration. SSV-K1 is known to integrate into multiple positions in the host genome (Wiedenheft et al., 2004), which may contribute to its higher copy number. Whether SSV- Δ d244 integrates into multiple positions in the host genome is under investigation. On the other hand, there may be a defect in SSV- Δ d244 replication or resolution of SSV replication intermediates that leads to accumulation of aberrant DNA, which, in turn, leads to slower host growth.

After multiple transfers of *Sulfolobus* cultures transfected with SSV- Δ d244 into fresh media, growth rates recover to near wild-type rates (unpublished data). The virus is still present in these cultures by PCR and is able to inhibit *Sulfolobus* growth on plates (unpublished data) so the virus is not lost or apparently rearranged (see Results). Whether there are other genetic changes in the virus or host under these conditions remains to be determined. One attractive possibility is changes to the CRISPR repeat structures that are proposed to be important for acquired immunity in *Sulfolobus* (Held and Whitaker, 2009).

SUMMARY AND OUTLOOK

Comparative and structural genomics has identified a number of targets for gene disruption in the SSV1 genome. Here precise gene disruptions of the poorly conserved VP2 gene, and the well-conserved ORFs b129 and d244 are described. Deletions in VP2 may allow insights into DNA packaging in the SSV1 genome. Deletion of ORF b129 may allow the identification of the second archaeal virus transcriptional activator. Deletion of ORF d244 may allow insight into copy number regulation in SSVs, previously thought to be regulated by ORF d63 (Lawrence et al., 2009). Clearly, there are many more genes to be analyzed in the SSV1

genome and more insights that can be gained by combining comparative genomics, structural biology, and genetics. In the future, biochemical work will be added to this suite of techniques to gain fundamental understanding of this fascinating, unique, and ubiquitous archaeal virus family.

REFERENCES

- Albers, S. V., Jonuscheit, M., Dinkelaker, S., Ulrich, T., Kletzin, A., Tampé, R., Driessen, A. J., and Schleper, C. (2006). Production of recombinant and tagged proteins in the hyperthermophilic archaeon *Sulfolobus solfataricus*. *Appl. Environ. Microbiol.* 72, 102–111.
- Anderson, A. F., and Banfield, J. F. (2008). Virus population dynamics and acquired virus resistance in natural microbial communities. *Science* 320, 1047–1050.
- Birnboim, H. C., and Doly, J. (1979). A rapid alkaline extraction procedure for screening recombinant plasmid DNA. *Nucleic Acids Res.* 7, 1513–1523.
- Cannio, R., Contursi, P., Rossi, M., and Bartolucci, S. (1998). An autonomously replicating transforming vector for *Sulfolobus solfataricus*. *J. Bacteriol.* 180, 3237–3240.
- Choli, T., Henning, P., Wittmann-Liebold, B., and Reinhardt, R. (1988). Isolation, characterization and microsequence analysis of a small basic methylated DNA-binding protein from the Archaeobacterium, *Sulfolobus solfataricus*. *Biochim. Biophys. Acta* 950, 193–203.
- Clore, A. J., and Stedman, K. M. (2006). The SSV1 viral integrase is not essential. *Virology* 361, 103–111.
- Fröls, S., Gordon, P. M., Panlilio, M. A., Schleper, C., and Sensen, C. W. (2007). Elucidating the transcription cycle of the UV-inducible hyperthermophilic archaeal virus SSV1 by DNA microarrays. *Virology* 365, 48–59.
- Guo, L., Brügger, K., Liu, C., Shah, S. A., Zheng, H., Zhu, Y., Wang, S., Lillestøl, R. K., Chen, L., Frank, J., Prangishvili, D., Paulin, L., She, Q., Huang, L., and Garrett, R. A. (2011). Genome analyses of Icelandic strains of *Sulfolobus islandicus* model organisms for genetic and virus-host interaction studies. *J. Bacteriol.* 193, 1672–1680.
- Guo, L., Feng, Y., Zhang, Z., Yao, H., Luo, Y., Wang, J., and Huang, L. (2008). Biochemical and structural characterization of Cren7, a novel chromatin protein conserved among crenarchaea. *Nucleic Acids Res.* 36, 1129–1137.
- Held, N. L., and Whitaker, R. J. (2009). Viral biogeography revealed by signatures in *Sulfolobus islandicus* genomes. *Environ. Microbiol.* 11, 457–466.
- Jonuscheit, M., Martusewitsch, E., Stedman, K. M., and Schleper, C. (2003). A reporter gene system for the hyperthermophilic archaeon *Sulfolobus solfataricus* based on a selectable and integrative shuttle vector. *Mol. Microbiol.* 48, 1241–1252.
- Kessler, A., Sezonov, G., Guijarro, J. I., Desnoves, N., Rose, T., Delepierre, M., Bell, S. D., and Prangishvili, D. (2006). A novel archaeal regulatory protein, Sta1, activates transcription from viral promoters. *Nucleic Acids Res.* 34, 4837–4845.
- Lawrence, C. M., Menon, S., Eilers, B. J., Bothner, B., Khayat, R., Douglas, T., and Young, M. J. (2009). Structural and functional studies of archaeal viruses. *J. Biol. Chem.* 284, 12599–12603.
- Martin, A., Yeats, S., Janekovic, D., Reiter, W. D., Aicher, W., and Zillig, W. (1984). SAV 1, a temperate u.v.-inducible DNA virus-like particle from the archaeobacterium *Sulfolobus acidocaldarius* isolate B12. *EMBO J.* 3, 2165–2168.
- Menon, S. K., Eilers, B. J., Young, M. J., and Lawrence, C. M. (2010). The crystal structure of D212 from *Sulfolobus* spindle-shaped virus ragged hills reveals a new member of the PD-(D/E)XK nuclease superfamily. *J. Virol.* 84, 5890–5897.
- Menon, S. K., Maaty, W. S., Corn, G. J., Kwok, S. C., Eilers, B. J., Kraft, P., Gillitzer, E., Young, M. J., Bothner, B., and Lawrence, C. M. (2008). Cysteine usage in *Sulfolobus* spindle-shaped virus 1 and extension to hyperthermophilic viruses in general. *Virology* 376, 270–278.
- Nadal, M., Mirambeau, G., Forterre, P., Reiter, W. D., and Duguet, M. (1986). Positively supercoiled DNA in a virus-like particle of an archaeobacterium. *Nature* 321, 256–258.
- Palm, P., Schleper, C., Grampp, B., Yeats, S., McWilliam, P., Reiter, W. D., and Zillig, W. (1991). Complete nucleotide sequence of the virus SSV1 of the archaeobacterium *Sulfolobus shibatae*. *Virology* 185, 242–250.
- Pina, M., Bize, A., Forterre, P., and Prangishvili, D. (2011). The archeoviruses. *FEMS Microbiol. Rev.* 35, 1035–1054.
- Redder, P., Peng, X., Brügger, K., Shah, S. A., Roesch, F., Greve, B., She, Q., Schleper, C., Forterre, P., Garrett, R. A., and Prangishvili, D. (2009). Four newly isolated fuselloviruses from extreme geothermal environments reveal unusual morphologies and a possible intervirion recombination mechanism. *Environ. Microbiol.* 11, 2849–2862.
- Reiter, W. D., Palm, P., Henschen, A., Lottspeich, F., and Zillig, W. D. (1987a). Identification and characterization of the genes encoding three structural proteins of the *Sulfolobus* virus-like particle SSV1. *Mol. Gen. Genet.* 206, 144–153.
- Reiter, W. D., Palm, P., Yeats, S., and Zillig, W. (1987b). Gene-expression in Archaeobacteria - Physical mapping of constitutive and UV-inducible transcripts from the *Sulfolobus* virus-like particle SSV1. *Mol. Gen. Genet.* 209, 270–275.
- Reno, M. L., Held, N. L., Fields, C. J., Burke, P. V., and Whitaker, R. J. (2009). Biogeography of the *Sulfolobus* pan-genome. *Proc. Natl. Acad. Sci. U.S.A.* 106, 8605–8610.
- Schleper, C., Kubo, K., and Zillig, W. (1992). The particle SSV1 from the extremely thermophilic archaeon *Sulfolobus* is a virus: Demonstration of infectivity and of transfection with viral DNA. *Proc. Natl. Acad. Sci. U.S.A.* 89, 7645–7649.
- She, Q., Singh, R. K., Confalonieri, F., Zivanovic, Y., Allard, G., Awayez, M. J., Chan-Weiher, C. C., Clausen, I. G., Curtis, B. A., De Moors, A., Erauso, G., Fletcher, C., Gordon, P. M., Heikamp-de Jong, I., Jeffries, A. C., Kozera, C. J., Medina, N., Peng, X., Thi-Ngoc, H. P., Redder, P., Schenk, M. E., Theriault, C., Tolstrup, N., Charlebois, R. L., Doolittle, W. F., Duguet, M., Gaasterland, T., Garrett, R. A., Ragan, M. A., Sensen, C. W., and Van der Oost, J. (2001). The complete genome of the crenarchaeon *Sulfolobus solfataricus* P2. *Proc. Natl. Acad. Sci. U.S.A.* 98, 7835–7840.
- Stedman, K. M., Schleper, C., Rumpf, E., and Zillig, W. (1999). Genetic requirements for the function of the archaeal virus SSV1 in *Sulfolobus solfataricus*: construction and testing of viral shuttle vectors. *Genetics* 152, 1397–1405.
- Stedman, K. M., She, Q., Phan, H., Arnold, H. P., Holz, I., Garrett, R. A., and Zillig, W. (2003). Relationships between fuselloviruses infecting the extremely thermophilic archaeon *Sulfolobus*: SSV1 and SSV2. *Res. Microbiol.* 154, 295–302.
- Wiedenheft, B., Stedman, K. M., Roberto, F., Willits, D., Gleske, A. K., Zoeller, L., Snyder, J., Douglas, T., and Young, M. (2004). Comparative genomic analysis of hyperthermophilic archaeal Fuselloviridae genomes. *J. Virol.* 78, 7438–7442.
- Yeats, S., McWilliam, P., and Zillig, W. (1982). A plasmid in the archaeobacterium *Sulfolobus acidocaldarius*. *EMBO J.* 1, 1035–1038.

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Received: 15 March 2012; paper pending published: 16 April 2012; accepted: 15 May 2012; published online: 05 June 2012.

Citation: Iverson E and Stedman K (2012) A genetic study of SSV1, the prototypical fusellovirus. *Front. Microbiol.* 3:200. doi: 10.3389/fmicb.2012.00200

This article was submitted to *Frontiers in Evolutionary and Genomic Microbiology*, a specialty of *Frontiers in Microbiology*. Copyright © 2012 Iverson and Stedman. This is an open-access article distributed under the terms of the Creative Commons Attribution Non Commercial License, which permits non-commercial use, distribution, and reproduction in other forums, provided the original authors and source are credited.