Portland State University

PDXScholar

University Honors Theses

University Honors College

2014

Effect of Mycorrhizal Colonization and Light Limitation on Growth and Reproduction of Lima Bean (*Phaseolus lunatus* L.)

Jess Annai Millar

Portland State University

Follow this and additional works at: <https://pdxscholar.library.pdx.edu/honorstheses>

Let us know how access to this document benefits you.

Recommended Citation

Millar, Jess Annai, "Effect of Mycorrhizal Colonization and Light Limitation on Growth and Reproduction of Lima Bean (*Phaseolus lunatus* L.)" (2014). *University Honors Theses*. Paper 62.

<https://doi.org/10.15760/honors.38>

This Thesis is brought to you for free and open access. It has been accepted for inclusion in University Honors Theses by an authorized administrator of PDXScholar. Please contact us if we can make this document more accessible: pdxscholar@pdx.edu.

Effect of Mycorrhizal Colonization and Light Limitation on Growth and
Reproduction of Lima Bean (*Phaseolus lunatus* L.)

by

Jess Annai Millar

An undergraduate honors thesis submitted in partial fulfillment of the
requirements for the degree of

Bachelor of Science

in

University Honors

and

Biology

Thesis Adviser

Daniel J. Ballhorn

Portland State University

2014

ABSTRACT

Plants can respond with sink stimulation of photosynthesis when colonized with fungal or bacterial root symbionts, compensating costs of carbohydrate allocation to the microbes. However, constraints may arise under light limitation when plants cannot extensively increase photosynthesis. We hypothesize that under such conditions the costs for maintaining the symbiosis outweigh the benefits, ultimately turning the mutualist microbes into parasites, resulting in reduced plant growth and reproduction.

Using lima bean (*Phaseolus lunatus*) as experimental plant, we applied two levels of light (full light, 75% shading) and microbial inoculation (sterile soil, mycorrhizal fungi) and quantified both vegetative and generative plant traits.

As expected, shaded plants produced less vegetative biomass and seeds than non-shaded plants. However, individual seeds were significantly heavier in shaded plants and required less time for germination. While under both light conditions mycorrhizal plants showed a significantly reduced belowground biomass, mycorrhizal fungi neither enhanced overall plants performance in terms of total biomass and seed production nor resulted in measurable costs in either light condition. Our study suggest that mycorrhizal colonization neither provided benefits to lima bean plants grown under full light, nor created costs when photosynthesis was limited.

ACKNOWLEDGEMENTS

I would like to thank my research advisor Daniel J. Ballhorn of the Biology Department at Portland State University. He has provided continual support but also generous freedom that has allowed me to design aspects and conduct much of this project independently. I would also like to acknowledge Daniel for his assistance in writing my thesis. His feedback has helped me become a better writer. I want to thank Tina Schroyer for her assistance with collecting greenhouse data and other lab duties. I also want to thank Tanya Cheeke for taking the time to teach me techniques for identifying and quantifying mycorrhizal fungi.

I would also like to thank my thesis committee: Michael Bartlett, Mitchell Cruzan, Kenneth Stedman, and Igor Lacan. I appreciate their time and effort reviewing my paper and supporting my research and education.

This work was supported by startup funds from PSU to Daniel J. Ballhorn. Fellowship and partial funding was provided by the Ronald E. McNair Scholars Program at PSU. Funding support was also provided by the Portland Bridges to Baccalaureate program at PSU and the Rose E. Tucker Trust Undergraduate Research Grant through the University Honors College at PSU.

TABLE OF CONTENTS

	PAGE
ABSTRACT.....	iv
ACKNOWLEDGMENTS	iii
TABLE OF CONTENTS.....	ii
LIST OF TABLES AND FIGURES.....	i
CHAPTERS	
I. INTRODUCTION	1
II. MATERIALS AND METHODS.....	5
III. RESULTS	8
IV. DISCUSSION	15
REFERENCES	21

LIST OF TABLES AND FIGURES

PAGE

TABLES

1.	Summary of ANOVA results of the effects of mycorrhizal fungi and light availability on vegetative plant traits	14
2.	Summary of ANOVA results of the effects of mycorrhizal fungi and light availability on reproductive plant traits	14
3.	Summary of ANOVA results of the effects of mycorrhizal fungi and light availability on seed traits and viability	14

FIGURES

1.	Effects of mycorrhizal fungi and light availability on vegetative plant and biomass traits	9
2.	Effects of mycorrhizal fungi and light availability on seed production.....	10
3.	Effects of mycorrhizal fungi and light availability on weight and viability of seeds.....	11
4.	Percent mycorrhizal colonization	12

CHAPTER I

INTRODUCTION

Mutualistic interactions between plants and microbes are an important component in the determination of plant diversity and ecosystem productivity. One of the most important groups of plant-associated microbial mutualists are mycorrhizal fungi (Smith and Read, 1997). The symbiosis between plants and mycorrhizal fungi is extremely widespread and ancient in the plant kingdom. Root colonization with mycorrhizal fungi occurs in >80% all plant species (Smith and Read, 1997) and has been observed in fossils dating back 400 million years ago (Remy and Taylor, 1994). Mycorrhizal fungi colonize the host plant's root, form extensive networks and participate in the acquisition of phosphorus (P) in case of arbuscular mycorrhizal fungi (AMF) and nitrogen in ectomycorrhizal fungi (EMF) (Smith and Read, 1997). Due to their critical impact on plant growth and species composition these microbial symbionts are considered keystone species in terrestrial ecosystems (Lodge et al., 1996).

Root colonization with mycorrhizal fungi generally has positive effects on plant growth (Chalk et al., 2006) and mycorrhiza inoculation is frequently applied to increase crop plant productivity in agricultural systems (Li et al., 2000, 2004; Ortas et al., 2003; Ortas, 2010). Positive effect of mycorrhiza on plants include increases in height (Hayman, 1986; Hoeksema et al., 2010; Safapour et al., 2011), biomass (Vejsadova et al., 1993; Mathur and Vyas, 2000; Ramana et al., 2010), shoot:root ratio (Gavito et al., 2000; Veresoglou et al., 2012), production of flowers (Dodd et al., 1983; Carey et al., 1992), and yield in crop plants such as *Phaseolus vulgaris*, *Glycine max*, and *Triticum aestivum* (Vejsadova et al., 1993; Bethlenfalvay et al., 1997; Abdel-Fattah, 1997; Li et al., 2005; Ramana et al., 2010; Safapour et al., 2011). There is an

extensive body of literature on the effects of mycorrhizal fungi in a broad range of plant families including legumes (Barea and Azcon-Aguilar, 1983; Yang et al., 1994; Olsen et al., 1999a; 1999b; Liu et al., 2003; Scheublin and Ridgway, 2004; Ortas, 2008; Muleta, 2010) but a detailed understanding of costs and benefits arising from the mycorrhizal symbiosis under different abiotic conditions is often lacking. Legumes are both important components in many terrestrial ecosystems and crop plants of world economic importance. Due to their association with another group of microbial symbionts—nitrogen-fixing rhizobia—legumes have been demonstrated to determine critically the productivity and species composition of ecosystems (Sprent and Sprent, 1990) and according to their key function in global nitrogen cycles legumes are considered ecosystem engineers (Carney and Matson, 2005; van der Heijden et al., 2008). In addition to their high impact on natural ecosystems, legumes are of high relevance in agro-ecosystems. Legumes critically enhance the sustainability of agroforestry systems (Muleta, 2010) or pastures (Haystead et al., 1988) and some legume crop plants are of world economic importance (*Glycine max*, *Pisum sativum*, *Phaseolus vulgaris*, *Phaseolus lunatus*).

Although in most cases clearly beneficial for the plants, the associations with mycorrhizal fungi also incur costs as the microbial symbionts may consume up to 16% of photosynthetically-fixed carbon, which otherwise could be allocated to growth and reproductive functions (Kaschuk et al., 2009). However, recent research demonstrated that plants can compensate this cost through sink stimulation of photosynthesis, which is thought to be an adaptation to take advantage of the nutrient supply provided without compromising the total amount of photosynthates available for plant growth and development (Mortimer et al., 2008). In natural ecosystems, light availability is often a variable resource due to competition among plant species and, depending on cultivation method, shows strong variation also in agricultural systems

(Chirko et al., 1996). While sink stimulation of photosynthesis is generally an efficient strategy to compensate for costs of carbohydrate allocation to microbial root symbionts, the question arises as of how plants respond to mycorrhizal inoculation under light limited conditions when photosynthesis cannot be increased easily.

The interacting effects of mycorrhizal colonization and light limitation on plant vegetative growth have been studied previously (Smith and Read, 1997), however, the impact of mycorrhiza on plant reproduction and the actual crop yield have received much less attention. One study found maize plants exposed to lower irradiance had a smaller percent mycorrhizal infection and smaller shoot weight, which was equated to yield (Daft and El-Giahmi, 1978). Another study showed that mycorrhizal infection did not affect *Pisum sativum* biomass, however increased light resulted in enhanced growth once the plant reached the flowering stage (Reinhard et al., 1994). Unfortunately these studies did not elucidate mycorrhizal fungi's effects on seed production. Analyzing the effects of mycorrhiza on actual seed set is of high importance due to two reasons: First, it is a much more precise fitness measure than plant biomass as mycorrhizal symbionts might lead to resource allocation and reduce plant biomass while increasing seed set. Second, for many agricultural plants, seed yield is of larger interest than vegetative parameters (Smith and Smith, 2011; Ronsheim, 2012). In the present study we used lima bean (Fabaceae: *Phaseolus lunatus*) as experimental plant. Lima bean represents an emerging model plant commonly used in studies on indirect and direct plant defense against herbivores (Ballhorn et al., 2008; 2009; 2010), but also bacterial (Yi et al., 2009) and fungal pathogens (Ballhorn et al., 2010). Furthermore, this plant is one of the most economically important *Phaseolus* species cultivated for food (Fofana et al., 1999; Alves et al., 2008; Bonifácio et al., 2012). To better understand the concerted effects of mycorrhizal colonization and light availability, we exposed

mycorrhizal and mycorrhiza-free lima bean plants to two levels of light each. We hypothesized that mycorrhizal fungi provide benefits for their host plant under full light regimes, but that reduced light availability might increase the costs of the symbiosis and shift a beneficial interaction to a detrimental one. To our knowledge, this study is the first to analyze the interactive effects of mycorrhizal fungi and light availability in lima bean in order to uncover potential costs of these microbes when photosynthesis is limited.

CHAPTER II

MATERIALS AND METHODS

Plants

Lima bean plants (*Phaseolus lunatus* cv. Henderson) were grown from seeds (American Meadows Inc., Williston, VT). Plants were cultivated in a greenhouse with light regime of 13:11 light:day. Light in the greenhouse was provided by a combination (1:1) of HQI-BT 400 W (Osram) and RNP-T/LR 400 W (Radium) lamps. Temperature was 27:19 °C (i.e. temperature of 27 °C in the light period and 19 °C in the dark period) and we maintained an air humidity of 70–80%. Plants were grown in containers (one plant per pot) with 12 cm in diameter in Sunshine Mix #1, LC1 (SunGro Horticulture®, Bellevue, WA) 175 g per pot and were watered daily.

Experimental Setup

In a full factorial split plot design with two whole blocks we applied four treatments, 15 replicates each (60 plants total) including two levels of light (full light, 75% shading) and two levels of mycorrhizal colonization (with mycorrhiza, mycorrhiza free). Light availability was measured at noon on a sunny day (+ additional lighting) and at table height was an average of 525 $\mu\text{mol photons s}^{-1} \text{ m}^{-2}$ in full light and an average of 138 $\mu\text{mol photons s}^{-1} \text{ m}^{-2}$ (LI-250 light meter; LI-COR Biosciences, Lincoln, Nebraska, USA) under the shade tent (205 cm $\times$ 113 cm $\times$ 113 cm). Experimental plants were inoculated with commercial mycorrhizal fungi [powder inoculant, Bio Organics™, La Pine, Oregon (*Glomus aggregatum*, *G. etunicatum*, *G. intraradices*, *G. mosseae*, *G. clarum*, *G. deserticola*, *G. monosporus*, *Gigaspora margarita*, and *Paraglomus brasilianum*), 10 cc (8 g) per plant], when they had developed completely unfolded

primary leaves. Plants were not inoculated with any other microbes (such as rhizobia) to avoid uncontrolled cross-effects between fungal and bacterial root symbionts.

Plant biomass and reproduction

Over the experimental period of 14 weeks we measured plant height, leaf number, and initial number of seed pods. At the end of the experiment we evaluated the above and belowground biomass, the final number of seed pods, and determined the shoot:root ratio. To obtain belowground biomass, plant root systems were carefully washed until all potting soil was removed. Above and belowground parts of plants were dried in an oven (IncuMax™ CV250 Convection Oven, Amerex Instruments, Inc., Lafayette, CA) at 70 °C for 5 days until constancy of weight. Seeds produced per plant were counted and weighed. To assess viability of seeds they were germinated by placing them between four 24 cm × 45 cm wet papers towels in the dark at 25 °C. Number of days to germination was recorded for each seed.

Microscopic analysis of mycorrhizal colonization

Mycorrhizal colonization was evaluated by taking 1 g fresh root samples, from each plant, from 4 separate locations of washed roots. Root segments were placed into histocassettes (VWR, West Chester, PA). All root samples were cleared with 10% KOH, acidified in 2% HCl, stained with 0.05% trypan blue solution, and preserved in lactoglycerol (Phillips and Hayman 1970). Roots were cut into 1 cm sections and at least 40 cm of roots from each plant were placed on a single microscope slide with lactoglycerol. Microscopic observations were conducted using an AmScope FM320 Trinocular Microscope in both 100x and 400x magnification. Roots were examined for mycorrhizal structures that intersected the microscope eyepiece crosshair at 100

random points using the Magnified Intersections Method (McGonigle et al. 1990). The presence or absence of mycorrhizal structures at 100 intersects was used to calculate percent root length colonization by mycorrhizal fungi per plant.

Data Analysis

The effects of 'Mycorrhizal inoculation' and 'Light availability' on plant traits were assessed with 2-way ANOVA split-plot analysis with 'Light Availability' as a blocking factor, using the GLM procedure in SAS. Square root transformation was performed on several plant growth parameters (number of leaves, buds, flowers, and seed pods) to control for normality in tests. 'Mycorrhizal inoculation' and 'Light availability' were set as fixed effects in all tests. Number of inflorescences per plant was set as a covariate in the 2-way ANOVA for number of seed pods at a defined time point (6 weeks after cultivation). All tests were performed in SAS version 9.2 (SAS, 2009).

CHAPTER III

RESULTS

Light availability

The colonization of experimental plants with mycorrhizal fungi was not affected by light availability (Table 1). Light availability significantly affected all considered vegetative plant traits including plant height ($F=144.59$, $P<0.001$), number of leaves ($F=38.93$, $P<0.001$), total biomass ($F=103.48$, $P<0.001$), above- ($F=70.10$, $P<0.001$) and belowground biomass ($F=290.16$, $P<0.001$), as well as shoot:root ratio ($F=4.99$, $P=0.034$; Table 1). Plants growing under full light were significantly taller, had more leaves, more total biomass, more above- and belowground biomass than plants cultivated under shaded conditions (Fig. 1).

Light availability also significantly affected all generative plants traits analyzed such as the initial number of pods per plant ($F=127.88$, $P<0.001$), the final number of pods ($F=34.28$, $P<0.001$), the number of pods dropped before maturation ($F=129.94$, $P<0.001$) as well as the number of seeds ($F=202.15$, $P<0.001$; Table 2). Plants growing under full sun had more initial pods, more final pods, and more seeds, but also dropped more pods than shaded plants (Fig. 2).

Seed quality parameters were significantly affected by light conditions including total seed weight ($F=43.56$, $P<0.001$), average seed weight ($F=10.72$, $P=0.002$), days to germination ($F=54.13$, $P<0.001$), but not the percentage of seeds that germinated (Table 3). Total and average seed weight were higher for shaded plants than for full light controls and seeds from plants under shaded conditions germinated significantly quicker than seeds from plants growing under full light conditions (Fig. 3).

Figure 1. Effects of mycorrhizal fungi and light availability on vegetative plant and biomass traits. Plant height (a) and number of leaves per plant (b) were determined at the end of the cultivation period. Total plant aboveground biomass (c), belowground biomass (d), total biomass (e) and plant shoot:root ratio (f) were determined on dry weight basis. MF = Mycorrhizal Fungi; Values shown are means + SE; n = 15 plants per treatment

Figure 2. Effects of mycorrhizal fungi and light availability on seed production. Number of initial pods per plant (a) was determined 6 weeks after planting. Final number of pods (b), dropped pods (c), and total seed production per plant (d) was determined at the end of the cultivation period. MF = Mycorrhizal Fungi; Values shown are means + SE; n = 15 plants per treatment

Figure 3. Effects of mycorrhizal fungi and light availability on weight and viability of seeds. Total weight of seeds per plant (a) and average seed weight (b) were determined on dry weight basis. Percent seeds germinated per plant (c) and days to seed germination per plant (d) were determined over an 8-day germination period. MF = Mycorrhizal Fungi; Values shown are means + SE; n = 15 plants per treatment for (a); for (b), (c), and (d), n = 27 seeds per treatment for shaded plants, n = 129 seeds for inoculated, no shade plants, and n = 147 seeds for control, no shade plants

AM fungal colonization

Microscopic analyses revealed successful mycorrhizal colonization in the inoculated group, whereas the control group showed little to no mycorrhizal fungi (Fig. 4). AM fungi influenced far fewer traits as light availability. Specifically, mycorrhizal fungi significantly affected

belowground biomass ($F=9.44$, $P=0.005$) and the shoot:root ratio of plants ($F=23.49$, $P<0.001$). Plants inoculated with mycorrhizal fungi had significantly lower belowground biomass than controls and accordingly a higher shoot:root ratio (Fig. 1). All other vegetative plant traits that we analyzed were not significantly changed in response to mycorrhizal colonization including plant height, number of leaves, aboveground biomass and total biomass (Table 1).

Figure 4. Percent mycorrhizal colonization. MF = Mycorrhizal Fungi; Values shown are means + SE; n = 15 plants per treatment

Reproductive plant traits, which were significantly affected by AM fungal colonization, were the number of initial pods ($F=7.38$, $P=0.011$) and the number of dropped pods ($F=12.14$, $P<0.001$). Plants with mycorrhizal fungi had less initial pods and dropped less pods than plants grown without the symbiotic partner (Fig. 2). However, the total number of final pods and total seed numbers did not show significant variation between mycorrhizal and mycorrhiza-free plants (Table 2).

Inoculation with mycorrhizal fungi significantly affected the average seed weight ($F=11.80$, $P=0.002$) as well as the days to germination ($F=5.27$, $P<0.022$). Inoculated plants had

significantly higher average seed weight per plant than controls, but seeds of these plants required more time to germinate (Fig. 3). In contrast, the total seed weight and the percent of seeds that germinated were not altered by mycorrhizal fungi (Table 3).

Interacting effects of ‘Light availability’ and ‘Mycorrhizal fungi’

The interaction of ‘Light availability’ and ‘Mycorrhizal fungi’ ($M \times L$) did not significantly affect any of the vegetative plant parameters assessed (Table 1). Of the reproductive plant traits analyzed, the number of initial pods ($F=5.37$, $P=0.028$) and the number of dropped pods ($F=10.28$, $P=0.003$) were significantly affected by the interaction of ‘Light availability’ and ‘Mycorrhizal fungi’, while the number of final pods and the number of seeds per pot were not affected (Table 2). The only seed trait to be significantly affected by the interacting term of both variables was the days to germination ($F=4.93$, $P=0.027$), while all other seed traits were not significantly altered (Table 3).

Table 1. Summary of ANOVA results of the effects of mycorrhizal fungi and light availability on vegetative plant traits.

Effects	Height			Total Leaves			Aboveground Biomass			Belowground Biomass			Total Biomass			Shoot:Root		
	df	F	P	F	P	F	F	P	F	F	P	F	F	P	F	F	P	
Mycorrhizae (M)	1	2.78	0.107	1.43	0.242	1.20	0.283		9.44	0.005	0.10	0.750	23.49	<0.001				
Light (L)	1	144.59	<0.001	38.93	<0.001	70.10	<0.001		290.16	<0.001	103.48	<0.001	4.99	0.034				
M × L	1	0.40	0.534	0.67	0.419	0.61	0.387		3.13	0.088	0.10	0.757	1.17	0.288				

Table 2. Summary of ANOVA results of the effects of mycorrhizal fungi and light availability on reproductive plant traits.

Effects	Initial Pods			Final Pods			Dropped Pods			Total Seeds		
	df	F	P	F	P	F	F	P	F	F	P	
Mycorrhizae (M)	1	7.38	0.011	0.05	0.821	12.14	0.002		1.14		0.294	
Light (L)	1	127.88	<0.001	34.28	<0.001	129.94	<0.001		202.15		<0.001	
M × L	1	5.37	0.028	1.54	0.225	10.28	0.003		1.14		0.294	

Table 3. Summary of ANOVA results of the effects of mycorrhizal fungi and light availability on seed traits and viability.

Effects	Total Seed Weight			Average Seed Weight			Percent Seed Germination			Days to Germination		
	df	F	P	F	P	F	F	P	F	F	P	
Mycorrhizae (M)	1	1.93	0.178	11.80	0.001	0.52	0.470	5.27	0.022			
Light (L)	1	543.56	<0.001	10.72	0.002	3.20	0.081	54.13	<0.001			
M × L	1	0.42	0.526	0.46	0.498	0.52	0.470	4.93	0.027			

Bolded *P*-values are statistically significant ($P < 0.05$)

CHAPTER IV

DISCUSSION

Even though the benefits of root-associated microbes are well studied in many cases, it remains elusive in what ways the outcome of this mutualism is affected by external abiotic conditions. In our study we quantitatively analyzed the effects of mycorrhizal fungi and light availability on growth and reproduction of lima bean. We hypothesized enhancing effects of growth and reproduction by mycorrhizal fungi under full light whereas we expected reduced plant development in mycorrhizal plants under shaded conditions due to constraints for plants to support both growth and the mycorrhizal partner when photosynthesis is limited. Studies in the past have shown decreases in plant growth in lower light conditions, mainly seen as decreased plant biomass (Daft and El-Giahmi, 1978; Bethlenfalvay and Pacovsky, 1983; Tester et al., 1986; Pearson et al., 1991). Thus, under such conditions, the costs for maintaining the mutualism with mycorrhizal fungi may outweigh the benefits, which ultimately turns the mutualistic microbes into parasites that exploit resources and reduce host fitness (Bethlenfalvay and Pacovsky, 1983; Reinhard et al., 1993). Contrary to our expectations, under full light, mycorrhizal fungi did not significantly enhance plant performance. Inoculated plants did not produce more biomass than the respective controls (Fig. 1), and mycorrhizal plants actually had significantly less pods than controls (Fig. 1). However, in our study the significantly negative effects of mycorrhizal fungi disappeared on level of actual number and weight of seeds, which showed no significant differences between the treatments (Figs. 2 & 3). These overall neutral effects of mycorrhizal fungi on plant reproductive structures observed in our study are in line with a recent meta-analysis looking at various legume species including *Cicer arietinum*, *Lens culinaris*, *Phaseolus*

vulgaris, *Pisum sativum*, and *Vicia faba*. In these plants, mycorrhizal fungi were found to have no effect on crop yield (Kaschuk et al., 2010). In our study the only factor affecting biomass and seed production was light availability. Under reduced light, plants produced less biomass and less seeds compared to plants growing under full light conditions.

There are several possible explanations for the limited impact of mycorrhizal fungi on plant growth and reproduction as observed in the present study. One possibility is that mycorrhizal fungi may play a less important role for grain legumes than for other plant species in general (Kaschuk et al., 2010) or that lima bean in particular is not adapted to form a symbiosis with mycorrhizal fungi. Even though lima bean represents an important crop plant, to the best of our knowledge there is no information available on the colonization of lima bean plants with mycorrhizal fungi under laboratory or field conditions. On the other hand, however, studies on a closely related plant species (snap bean, *Phaseolus vulgaris*) showed positive effects of mycorrhizal fungi on growth and nutrient uptake (Hacisalihoglu et al., 2005; Ciftci et al., 2010). Another possibility that could explain the limited impact of mycorrhizal fungi on lima bean observed in our study is that the commercial inoculum we used may not have contained fungal strains that provide a benefit to lima bean plants. Although we could show mycorrhizal colonization of the plants roots and formation of haustoria microscopically, this does not necessarily mean the fungi efficiently provided nutrients for their host plant. Furthermore, the rate of mycorrhizal colonization was relatively low compared to other plant species. However, this might be due to sampling roots towards the end of the plants' life cycle, which was required, as we needed to collect data on seed production of the same individual plants. At earlier stages of plant development, the rate of mycorrhizal colonization likely might have been higher as colonization rates decrease with plant age as it has been reported for *Hordeum vulgare*, *Secale*

cereale and *Triticum aestivum* (Dodd and Jeffries, 1986; Boswell et al., 1998; Li et al., 2005).

Furthermore, together with our microscopic evidence of successful colonization, the observation that mycorrhizal colonization of lima bean resulted in a significantly increased shoot:root ratio compared to the respective controls supports a good matching quality of plants and fungi rather than limited compatibility. A relatively lower root biomass in mycorrhizal plants compared to uncolonized conspecifics is a common phenomenon and indicates an efficient transport of minerals and water via mycorrhizal fungi (Kothari et al., 1990). Thus, the increased shoot:root ratio in mycorrhizal lima bean plants we observed here suggests an actual interaction between both partners.

In addition to effects of mycorrhizal fungi on above- and belowground biomass of the lima bean plants themselves we also considered effects of mycorrhizal inoculation on germination and viability of produced seeds, that is, on the next generation of host plants. Compared to plants growing under full light conditions, shaded plants produced overall significantly fewer but heavier seeds. Nevertheless, within each light treatment, mycorrhizal inoculation had no significant effect on the total seed weight produced per plant (Fig. 3a). Mycorrhiza, however, increased the average seed weight with heaviest seeds in shaded plants (Fig. 3b); in plants grown in full light and in shade, this increase was statistically significant. Increases in average seed weight due to mycorrhizal inoculation has been described previously for various plants such as *Triticum aestivum* and *Abutilon theophrasti* (Lu and Koide, 1994; Karagiannidis and Hadjisavva-Zinoviadi, 1998). While the underlying mechanisms are little understood, changes in resource allocation within the plant are likely as mycorrhiza fungi represent a significant carbon sink (Mathur and Vyas, 2000; Chalk et al., 2006; Kaschuk et al., 2009). Based on the results of our study we cannot make predictions whether the observed

mycorrhiza-mediated changes are positive or negative for plants in nature, however, other studies showed that changes in seed traits can have a far reaching impact on plant fitness. Seed weight and size have been identified as plant traits strongly influencing the dispersal, establishment, survival and growth of seedlings (Harper et al., 1970; Harper, 1977; Westoby, 1998; Weiher et al., 1999; Leishman et al., 2000; Moles and Westoby, 2004), particularly at early seedling stages (Leishman et al., 2000; Coomes and Grubb, 2003). Fenner and Thompson (2005) showed that large seeds had an increased probability of establishment under detrimental conditions. Generally seedlings developing from large seeds cope better than those of smaller seeded species under competition (Parrish and Bazzaz, 1985; Rees, 1995), drought, nutrient limitation (Lee and Fenner, 1989; Jurado and Westoby, 1992; Leishman and Westoby, 1994) and depth of seedling emergence (Gulmon and Url, 1992; Peterson and Facelli, 1992; Vázquez-Yanes and Orozco-Segovia, 1992). In line with our study, also deep shade has been identified as a factor selecting against small seed size (Grime and Jeffrey, 1965; Leishman and Westoby, 1994). Thus, development of larger seeds under shaded conditions and the increases in average seed weight due to mycorrhizal colonization as we observed in our study might represent fitness relevant parameters.

Beyond changes in seed weight, seeds produced by mycorrhiza-colonized and mycorrhiza-free plants showed differences regarding germination time (Table 3). While seeds from mycorrhizal inoculated parent plants took equal time to germinate compared to mycorrhiza-free controls when plants were grown in full light, seeds produced by mycorrhiza-colonized plants under shaded conditions germinated significantly later than seeds produced by mycorrhiza-free plants (Fig. 3). Whether these changes were due to variation in the thickness of seed coats determining water intake as the first step of the germination process or whether the

observed variation is due to different enzymatic activities in seeds derived from the different treatments remains elusive and, again, it is difficult to predict if these changes germination time increase or decrease plant fitness as this depends on the specific environmental conditions. In general, the benefits to shorter germination time in plants is variable, however in large-seeded plants shorter germination time has been demonstrated to enhance growth and fecundity (Verdú and Traveset, 2005).

Conclusions. Mycorrhizal fungi play a key role for plant performance and productivity in natural and agricultural ecosystems yet their effects on actual seed production in interdependence with variable abiotic conditions remains elusive in many cases. Thus, as mycorrhizal symbiosis holds great potential to improve sustainable crop production (Plenchette et al., 2005) there is an urgent need to functionally study this almost ubiquitous interaction and analyze the effects of environmental factors on the symbiosis (Ortas, 2012). Using lima bean (*Phaseolus lunatus*) as experimental plant, we hypothesized that for plants under light limitation the costs for maintaining the symbiosis outweigh the benefits of the fungal partner, ultimately turning the beneficial microbes into parasites, resulting in reduced plant growth and reproduction. Contrary to our expectations, we found that mycorrhizal colonization neither provided benefits in terms of increased biomass and total seed number and total seed weight to plants grown under full light, nor created costs under shaded conditions. Mycorrhizal colonization did, however, significantly increase the average seed weight in both light treatments with heaviest seeds in shaded plants and lengthened germination time of seeds produced by shaded plants. Our study shows that the effects of mycorrhizal symbionts go beyond mere effects on plant biomass production as they significantly alter plant reproductive traits. Results of our study suggest that even though mycorrhiza commonly enhances plant growth, prior to costly inoculation of agricultural systems

with commercial mycorrhizal strains, preceding experiments are required to test for actual effects of mycorrhiza on the specific crop plant.

REFERENCES

- Abdel-Fattah GM (1997) Functional activity of VA-mycorrhiza (*Glomus mosseae*) in the growth and productivity of soybean plants grown in sterilized soil. *Folia Microbiol* 42: 495–502. doi:10.1007/bf02826560.
- Alves AU, Oliveira AP De, Alves AU, Dornelas CS, Alves EU, et al. (2008) Lima beans production and economic revenue as function of organic and mineral fertilization. *Hortic Bras* 26: 251–254. doi:10.1590/s0102-05362008000200024.
- Ballhorn DJ, Kautz S, Heil M, Hegeman AD (2009) Cyanogenesis of wild lima bean (*Phaseolus lunatus* L.) is an efficient direct defense in nature. *PLoS ONE* 4: e5450. doi:10.1371/journal.pone.0005450.
- Ballhorn D, Pietrowski A, Lieberei R (2010) Direct trade-off between cyanogenesis and resistance to a fungal pathogen in lima bean (*Phaseolus lunatus* L.). *J Ecol* 98: 226–236. doi:10.1111/j.1365-2745.2009.01591.x.
- Ballhorn DJ, Schiwy S, Jensen M, Heil M (2008) Quantitative variability of direct chemical defense in primary and secondary leaves of lima bean (*Phaseolus lunatus*) and consequences for a natural herbivore. *J Chem Ecol* 34: 1298–1301. doi:10.1007/s10886-008-9540-1.
- Barea J, Azcon-Aguilar C (1983) Mycorrhizas and their significance in nodulating nitrogen-fixing plants. *Adv Agron* 36: 1–54. doi:10.1016/s0065-2113(08)60351-x.
- Bethlenfalvay GJ, Pacovsky RS (1983) Light effects in mycorrhizal soybeans. *Plant Physiol* 73: 969–972. doi:10.1104/pp.73.4.969.

- Bethlenfalvay GJ, Schreiner RP, Mihara KL (1997) Mycorrhizal fungi effects on nutrient composition and yield of soybean seeds. *J Plant Nutr* 20: 581–591. doi:10.1080/01904169709365276.
- Bonifácio EM, Fonsêca A, Almeida C, Dos Santos KGB, Pedrosa-Harand A (2012) Comparative cytogenetic mapping between the lima bean (*Phaseolus lunatus* L.) and the common bean (*P. vulgaris* L.). *Theor Appl Genet* 124: 1513–1520. doi:10.1007/s00122-012-1806-x.
- Boswell EP, Koide RT, Shumway DL, Addy HD (1998) Winter wheat cover cropping, VA mycorrhizal fungi and maize growth and yield. *Agr Ecosyst Environ* 67: 55–65. doi:10.1016/s0167-8809(97)00094-7.
- Carey PD, Fitter AH, Watkinson AR, Url S, Carey D (1992) A field study using the fungicide benomyl to investigate the effect of mycorrhizal fungi on plant fitness. *Oecologia* 90: 550–555. doi:10.1007/bf01875449.
- Carney KM, Matson PA (2005) Plant communities, soil microorganisms, and soil carbon cycling: does altering the world belowground matter to ecosystem functioning? *Ecosystems* 8: 928–940. doi:10.1007/s10021-005-0047-0.
- Chalk P, Souza R, Urquiaga S, Alves B, Boddey R (2006) The role of arbuscular mycorrhiza in legume symbiotic performance. *Soil Biol Biochem* 38: 2944–2951. doi:10.1016/j.soilbio.2006.05.005.
- Chirko CP, Gold MA, Nguyen P, Jiang JP (1996) Influence of direction and distance from trees on wheat yield and photosynthetic photon flux density (Qp) in a Paulownia and wheat intercropping system. *For Ecol Manage* 83: 171–180. doi:10.1016/0378-1127(96)03721-8.

- Ciftci V, Turkmen O, Erdinc C, Sensoy S (2010) Effects of different arbuscular mycorrhizal fungi (AMF) species on some bean (*Phaseolus vulgaris* L.) cultivars grown in salty conditions. *Afr J Agric Res* 5: 3408–3416. doi:10.5897/ajar10.655.
- Coomes DA, Grubb PJ (2003) Colonization, tolerance, competition and seed-size variation within functional groups. *Trends Ecol Evol* 18: 283–291. doi:10.1016/S0169-5347(03)00072-7.
- Daft M, El-Giahmi A (1978) Effect of arbuscular mycorrhiza on plant growth. VIII. Effects of defoliation and light on selected hosts. *New Phytol* 80: 365–372. doi:10.1111/j.1469-8137.1978.tb01570.x.
- Dodd J, Jeffries P (1986) Early development of vesicular-arbuscular mycorrhizas in autumn-sown cereals. *Soil Biol Biochem* 18: 149–154. doi:10.1016/0038-0717(86)90019-2.
- Dodd J, Krikun J, Haas J (1983) Relative effectiveness of indigenous populations of vesicular-arbuscular mycorrhizal fungi from four sites in the Negev. *Isr J Bot* 32: 10–21.
- Fenner M, Thompson K (2005) *The ecology of seeds*. 1st ed. Cambridge: Cambridge University Press.
- Fofana B, Baudoin JP, Vekemans X, Debouck DG, Du Jardin P (1999) Molecular evidence for an Andean origin and a secondary gene pool for the Lima bean (*Phaseolus lunatus* L.) using chloroplast DNA. *Theor Appl Genet* 98: 202–212. doi:10.1007/s001220051059.
- Gavito ME, Curtis PS, Mikkelsen TN, Jakobsen I (2000) Atmospheric CO₂ and mycorrhiza effects on biomass allocation and nutrient uptake of nodulated pea (*Pisum sativum* L.) plants. *J Exp Bot* 51: 1931–1938. doi:10.1093/jexbot/51.352.1931.
- Grime AJP, Jeffery DW (1965) Seedling establishment in vertical gradients of sunlight. *J Ecol* 53:621–642.

- Gulmon ASL, Uri S (1992) International association for ecology patterns of seed germination in Californian serpentine grassland species. *Oecologia* 89: 27–31.
- Hacisalihoglu G, Duke ER, Longo LM (2005) Differential response of common bean genotypes to mycorrhizal colonization. *Proc Fla State Hort Soc* 118: 150–152.
- Harper JL (1977) Population biology of plants. London, GB: Academic Press.
- Harper JL, Lovell PH, Moore KG (1970) The shapes and sizes of seeds. *Annu Rev Ecol Syst* 1: 327–356. doi:10.1146/annurev.es.01.110170.001551.
- Hayman DS (1986) Mycorrhizae of nitrogen-fixing legumes. *MIRCEN J Appl Microbiol Biotechnol* 2: 121–145. doi:10.1007/bf00937189.
- Haystead A, Malajczuk N, Grove T (1988) Underground transfer of nitrogen between pasture plants infected with vesicular-arbuscular mycorrhizal fungi. *New Phytol* 108: 417–423. doi:10.1111/j.1469-8137.1988.tb04182.x.
- Hoeksema JD, Chaudhary VB, Gehring CA, Johnson NC, Karst J, et al. (2010) A meta-analysis of context-dependency in plant response to inoculation with mycorrhizal fungi. *Ecol Lett* 13: 394–407. doi:10.1111/j.1461-0248.2009.01430.x.
- Jurado E, Westoby M (1992) Seedling growth in relation to seed size among species of arid Australia. *J Ecol* 80: 407–416. doi:10.2307/2260686.
- Karagiannidis N, Hadjisavva-Zinoviadi S (1998) The mycorrhizal fungus *Glomus mosseae* enhances growth, yield and chemical composition of a durum wheat variety in 10 different soils. *Nutr Cycl Agroecosys* 52: 1–7. doi:10.1023/a:1016311118034.

- Kaschuk G, Kuyper TW, Leffelaar PA, Hungria M, Giller KE (2009) Are the rates of photosynthesis stimulated by the carbon sink strength of rhizobial and arbuscular mycorrhizal symbioses? *Soil Biol Biochem* 41: 1233–1244. doi:10.1016/j.soilbio.2009.03.005.
- Kaschuk G, Leffelaar PA, Giller KE, Alberton O, Hungria M, et al. (2010) Responses of legumes to rhizobia and arbuscular mycorrhizal fungi: A meta-analysis of potential photosynthate limitation of symbioses. *Soil Biol Biochem* 42: 125–127. doi:10.1016/j.soilbio.2009.10.017.
- Kothari SK, Marschner H, George E (1990) Effect of VA mycorrhizal fungi and rhizosphere microorganisms on root and shoot morphology, growth and water relations in maize. *New Phytol* 116: 303–311. doi:10.1111/j.1469-8137.1990.tb04718.x.
- Lee WG, Fenner M (1989) Mineral nutrient allocation in seeds and shoots of twelve *Chionochloa* species in relation to soil fertility. *J Ecol* 77: 704–716.
- Leishman MR, Westoby M (1994) The role of large seed size in shaded conditions: experimental evidence. *Funct Ecol* 8: 205–214. doi:10.2307/2389903.
- Leishman MR, Wright IJ, Moles, AT, Westoby M (2000) The evolutionary ecology of seed size. In: Fenner M, editor. *Seeds: the ecology of regeneration in plant communities*. 2nd ed. Wallingford, UK: CAB International. pp. 31–58.
- Li HY, Zhu YG, Marschner P, Smith FA, Smith SE (2005) Wheat responses to arbuscular mycorrhizal fungi in a highly calcareous soil differ from those of clover, and change with plant development and P supply. *Plant Soil* 277: 221–232. doi:10.1007/s11104-005-7082-7.

- Li M, Meng XX, Jiang JQ, Liu RJ (2000) A preliminary study on relationship between arbuscular mycorrhizal fungi and *Fusarium* wilt of watermelon (in Chinese). *Acta Phytotaxon Sin* 30: 327–331.
- Li M, Liu RJ, Li XL (2004) Influence of arbuscular mycorrhizal fungi on growth and *Fusarium*-wilt disease of watermelon grown in the field (in Chinese). *Acta Phytotaxon Sin* 34: 456–457.
- Liu A, Hamel C, Elmi AA, Zhang T, Smith DL (2003) Reduction of the available phosphorus pool in field soils growing maize genotypes with extensive mycorrhizal development. *Can J Plant Sci* 83: 737–744. doi:10.4141/P02-199.
- Lodge DJ, Hawksworth DL, Ritchie BJ (1996) Microbial diversity and tropical forest functioning. In: Oriens GH, Dirzo R, Cushman JH, editors. *Biodiversity and ecosystem processes in tropical forests*. Berlin: Springer-Verlag, Vol. 122. pp. 69–100.
- Lu X, Koide RT (1994) The effects of mycorrhizal infection on components of plant growth and reproduction. *New Phytol* 128: 211–218. doi:10.1111/j.1469-8137.1994.tb04004.x.
- Mathur N, Vyas A (2000) Influence of arbuscular mycorrhizae on biomass production, nutrient uptake and physiological changes in *Ziziphus mauritiana* Lam. under water stress. *J Arid Environ* 45: 191–195. doi:10.1006/jare.2000.0644.
- McGonigle TP, Miller MH, Evans DG, Fairchild GL, Swan JA (1990) A new method which gives an objective measure of colonization of roots by vesicular-arbuscular mycorrhizal fungi. *New Phytol* 115: 495–501. doi:10.1111/j.1469-8137.1990.tb00476.x.
- Moles AT, Westoby M (2004) Seedling survival and seed size: a synthesis of the literature. *J Ecol* 92: 372–383. doi:10.1111/j.0022-0477.2004.00884.x.

- Mortimer PE, Pérez-Fernández MA, Valentine AJ (2008) The role of arbuscular mycorrhizal colonization in the carbon and nutrient economy of the tripartite symbiosis with nodulated *Phaseolus vulgaris*. *Soil Biol Biochem* 40: 1019–1027. doi:10.1016/j.soilbio.2007.11.014.
- Muleta D (2010) Legume responses to arbuscular mycorrhizal fungi inoculation in sustainable agriculture. In: Khan MS, Musarrat J, Zaidi A, editors. *Microbes for legume improvement*. Vienna: Springer Vienna. pp. 293–323. doi:10.1007/978-3-211-99753-6_12.
- Olsen JK, Schaefer JT, Edwards DG, Hunter MN, Galea VJ, Muller LM (1999a) Effects of a network of mycorrhizae on capsicum (*Capsicum annuum* L.) grown in the field with five rates of applied phosphorus. *Aust J Agric Res* 50: 239–252. doi:10.1071/A98008.
- Olsen JK, Schaefer JT, Edwards DG, Hunter MN, Galea VJ, Muller LM (1999b) Effects of mycorrhizae, established from an existing intact hyphal network, on the growth response of capsicum (*Capsicum annuum* L.) and tomato (*Lycopersicon esculentum* Mill.) to five rates of applied phosphorus. *Aust J Agric Res* 50: 223–237. doi:10.1071/A97167.
- Ortas I (2010) Effect of mycorrhiza application on plant growth and nutrient uptake in cucumber production under field conditions. *Span J Agric Res* 8: S116–S122. doi:10.5424/sjar/201008S1-1230.
- Ortas I (2012) The effect of mycorrhizal fungal inoculation on plant yield, nutrient uptake and inoculation effectiveness under long-term field conditions. *Field Crop Res* 125: 35–48. doi:10.1016/j.fcr.2011.08.005.

- Ortas I, Sari N, Akpınar C (2003) Effects of mycorrhizal inoculation and soil fumigation on the yield and nutrient uptake of some solanaceas crops (tomato, eggplant and pepper) under field conditions. *Agric Mediterr* 133: 249–258.
- Ortas İ (2008) The effect of mycorrhizal inoculation on forage and non-forage plant growth and nutrient uptake under field conditions. In: *Options méditerranéennes.sustainable mediterranean grasslands and their multi-functions*. Zaragoza, Spain: Ciheam. pp. 463–469.
- Parrish J, Bazzaz FA (1985) Nutrient content of *Abutilon theophrasti* and the competitive ability of the resulting plants. *Oecologia* 65: 247–251. doi:10.1007/BF00379224.
- Pearson J, Smith S, Smith F (1991) Effect of photon irradiance on the development and activity of VA mycorrhizal infection in *Allium porrum*. *Mycol Res* 95: 741–746. doi:10.1016/s0953-7562(09)80824-1.
- Phillips J, Hayman D (1970) Improved procedures for clearing roots and staining parasitic and vesicular-arbuscular mycorrhizal fungi for rapid assessment of infection. *Trans Br Mycol Soc* 55: 158–161. doi:10.1016/s0007-1536(70)80110-3.
- Plenchette C, Clermont-Dauphin C, Meynard JM, Fortin JA (2005) Managing arbuscular mycorrhizal fungi in cropping systems. *Can J Plant Sci* 85: 31–40. doi:10.4141/P03-159.
- Ramana V, Ramakrishna M, Purushotham K, Reddy KB (2010) Effect of bio-fertilizers on growth, yield attributes and yield of french bean (*Phaseolus vulgaris* L.). *Legume Res* 33: 178–183.
- Rees M (1995) Community structure in sand dune annuals: is seed weight a key quantity? *Oecologia* 83: 857–863.

- Reinhard S, Martin P, Marschner H (1993) Interactions in the tripartite symbiosis of pea (*Pisum sativum* L.), *Glomus* and *Rhizobium* under non-limiting phosphorus supply. J Plant Physiol 141: 7–11. doi:10.1016/S0176-1617(11)80844-8.
- Reinhard S, Weber E, Martin P, Marschner H (1994) Influence of phosphorus supply and light intensity on mycorrhizal response in *Pisum-Rhizobium-Glomus* symbiosis. Experientia 50: 890–896.
- Remy W, Taylor T (1994) Four hundred-million-year-old vesicular arbuscular mycorrhizae. Proc Natl Acad Sci USA 91: 11841–11843. doi:10.1073/pnas.91.25.11841.
- Ronsheim M (2012) The effect of mycorrhizae on plant growth and reproduction varies with soil phosphorus and developmental stage. Am Midl Nat 167: 28–39. doi:10.1674/0003-0031-167.1.28.
- Safapour M, Ardakani M, Khaghani S, Rejali F, Zargari K, et al. (2011) Response of yield and yield components of three red bean (*Phaseolus vulgaris* L.) genotypes to co-inoculation with *Glomus intraradices* and *Rhizobium phaseoli*. Am Eurasian J Agric Environ Sci 11: 398–405.
- SAS Institute Inc (2009) Base SAS® 9.2 procedures guide. 1st ed. Cary, NC: SAS Institute Inc.
- Scheublin T, Ridgway K (2004) Nonlegumes, legumes, and root nodules harbor different arbuscular mycorrhizal fungal communities. Appl Environ Microbiol 70: 6240–6246. doi:10.1128/aem.70.10.6240.
- Smith FA, Smith SE (2011) What is the significance of the arbuscular mycorrhizal colonisation of many economically important crop plants? Plant Soil 348: 63–79. doi:10.1007/s11104-011-0865-0.
- Smith S, Read D (1997) Mycorrhizal symbiosis. 2nd ed. San Diego: Academic Press.

- Sprent JI, Sprent P (1990) Nitrogen fixing organisms: pure and applied aspects. 1st ed. London: Charman and Hall.
- Tester AM, Smith SE, Smith FA, Walker NA (1986) Effects of photon irradiance on the growth of shoots and roots, on the rate of initiation of mycorrhizal infection and on the growth of infection units in *Trifolium subterraneum* L. New Phytol 103: 375–390. doi:10.1111/j.1469-8137.1986.tb00623.x.
- Van der Heijden MGA, Bardgett RD, Van Straalen NM (2008) The unseen majority: soil microbes as drivers of plant diversity and productivity in terrestrial ecosystems. Ecol Lett 11: 296–310. doi:10.1111/j.1461-0248.2007.01139.x.
- Vázquez-Yanes C, Orozco-Segovia A (1992) Effects of litter from a tropical rainforest on tree seed germination and establishment under controlled conditions. Tree Physiol 11: 391–400. doi:10.1093/treephys/11.4.391.
- Vejsadova H, Siblikova D, Gryndler M, Simon T, Miksik I (1993) Influence of inoculation with *Bradyrhizobium japonicum* and *Glomus claroideum* on seed yield of soybean under greenhouse and field conditions. J Plant Nutr 16: 619–629. doi:10.1080/01904169309364561.
- Verdú M, Traveset A (2005) Early emergence enhances plant fitness: a phylogenetically controlled meta-analysis. Ecology 86: 1385–1394. doi:10.1890/04-1647.
- Veresoglou SD, Menexes G, Rillig MC (2012) Do arbuscular mycorrhizal fungi affect the allometric partition of host plant biomass to shoots and roots? A meta-analysis of studies from 1990 to 2010. Mycorrhiza 22: 227–235. doi:10.1007/s00572-011-0398-7.

- Weiher E, van der Werf A, Thompson K, Roderick M, Garnier E, Eriksson O (1999)
Challenging Theophrastus: A common core list of plant traits for functional ecology. *J Veg Sci* 10: 609–620. doi:10.2307/3237076.
- Westoby M (1998) A leaf-height-seed (LHS) plant ecology strategy scheme. *Plant Soil* 199: 213–227. doi:10.1023/A:1004327224729.
- Yang XH, Luo XS, Liu RJ (1994) Effect of vesicular-arbuscular mycorrhiza on yield and quality of watermelon (in Chinese). *Fruit Sci* 11: 117-119.
- Yi H-S, Heil M, Adame-Alvarez RM, Ballhorn DJ, Ryu C-M (2009) Airborne induction and priming of plant defenses against a bacterial pathogen. *Plant Physiol* 151: 2152–2161. doi:10.1104/pp.109.144782.