

Portland State University

PDXScholar

Mechanical and Materials Engineering Faculty
Publications and Presentations

Mechanical and Materials Engineering

9-1998

Water Balloon Rupture in Low-G

Mark M. Weislogel

Portland State University, weisloge@pdx.edu

S. Lichter

Northwestern University

Follow this and additional works at: https://pdxscholar.library.pdx.edu/mengin_fac



Part of the [Fluid Dynamics Commons](#)

Let us know how access to this document benefits you.

Citation Details

Weislogel, M.M., Lichter, S., Water Balloon Rupture in Low Gravity, *Phys. of Fluids*, 10, No. 9, Gallery of Fluid Motion, p. S4, Sept., 1998.

This Article is brought to you for free and open access. It has been accepted for inclusion in Mechanical and Materials Engineering Faculty Publications and Presentations by an authorized administrator of PDXScholar. Please contact us if we can make this document more accessible: pdxscholar@pdx.edu.



Fig. 1



Fig. 3



Fig. 2



Fig. 4

WATER BALLOON RUPTURE IN LOW-G

Submitted by M. M. Weislogel¹ and S. Lichter²
(NASA Lewis Research Center,¹ Northwestern University²)

A qualitative study of the bursting of water balloons in a simulated low-gravity environment was conducted aboard NASA Lewis's DC-9 aircraft. Following rupture by a syringe needle, the balloon retracts tangent to the water surface leaving a smooth surface near the puncture location, Fig. 1. Asymmetries soon develop as the membrane rips apart, ejecting a directional spray from the surface. When the balloon parts from the drop entirely, it causes a large deformation of the remaining liquid mass leading to significantly underdamped oscillations which persist for the duration of the simulation.

In Fig. 2 an approximately 2.5 liter blob of undulating water hovers after rupture of the membrane. A free floating, red-dyed water blob flattens, crowns, and breaks up after impact by an impinging blue-dyed water jet, Fig. 3. In Fig. 4 a large air bubble is blown into a free floating blob using a straw. Such flows are *unearthly* in that a balance is struck between capillary and inertial forces over truly large length scales.

The tests were performed to develop techniques to rapidly deploy large liquid drops in a microgravity environment. The footage has also proven of general interest and is used to introduce students to low-g phenomena. Quicktime movies may be found at <http://zeta.lerc.nasa.gov/balloon/blob.htm>. Related ground tests were also performed using high-speed video photography and may be found at <http://zeta.lerc.nasa.gov/balloon/hs.htm>.