

Portland State University

PDXScholar

Chemistry Faculty Publications and
Presentations

Chemistry

2009

Exploring the pH Dependence of Viologen Reduction by Alpha-Carbon Radicals Derived from HCy and Cys

Dong Wang

Louisiana State University

William E. Crowe

Louisiana State University

Robert M. Strongin

Portland State University, strongin@pdx.edu

Martha Sibrian-Vazquez

Portland State University

Follow this and additional works at: https://pdxscholar.library.pdx.edu/chem_fac

 Part of the [Chemistry Commons](#)

Let us know how access to this document benefits you.

Citation Details

Wang, Dong; Crowe, William E.; Strongin, Robert M.; and Sibrian-Vazquez, Martha, "Exploring the pH Dependence of Viologen Reduction by Alpha-Carbon Radicals Derived from HCy and Cys" (2009). *Chemistry Faculty Publications and Presentations*. 57.

https://pdxscholar.library.pdx.edu/chem_fac/57

This Post-Print is brought to you for free and open access. It has been accepted for inclusion in Chemistry Faculty Publications and Presentations by an authorized administrator of PDXScholar. Please contact us if we can make this document more accessible: pdxscholar@pdx.edu.

Published in final edited form as:

Chem Commun (Camb). 2009 April 14; (14): 1876–1878. doi:10.1039/b819746f.

Exploring the pH dependence of viologen reduction by α -carbon radicals derived from Hcy and Cys[†]

Dong Wang^a, William E. Crowe^a, Robert M. Strongin^b, and Martha Sibrian-Vazquez^b

William E. Crowe: wcrowe1@lsu.edu

^aDepartment of Chemistry, Louisiana State University, Baton Rouge, LA 70803, USA. Fax: +1 225 578 3458; Tel: +1 225 578 2060

^bDepartment of Chemistry, Portland State University, Portland, OR 97201, USA. Fax: +1 503 725 9525; Tel: +1 503 725 9724

Abstract

The colorimetric reaction of homocysteine (HCy) with a series of viologen salts suggests a linear correlation between the mid-point reduction potential of Hcy-derived α -carbon radicals and pH.



There has been great interest in the development of selective methods for the detection of homocysteine (Hcy) as it is a biomarker for a wide range of diseases.¹ Interference from structurally related biological thiols such as Cys and GSH, typically present in much higher concentration, hinders selective detection methods. Unlike Hcy, the relatively more common biological thiols, such as GSH, are typically associated with beneficial antioxidant activity. It is thus also important to understand the differences between the fundamental chemistry of Hcy and other biothiols.

Oxidizing thiyl radicals formed from biological thiols interconvert with reducing, α -amino carbon-centered radicals under physiological, aerobic conditions (eqn (1)). Strongin *et al.* recently reported a colorimetric detection method based on the selective reduction of colorless methyl viologen (MV²⁺) to its blue radical cation (MV^{+•}) by Hcy at neutral pH.² The selectivity of this detection method for Hcy was proposed to arise from an intramolecular hydrogen atom abstraction converting the oxidizing Hcy thiyl radical to the reducing Hcy α -amino carbon-centered radical through a kinetically favorable, five-membered ring transition state (eqn (2)).³ Earlier work by Zhao *et al.* had shown that at pH 10.5 colorimetric detection of GSH, Cys, and Hcy was observed with no discernible selectivity.^{3,4} Interconversion of thiyl radicals with reducing disulfide radical anions (eqn (3)) becomes more prominent with increasing pH, and it is possible that at pH 10.5 nonselective MV²⁺ reduction results from the equally facile formation of disulfide radical anions from all three biological thiols.

[†]Electronic supplementary information (ESI) available: Experimental details and spectroscopic data of prepared viologens.

© The Royal Society of Chemistry 2009

Correspondence to: William E. Crowe, wcrowe1@lsu.edu.



Conducting the viologen-mediated colorimetric assay at neutral pH has the advantage of minimizing interference from nonselective MV^{2+} reduction by disulfide radical anions. However, captodative stabilization of α -amino carbon-centered radicals requires lone pair donation from nitrogen, and at neutral pH only a small fraction of the amino groups is not protonated. This begs the question whether the reducing species producing the colorimetric response in the MV^{2+} assay could really be an α -carbon radical. To address this question we propose herein a thermodynamic model for the pH dependence of the effective reduction potential of Hcy derived α -carbon radicals and, using viologen indicators of varying reduction potentials, demonstrate that colorimetric changes occur at experimental pH values consistent with this model.

We also report the first examples, to our knowledge, of viologen reduction mediated by α -amino carbon-centered radicals occurring at acidic pH.

The ability of the carbon radicals to reduce a given substrate *e.g.*, methyl viologen) has pH-dependence which is determined by Scheme 1.

The mid-point potential $E_m(RC^+/RC^\bullet)$ at a given pH determines whether the α -carbon radical will be able to reduce a particular substrate. The following relationship between $E_m(RC^+/RC^\bullet)$ and $[H^+]$ was derived from the Nernst equation:⁵

$$E_m(RC^+/RC^\bullet) = E^\circ(RC^+/RC^\bullet) + \frac{RT \ln 10}{F} \log \left[1 + \frac{H^+}{K_{obs}} \right] \quad (4)$$

where $E^\circ(RC^+/RC^\bullet)$ is the standard potential, which is also the minimum value possible for $E_m(RC^+/RC^\bullet)$, and the pseudo-equilibrium constant, K_{obs} , is given by:

$$K_{obs} = \frac{K_1 K_2 K_3}{K_1 K_2 + K_3}$$

When $[H^+] \gg K_{obs}$ (low pH), E_m varies linearly with pH:

$$E_m(\text{RC}^+/\text{RC}^\bullet) \approx E^\circ(\text{RC}^+/\text{RC}^\bullet) + \frac{RT \ln 10}{F} [\log[\text{H}^+] - \log K_{\text{obs}}] \quad (5)$$

At 298 K, $RT \ln 10/F = 59.2$ mV. Since $\text{pH} = -\log[\text{H}^+]$ and we can define $\text{p}K_{\text{obs}} = -\log K_{\text{obs}}$, the above equation can be re-written as:

$$E_m(\text{RC}^+/\text{RC}^\bullet) = E^\circ(\text{RC}^+/\text{RC}^\bullet) + 59.2 [\text{p}K_{\text{obs}} - \text{pH}] \quad (6)$$

When $[\text{H}^+] \ll K_{\text{obs}}$ (high pH), the pH-dependent part of the function goes to zero and E_m becomes constant, which is also the minimum value of E_m .

$$E_m(\text{RC}^+/\text{RC}^\bullet) = E^\circ(\text{RC}^+/\text{RC}^\bullet) \quad (7)$$

To test this model the effect of pH on the aminothiols-mediated reduction of a series of viologen indicators was determined. This was accomplished by gently refluxing a buffered,⁶ aqueous solution of the aminothiol (Hcy or Cys, 17 mM) and viologen (4.0 mM). Typically, color formation was observed (*e.g.*, colorless to blue for MV^{2+}) above a certain pH. As the pH was increased, faster onset of color formation and deeper color change were observed as well as longer persistence of color after the sample was removed from heating. For several viologens there was a pH range where color formation is observed for Hcy but not Cys (selective Hcy detection). Hcy should have a stronger reduction potential because of the particularly favorable formation of α -carbon radicals.

As solution pH decreases, we predict that $E_m(\text{RC}^+/\text{RC}^\bullet)$ should increase until $E_m(\text{RC}^+/\text{RC}^\bullet) > E^0(\text{viologen})$ at which point no color formation should be observed. Upon lowering the pH we observed slower onset of color formation and less intense color change as well as shorter persistence of the color after the sample was removed from heating. For six of the eight viologens⁷⁻⁹ examined we were able to obtain a pH endpoint (minimum pH where color formation was observable) below which no color formation occurred. As predicted, lower pH endpoints were observed for viologens possessing higher (*i.e.*, less negative) reduction potentials (E^0 values). No pH endpoint was observed (color formation for all pH values ≥ 0) for the two most easily reduced viologens. These pH endpoint results are summarized in Table 1. We propose that $E_m(\text{RC}^+/\text{RC}^\bullet) \approx E^0(\text{viologens})$ for the pH endpoints obtained in entries 4–8 of this table. As can be seen from Table 1, entry 3 through entry 8, viologens can be used to detect Hcy and Cys selectively during lower pH. For example, benzylic viologen (entry 5) can be used to detect Hcy selectively between pH 3.6 and pH 4.8 since there is no color change for Cys.

Interestingly, phenyl viologen (entry 4) is found to be an excellent reagent to selectively detect Hcy and Cys. It can selectively detect Hcy and Cys not only in acidic conditions, but also in basic conditions. From pH 9 to pH 12, Hcy shows a green color at first (and finally turned blue) whereas Cys shows blue. Especially at pH 11, Cys turned blue within 5 min of gentle refluxing, whereas Hcy turned green after 10 min of gentle refluxing.

It is worth noting that most of the viologens (from entry 4 to entry 8) we tested showed a linear correlation between pH endpoint and E^0 (standard reduction potential of α -carbon radicals). This linear relation exactly matches what we expected from eqn (6). Using the experimental data, we are able to get a line between pH_{min} and E^0 (Fig. 1).

In summary, α -aminoalkyl radicals can be generated under acidic conditions. The reduction potential of α -carbon radicals was found to be pH dependent, and it shows a linear correlation between the reduction potential and pH for Hcy. The pH profiles of Cys and Hcy have been established for six viologens. Most of the viologens can be used to selectively detect Hcy and Cys under acidic conditions. The determination of standard reduction potentials of α -carbon radicals is currently underway in our lab.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

Support from the National Institutes of Health (R01 EB002044) is gratefully acknowledged.

Notes and references

1. (a) Nekrassova O, Lawrence NS, Compton RG. *Talanta* 2003;60:1085. [PubMed: 18969134] (b) Refsum H, Smith AD, Ueland PM, Nexø E, Clarke R, McPartlin J, Johnston C, Engbaek F, Schneede J, McPartlin C, Scott JM. *Clin. Chem* 2004;50:3. [PubMed: 14709635] (c) De Bree A, Verschuren WMM, Kromhout D, Kluijtmans LAJ, Blom HJ. *Pharmacol. Rev* 2002;54:599. [PubMed: 12429870]
2. (a) Wang WH, Rusin O, Xu XY, Kim KK, Escobedo JO, Fakayode SO, Fletcher KA, Lowry M, Schowalter CM, Lawrence CM, Fronczek FR, Warner IM, Strongin RM. *J. Am. Chem. Soc* 2005;127:15949. [PubMed: 16277539] (b) Wang WH, Escobedo JO, Lawrence CM, Strongin RM. *J. Am. Chem. Soc* 2004;126:3400. [PubMed: 15025448]
3. Zhao R, Lind J, Merenyi G, Eriksen TE. *J. Am. Chem. Soc* 1994;116:12010.
4. Zhao R, Lind J, Merenyi G, Eriksen TE. *J. Chem. Soc., Perkin Trans* 1997;2:569.
5. See, for example: Wardman P. *J. Phys. Chem. Ref. Data* 1989;18:1637.
6. Components of buffer solutions were as follows: pH 0.0–2.2: HCl + KCl; pH 2.3–6.9: citric acid + Na₂HPO₄; pH 7.0–9.0: Tris + HCl; pH 10.0–11.0: NaHCO₃ + NaOH; pH 12.0–13.0: NaOH + KCl.
7. While viologens in entries 1, 4, 5, 7 and 8 are commercially available, others were prepared by nucleophilic alkylation of 4,4'-bipyridine as described in the literature.⁸ Dimethoxy viologen was prepared by alkylation of *N*-oxide according to a known process⁹.
8. Bruinink J, Kregting CGA, Ponjee JJ. *J. Electrochem. Soc* 1977;124:1854.
9. Lorance ED, Kramer WH, Gould IR. *J. Am. Chem. Soc* 2002;124:15225. [PubMed: 12487598]
10. Feely W, Lehn WL, Boekelheide V. *J. Org. Chem* 1957;22:1135.




Fig. 1.
Reduction potential vs. pH plot.





Fig. 2.
Reduction potential vs. whole pH range.




Scheme 1.

The equilibrium between thiyl radicals and carbon radicals.

Table 1

The minimum pH that α -carbon radicals can reduce viologens



Entry	R (viologen)	E^0 /mV	pH endpoint ^a of Hcy	pH(calc.) ^b of Hcy	pH endpoint of Cys
1	2,4-Dinitrophenyl	ND	None	<0	≤0
2	-CH ₂ CN	-150	ND	0	≤0
3	-CH ₂ CO ₂ Et	-267	3.8	1.9	5.8
4	-Ph	-288	2.5	2.3	3.8
5	-CH ₂ Ph	-370	3.6	3.7	4.8
6	-CH ₂ CH=CH ₂	-408	4.2	4.3	4.7
7	-CH ₂ (CH ₂) ₅ CH ₃	-415	4.3	4.4	5.8-7.0, 8.8
8	-CH ₃	-446	5.0	4.9	6.0-7.0, 8.2

^aData from entries 4–8 show a linear correlation between E^0 and pH endpoint. The best linear fit of this data gives the empirical equation: $E^0 = -132.57 - 64.498 \text{ pH}$ ($R^2 = 0.9872$). See Fig. 1.

^bpH(calc.) is calculated from $(-153.3 - E^0)/59.2$ which represents the best fit of the data from entries 4–8 to the theoretical equation $E^0(\text{viologen}) = E^0(\text{Hcy}) + 59.2(\text{pK}_{\text{obs}} - \text{pH})$. We calculate an average value: $[E^0(\text{Hcy}) + 59.2 \text{ pK}_{\text{obs}}]_{\text{avg}} = [E^0(\text{viologen}) + 59.2\text{pH}]_{\text{avg}} = -153.3 \text{ mV}$.