

Portland State University

PDXScholar

Environmental Science and Management
Faculty Publications and Presentations

Environmental Science and Management

12-2012

Long-Term Integrated Studies Show Complex and Surprising Effects of Climate Change in the Northern Hardwood Forest

Peter M. Groffman

Cary Institute of Ecosystem Studies

Lindsay E. Rustad

USDA Forest Service's Northern Research Station

Pamela H. Templer

Boston University

John L. Campbell

USDA Forest Service's Northern Research Station

Lynn M. Christenson

Vassar College

Follow this and additional works at: https://pdxscholar.library.pdx.edu/esm_fac



Part of the [Forest Science Commons](#), [Forest Sciences Commons](#), and the [Sustainability Commons](#)

Let us know how access to this document benefits you.

Citation Details

GROFFMAN, P. M., RUSTAD, L. E., TEMPLER, P. H., CAMPBELL, J. L., CHRISTENSON, L. M., LANY, N. K., & ... PARDO, L. H. (2012). Long-Term Integrated Studies Show Complex and Surprising Effects of Climate Change in the Northern Hardwood Forest. *Bioscience*, 62(12), 1056-1066. doi:10.1525/bio.2012.62.12.7

This Article is brought to you for free and open access. It has been accepted for inclusion in Environmental Science and Management Faculty Publications and Presentations by an authorized administrator of PDXScholar. Please contact us if we can make this document more accessible: pdxscholar@pdx.edu.

Authors

Peter M. Groffman, Lindsay E. Rustad, Pamela H. Templer, John L. Campbell, Lynn M. Christenson, Nina K. Lany, Anne M. Socci, Matthew A. Vadeboncoeur, Paul G. Schaberg, Geoffrey W. Wilson, Charles T. Driscoll, Timothy J. Fahey, Melany C. Fisk, Christine L. Goodale, Mark B. Green, Steven P. Hamburg, Chris E. Johnson, Mryon J. Mitchell, Jennifer L. Morse, Linda H. Pardo, and Nicholas L. Rodenhouse

Authors

Peter M. Grozman, Jennifer L. Morse, Lindsay E. Rustad, John L. Campbell, Pamela H. Templer, Lynn M. Christenson, Nina K. Lany, Anne M. Soci, Matthew A. Vadeboncoeur, Paul G. Schaberg, and Geoffrey W. Wilson

Long-Term Integrated Studies Show Complex and Surprising Effects of Climate Change in the Northern Hardwood Forest

PETER M. GROFFMAN, LINDSEY E. RUSTAD, PAMELA H. TEMPLER, JOHN L. CAMPBELL, LYNN M. CHRISTENSON, NINA K. LANY, ANNE M. SOCCI, MATTHEW A. VADEBONCOEUR, PAUL G. SCHABERG, GEOFFREY F. WILSON, CHARLES T. DRISCOLL, TIMOTHY J. FAHEY, MELANY C. FISK, CHRISTINE L. GOODALE, MARK B. GREEN, STEVEN P. HAMBURG, CHRIS E. JOHNSON, MYRON J. MITCHELL, JENNIFER L. MORSE, LINDA H. PARDO, AND NICHOLAS L. RODENHOUSE

Evaluations of the local effects of global change are often confounded by the interactions of natural and anthropogenic factors that overshadow the effects of climate changes on ecosystems. Long-term watershed and natural elevation gradient studies at the Hubbard Brook Experimental Forest and in the surrounding region show surprising results demonstrating the effects of climate change on hydrologic variables (e.g., evapotranspiration, stream flow, soil moisture); the importance of changes in phenology on water, carbon, and nitrogen fluxes during critical seasonal transition periods; winter climate change effects on plant and animal community composition and ecosystem services; and the effects of anthropogenic disturbances and land-use history on plant community composition. These studies highlight the value of long-term integrated research for assessments of the subtle effects of changing climate on complex ecosystems.

Keywords: climate change, forests, long-term studies, northeastern United States, winter

A dominant approach in climate change research has been to focus on the effects of changes in temperature and precipitation on broadscale ecosystem properties over large areas and long periods. This body of research suggests that climate change will substantially alter the distribution of species and the function of ecosystems (e.g., Iverson and Prasad 2001), with important effects on ecosystem services. These analyses are based on well-described effects of temperature and precipitation on the distribution and activity of organisms. However, climate change is playing out over the complex and dynamic hydrobiogeological structure of the landscape—that is, the intertwined patterns of soils, vegetation, and hydrologic flowpaths, with a spatially variable history of land use and a wide range of current human activities and concurrent environmental changes. The climate effects on ecosystem structure and function may be modified by interactions with these patterns and histories over a range of time scales. We assert that a key approach to

unraveling this daunting complexity is long-term studies, including those in which natural elevation gradients are exploited, as a foundation for detailed studies of critical and often unexpected climate-induced changes in forest structure and function.

In this article, results from the Hubbard Brook Experimental Forest (HBEF) and the surrounding region are used to illustrate how long-term studies can serve as a foundation for addressing the complex interactions that ultimately determine the effects of climate change on ecosystems. We combine data from long-term (50-year) measurements of multiple aspects of climate and ecosystem structure and function to highlight important but poorly studied interactions that could be critical determinants of the responses of plant and animal communities, fluxes of water, element dynamics, and services in northern hardwood forest ecosystems. Our objective is to demonstrate how a combination of long-term and in-depth measurements facilitates

the understanding of the interplay between climate and forest ecosystem dynamics and how this interplay produces unanticipated and important new results.

In the sections below, we first present basic information on changing climate at the HBEF and show how this change acts on a hydrobiogeological template in ways that alter key hydrologic variables (e.g., evapotranspiration, stream flow, soil moisture) that influence multiple aspects of ecosystem structure and function. Climate change is also interacting with a variety of temporally dynamic ecosystem processes, and we describe how changes in the phenology of vegetation interact with climate change to affect fluxes of water, carbon, and nitrogen in the transitions between seasons. Temporal complexities in ecosystem processes are also highlighted by a discussion of unexpected effects of climate change in winter, which has long been considered to be a "dormant" season but has recently been shown to be functionally important and to have legacy effects during the growing season (Campbell et al. 2005). We end with a description of how the complex expression of climate change over landscape (approximately 3000 hectares [ha]) and regional (100,000 square kilometers) scales and over 50–300-year time frames influences plant community composition and likely future ecotone shifts in response to broad changes in climate zones (Iverson and Prasad 2001, Mohan et al. 2009).

The HBEF was established by the US Department of Agriculture Forest Service in 1955 as a center for hydrologic research in New England. The Hubbard Brook Ecosystem Study (HBES) originated in 1960, centered on the idea of the small-watershed approach to studying element fluxes and cycling (Likens and Bormann 1995). The site has been a component of the US National Science Foundation's Long Term Ecological Research (LTER) Network since 1988. The long-term studies in the HBES have maintained a balance between the original research interests (e.g., air pollution, land disturbance) and new foci, such as climate change, that have emerged in recent decades. Maintaining this balance is facilitated by ongoing scientific peer-review cycles, especially those associated with the LTER Network, and an open governance structure that encourages the participation of new investigators and the incorporation of new research ideas (Groffman et al. 2004).

The 3037-ha Hubbard Brook Valley is located in the White Mountain National Forest, in New Hampshire. The climate is humid continental, with short, cool summers and long, cold winters. The mean air temperature (measured between 1955 and 2000) is 19 degrees Celsius (°C) in July and -9°C in January (Campbell et al. 2007). The annual precipitation averages approximately 140 centimeters (cm), and a continuous snowpack usually develops early each winter to a depth of approximately 1.5 meters (Campbell et al. 2007). Vegetation is dominated by northern hardwood forests (e.g., American beech [*Fagus grandifolia*], sugar maple [*Acer saccharum*], yellow birch [*Betula alleghaniensis*]) that grade into red spruce (*Picea rubens*) and balsam fir

(*Abies balsamea*) along a dynamic ecotone at higher elevations (Schwarz et al. 2003). Soils are 75–100 cm deep, acidic (pH ~4.0 in the surface organic horizons) typical and aquic haplorthods developed from unsorted basal tills. The HBEF was subjected to logging from 1880 to 1920 and then a salvage logging project following a hurricane in 1938. Grafton County, in which the HBEF is located, has a diverse history of agricultural activities beginning in the late eighteenth century, peaking in the 1860s, followed by almost complete abandonment of agricultural activities by the mid-twentieth century.

Climate change effects on the hydrobiogeological template that underlies the structure and function of the northern hardwood forest ecosystem

The climate of the HBEF has changed over the last half century. At the four weather stations with the longest records (43–52 years), the average annual air temperature has increased by 0.17°C–0.29°C per decade, with more marked warming in winter than in summer (Campbell et al. 2007, Hamburg et al. 2012). These local trends in air temperature are characteristic of the region and are expected to continue into the future, with projected increases of 2.1°C–5.3°C by 2100 (Hayhoe et al. 2007, Huntington et al. 2009). Precipitation volume has also increased at the HBEF and is indicative of regional trends. The long-term average annual precipitation ranges from 131 to 149 cm across the stations within the HBEF and has increased by 3.5–6.7 cm per decade, or 13%–28% over 50 years (Campbell et al. 2007). The trends in precipitation at the rain gauges with the longest records are stronger because of the influence of a protracted drought in the mid-1960s.

Winter precipitation has changed less than precipitation in other seasons, which, when it is combined with warmer winter air temperatures, has led to significant reductions in snowpack accumulation. Long-term (since 1959) weekly measurements of snow depth and snow water equivalent indicate that the maximum annual snowpack depth has declined by 4.8 cm per decade (1.4 cm snow water equivalent), and the number of days with snow cover has declined by 3.9 days per decade (Campbell et al. 2010). The annual and winter precipitation are projected to increase by 7%–14% and 12%–30%, respectively, by 2100, whereas summer precipitation is expected to show little change (Campbell et al. 2009). The climate projections for the northeastern United States are based on regionally downscaled, coupled atmosphere–ocean general-circulation models that are driven by scenarios describing how population, technology, and energy use might develop in the future (Hayhoe et al. 2007).

Snowpack decline and precipitation increases have, in turn, influenced stream flow. Measurements of daily stream flow from the gauged reference watersheds at the HBEF since 1956 show that snowmelt-induced peak flows in spring have declined (Campbell et al. 2011) and have occurred earlier (Hamburg et al. 2012). In contrast, both winter and summer stream flow have increased. In winter, the increase is due to

reduced storage of precipitation in the snowpack, whereas in summer (typically a low-flow season), the increase is due to increases in precipitation and concomitant declines in evapotranspiration, as is described below. The increase in summer streamflow at the HBEF may be beneficial to some aquatic organisms, such as brook trout, whose upstream movement is limited by physical obstructions during low flows (Warren et al. 2008). The increase may have negative effects on nitrogen uptake or retention by streams, which is strongly flow dependent.

Interestingly, evapotranspiration, when measured indirectly as the difference between annual precipitation and annual streamflow, has shown slight but significant declines over the period of record beginning in 1959 (Figure 1a; Campbell et al. 2011). The cause of the decline in evapotranspiration is not known but probably reflects decreases in transpirational water loss by vegetation that result from changes in vegetation composition, structure, or productivity or from possible reductions in stomatal conductance owing to increasing atmospheric carbon dioxide concentrations. Also probably influencing this trend are reductions in evaporative demand from the atmosphere driven by declines in solar radiation, vapor pressure deficit, and wind speed that are occurring at the HBEF as they are elsewhere (Roderick et al. 2009).

Changes in evapotranspiration and precipitation have effects on soil moisture. Although changes in soil moisture have clear consequences for ecological processes, the future direction of change is not well established. Although historical observations (since 1959) from the HBEF suggest that declines in evapotranspiration and increases in precipitation have led to increases in soil moisture (Figure 1b), projections of future change from models indicate that soil moisture may decline because of increases in transpiration associated with enhanced net primary productivity in a warmer, wetter climate (Campbell et al. 2009, Huntington et al. 2009). These projected changes are highly uncertain, however, and indicate the need for continued monitoring and mechanistic studies (e.g., of carbon-dioxide-induced changes in stomatal conductance) in order to develop a better predictive capability for this key driver of climate change impacts on ecosystems. Moreover, soil moisture is not influenced by climate alone; its spatial variation is strongly controlled by the physical effects of geomorphology and soil texture. Therefore, changes in climate are superimposed on a geologic template that mediates the effects of climatic drivers on ecological processes.

Insights into the potential structural and functional ecosystem changes arising from changes in soil moisture and temperature can also be gained by exploiting the natural elevation gradient that creates a difference of approximately 20°C in mean annual temperatures between low- and high-elevation locations at the HBEF that show few differences in soil and vegetation types (Campbell et al. 2007). Observed differences in soil moisture along this gradient are large enough to affect the abundance, composition, and dynamics of soil biota, including heterotrophs such as salamanders, insects, and snails (Rodenhouse et al. 2009). Hydrologic and gaseous nitrogen losses, as well as soil microbial processes (e.g., mineralization, nitrification) that increase nitrogen availability to plants have also been shown to be positively correlated with soil moisture at the HBEF (Groffman et al. 2009). Therefore, if drier soil conditions result from continued climatic warming, nitrogen availability to vegetation may decrease at the HBEF, which would exacerbate the nitrogen limitation of productivity but would decrease the potential for nitrogen saturation and losses to the environment. Greenhouse gas fluxes are also affected, with lower production of nitrous oxide and increased consumption of methane with lower levels of




Figure 1. Changes in precipitation, streamflow, and evapotranspiration (a) and soil moisture at three depths (b) in the hydrologic reference watershed at the Hubbard Brook Experimental Forest, New Hampshire. Abbreviations: Avg., average; cm, centimeters; mm, millimeters.

soil moisture (Groffman et al. 2009). Soil moisture can also have a substantial effect on sulfur dynamics. For example, with decreasing atmospheric sulfur deposition, annual discrepancies in the watershed mass balances of sulfur are increasingly regulated by watershed moisture, because high-moisture soil conditions stimulate the net mineralization of soil sulfur pools (Mitchell and Likens 2011).

Life on the edges: Climate change amplifies temporal asynchronies between plant and soil microbial processes

A particularly intriguing aspect of climate change is the lengthening of growing seasons: Spring is advancing and fall is retreating (Huntington et al. 2009, Ibanez et al. 2010). In the spring, the onset of the growing season in northern latitudes is heralded by a complex suite of environmental drivers, including a lengthening of the photoperiod; warming of the air and soil; shifts in precipitation from snow to rain; melting of the snowpack; thawing of surface soils; increases in available soil water; and, ultimately, budburst and leaf development (Jeong et al. 2012). The complete process of canopy expansion in spring may take several weeks in northern hardwood forest ecosystems.

Previous studies have addressed the asynchrony in the early spring when the amount of microbial mineralization of organic matter and solute transport through soil may be substantial but nutrient uptake by roots and movement to developing aboveground canopies may be minimal (Muller and Bormann 1976). Nutrient immobilization by microbes and uptake by spring ephemerals and understory vegetation can retard the loss of nutrients during this early spring period (Muller and Bormann 1976, Zak et al. 1990). Although it is now widely recognized that many belowground biological processes such as nutrient mineralization and immobilization continue under the snowpack and in soils with temperatures slightly above freezing, the rates of these processes are, in general, relatively low in winter (Brooks et al. 2011). Rapid biological activity is thought to commence when soil temperatures reach or exceed 4°C. The advance in spring snowmelt and warming air temperatures with climate change result in soil temperatures reaching biologically favorable conditions at earlier dates as well. It is not clear, however, whether the advance in soil warming is or will continue to be synchronous with advances in aboveground plant phenology (Polgar and Primack 2011). If changes in soil warming and plant phenology are not synchronous, there is the potential for greater decoupling of belowground microbial activity and the onset of rapid plant nutrient and water uptake that could lead to an increase in nutrient losses from the ecosystem.

At the HBEF, the snow-free period has increased by approximately 20 days over the last half century (Campbell et al. 2010). Green canopy duration has increased by approximately 10 days (Richardson et al. 2006), and the timing of the last measurable snowpack (around 13 days), the date of the last frost (about 8 days), and the timing of leaf expansion

(approximately 8 days) are earlier than they were in the mid-1950s (Richardson et al. 2006, Hamburg et al. 2012). In the fall, the dates of the first frost and the first measurable snowpack have been delayed by approximately 10 and 8 days, respectively (Hamburg et al. 2012). These observations illustrate the marked changes that have occurred in the physical environment in the spring and fall seasons, which may exacerbate existing seasonal asynchronies between microbial activity and plant uptake.

A closer examination of soil temperature data coupled with detailed phenological studies provides a new perspective on changes in spring asynchrony. Continuously recorded (hourly) soil temperature data are available at four northern hardwood Soil Climate Analysis Network sites—two in New Hampshire (including HBEF) and two in Vermont—for the period of 2002–2011. Data for all four sites and nine full years of record indicate a rapid increase of up to 8°C in surface soil temperature within 48 hours in early spring, usually occurring immediately after the snowpack melts completely (as is shown in Figure 2 for a midelevation site measured in 2003). The likely cause of this sudden increase in soil temperature is the abrupt decrease in surface albedo in response to the loss of the reflective snowpack and exposure of dark soils to solar radiation unhindered by the forest canopy in these deciduous forest sites. This rapid and abrupt spike in soil temperature probably triggers immediate increases in biological activity (Campbell et al. 2005) and may serve as an index marking the onset of spring soil conditions under deciduous canopies, similar to the use of the timing of ice-out as an indicator of dramatic changes in the physical environment of lakes (Magnuson et al. 2000).

At the HBEF, discrete measurements of soil temperature and snowpack depth dating back to 1956 allow us to reconstruct the approximate timing of this soil temperature trigger in early spring (see Figure 2b). For this analysis, time-series data were evaluated using the nonparametric Mann–Kendall test for trends. Prior to this analysis, the autocorrelation function test was used to ensure that the data met the assumption of serial independence. Trend lines were fitted by calculating the median slope of all the possible pairs in the data set. These records suggest that the date of snowmelt and the rapid increase in soil warming has advanced by 14.3 days over the period of record (1956–2010), or an average of 2.6 days per decade. Reconstructions of the concurrent trends in leaf phenology following Richardson and colleagues (2006) show that the timing of nearly complete canopy expansion for sugar maple, one of the dominant tree species at the site, advanced by only 6 days during the same period, or 1.1 days per decade. These data suggest that the period during which belowground temperatures are most conducive to biological activity (i.e., higher than 4°C) but the overstory vegetation is not maximally transpiring and taking up water and nutrients has increased by approximately 8 days over the past half century. This lengthening gap between belowground and canopy activity ought to elevate water and nutrient availability beyond plant uptake




Figure 2. An example of (a) the relationship between the loss of snowpack (snow depth, in centimeters [cm]), soil temperature (in degrees Celsius), and spring leaf phenology and (b) changes in the modeled day of nearly complete canopy expansion and last snowpack at a middle elevation site in the Hubbard Brook Experimental Forest. In panel (a), data from 2003 show how the disappearance of snowpack (shown by the horizontal shading) coincides with an abrupt increase in soil temperature at 4 cm (the solid line). The closed circles represent spring leaf phenology, and the arrow denotes the period of vernal asynchrony (see the text) that has increased over time. Spring canopy expansion (the closed circles) is becoming earlier, at a rate of 1.1 days per decade ($p = .02$), whereas the day of last snowpack (the open squares) is becoming earlier at a rate of 2.6 days per decade ($p = .004$); therefore, the length of the period between snowmelt and full canopy development has increased by over 8 days in 55 years.

capacities and could exacerbate nutrient leaching. Further studies are needed to evaluate the impacts of this vernal asynchrony on above- and below-ground plant and microbial dynamics, carbon and nutrient retention, and plant growth in north temperate forest ecosystems.

These climate-change-driven seasonal asynchronies may be enhanced regionally if deciduous cover increases at the expense of conifer cover, as is expected with climate change in northeastern North America (Mohan et al 2009). For example, spring snowmelt is earlier in deciduous than in coniferous forests because of radiation absorption and reflection by the evergreen canopy; therefore, soil warming is delayed to a time when air temperature is generally higher. Moreover, the evergreen canopy, responding directly to increasing soil and air temperatures (Schwarz et al. 1997), demands water and nutrients earlier than the deciduous forest does, which thereby reduces the potential for asynchrony. Therefore, the spatial extent of the seasonal decoupling of water, carbon, and nitrogen cycles is likely to be larger if conifer abundance decreases across the region.

More surprises in winter

The well-documented past and projected changes in the duration and depth of the winter snowpack in northern forests (Hayhoe et al. 2007, Campbell et al. 2010) have stimulated recent interest in winter ecology, biogeochemistry, and trophic interactions in northern forest ecosystems (Campbell et al. 2005). There has, however, been less attention focused on potential changes in tree species composition caused by both limits in tolerance to winter climate change and shifts in competitive outcomes (Kreyling 2010) and on the impacts on human activities (e.g., logging, recreation, fall foliage viewing, maple syrup production, the potential lengthening of the spring mud season).

Winter biogeochemistry studies at the HBEF have built on the long-term snow cover record and have included analyses of the effects of soil-freezing events caused by low snow cover on

watershed nitrate exports (Mitchell et al. 1996, Fitzhugh et al. 2003), plot manipulation studies of snow cover and soil freezing (Groffman et al. 2001, 2010, Templer et al. 2012), and studies of winter processes along the natural elevation gradient at the site (Groffman et al. 2009). These studies have shown that soil-freezing events can lead to significant increases in nitrate, phosphate, and base cation losses (Fitzhugh et al. 2001) that appear to be driven by increases in the root mortality of some tree species, such as sugar maple (Tierney et al. 2001, Cleavitt et al. 2008). However, the response to soil-freezing events varies—that is, there are not always marked increases in nitrate losses (Hentschel et al. 2009, Groffman et al. 2010). This variation in nitrate loss may be driven by variation in the response of dissolved organic carbon dynamics to soil freezing. In some cases, soil freezing mobilized dissolved organic carbon, which stimulates immobilization or denitrification, which, in turn, prevents a nitrogen response (Groffman et al. 2010). Moreover, the future trajectory of soil frost is also uncertain. Although a decrease in snow cover certainly increases the vulnerability of the ecosystem to soil-freezing events, at the same time, warming air temperatures decrease the potential for soil freezing (Campbell et al. 2010).

Evaluating the long-term effects of winter climate change on plant communities is particularly challenging. A reduction in the depth or duration of the winter snowpack could alter the reproduction and survivorship of some tree species, which could lead to changes in forest composition. As was mentioned above, sugar maple roots appear to be particularly susceptible to soil-freezing damage, which has contributed to episodes of sugar maple decline in northeastern North America (Auclair et al. 2010). Yellow birch benefits from enhanced seed dispersal on snow (Greene and Johnson 1997), and therefore, the regenerative success

of this species could also be diminished if the snowpack is reduced.

In addition to plants' physiological responses to climate shifts, animals may generate novel stresses on plant communities. For example, a decreased depth of snowpack can generate increased stress on young trees from foraging ungulates, such as moose (*Alces alces*; Faison et al. 2010). Either competitive outcomes or regeneration success could be affected by these compounded stresses (Crone et al. 2001), which would result in changes in forest composition that would not be predicted from simple climate-vegetation relationships alone (Mohan et al. 2009, Iverson et al. 2011). Similarly, even though climate envelope models project an increasing abundance of eastern hemlock (*Tsuga canadensis*) throughout the eastern United States, this species is threatened by the hemlock woolly adelgid (*Adelges tsugae*). The distribution of the adelgid is currently limited in the north by winter temperatures, because it cannot tolerate temperatures below -25°C (Paradis et al. 2008). However, there has been a 40% decline in the number of days with a mean temperature of -25°C from the first to the last decade in the 50-year temperature record at the HBEF (Hamburg et al. 2012). With the combination of increasing temperatures and the potential selection for greater cold tolerance within hemlock woolly adelgid populations, eastern hemlock trees could be affected by this pest throughout their entire range within the next 30 years (Albani et al. 2010). This could lead to an increase in the abundance of hardwood tree species (Albani et al. 2010) and to increases in the rates of nutrient cycling, transpiration, and nutrient loss throughout this region (Campbell et al. 2009).

Shifts in winter climate will undoubtedly continue to influence wildlife in the HBEF region (Rodenhouse et al. 2009). As with vegetation, the impact of winter climate change

on wildlife populations plays out in concert with other environmental and societal stressors, including historical and current land use, forest regeneration patterns, and hunting pressure on game species (Christenson et al. 2010). For example, as snow depth declines, a shift in the dominant ungulate herbivore could occur. Moose are currently the dominant herbivore at the HBEF (Figure 3). They remove large volumes of browse material (approximately 25 kilograms dry weight per day) with preferential consumption of balsam fir during winter (Christenson et al. 2010). As snow depth declines below 45 cm, white-tailed deer (*Odocoileus virginianus*) will have greater access to these forests, because their movement is generally limited by deep snow (Morrison et al. 2003). We may be observing this transition currently;




Figure 3. Average snow depth (in meters) at low (less than 1500 feet; the dark shading) and high (above 2500 feet; the light shading) elevations at the Hubbard Brook Experimental Forest (HBEF) and estimated deer, moose, and bear populations (in number of individuals per square kilometer) for the HBEF (the solid line) and for the state of New Hampshire (the dashed lines) from 1980 to 2010.

observations at the HBEF and by the State of New Hampshire indicate a slight decline in the number of moose and an increase in the number of white-tailed deer beginning in 2005, which coincides with less snow reported at the HBEF at lower elevations (Figure 3). In addition to competing for resources, white-tailed deer have a negative influence on moose through the transport of a parasite lethal to moose (*Parelaphostrongylus tenuis*; Whitlaw and Lankester 1994). White-tailed deer preferentially browse a large number of woody species, including balsam fir, sugar maple, and oak, and avoid or positively affect striped maple, American beech, and black cherry (Crife et al. 2001). This change in browsing could profoundly influence tree species composition (e.g., the continued presence of spruce and beech and a reduction or elimination of selected hardwoods) in ways not predicted by models based on simple climate-vegetation relationships that suggest that forest composition in New Hampshire will transition to an oak-dominated forest.

Not all wildlife species will respond to changes in climate. Black bear (*Ursus americanus*) populations, for example, have remained relatively stable at the HBEF and across New Hampshire over the past two decades (Figure 3). Bears are sometimes referred to as a landscape species (Samson and Huot 1998), because they can live in landscapes with a variety of vegetation; therefore, they are less susceptible to climate-induced changes in plant community composition. Bears also den during the winter season, so snow levels at this time of year do not have as much of an impact on them as on other animals, including deer. Together, these characteristics will allow bears to remain in the HBEF region, barring increased disturbance by humans.

Similar to aboveground plants and animals, belowground organisms are also vulnerable to shifts in winter climate. Soil frost decreases soil arthropod richness and diversity and reduces the abundance of many arthropod groups in northern hardwood forests (Templer et al. 2012). These responses may adversely affect other forest floor taxa and the entire forest food web. For example, reduced food availability could have negative consequences for the reproduction, winter survival, or migration of vertebrate insectivores that feed from the forest floor (e.g., salamanders [e.g., *Plethodon cinereus*], ovenbirds [*Seiurus aurocapilla*], or the multiple species of thrushes [*Catharus* spp.] that breed in the forest; Holmes and Sherry 2001). Conversely, some migratory birds (e.g., those feeding from foliage or on emerging aquatic insects) may benefit from advancing phenology. Warm spring conditions lengthen the breeding season and can allow facultative double brooders such as the black-throated blue warbler (*Setophaga caerulescens*) to increase fecundity by attempting a second brood later in the season, provided sufficient food is available (Holmes et al. 1992). Although food may become available earlier under warm spring conditions, the mismatch between the arrival or breeding of migratory songbirds and a peak in food abundance that occurs in European oak woodlands (Both 2006), where the larvae of a single species—the winter moth (*Operophtera*




Figure 4. Possible changes in maple sap production as winter climates warm in the northeastern United States and adjacent Canada during this century. The projections are mean estimates based on modeled results for changes in the number of days of sap flow (Duchesne et al. 2009) and the overall yield of sap (Skinner et al. 2010) under various climate change scenarios.

brumata) dominates bird diets, is not expected at the HBEF. Similar to patterns found in deciduous European forests that are not dominated by oak (*Quercus*) species (Burger et al. 2012), foliage-gleaning birds in the northern hardwood forests at the HBEF feed on a diverse community of Lepidoptera larvae and adult insects of other species that develop during a wide window encompassing the entire avian breeding season. However, the early emergence of arthropods and earlier breeding by birds may put both at greater risk to losses associated with late-season frosts.

Climate change in winter could have a variety of substantial and potentially important human dimensions. For example, reduced snowpack and warmer winter temperatures are predicted to lead to a reduction of 20% or more in sugar maple sap yield (Duchesne et al. 2009) and to shorten the sap season by about half (Skinner et al. 2010) as the climate warms over the twenty-first century (Figure 4), which will challenge the economic viability of the regional maple sugar industry. A decreased snowpack could also hasten the appearance of the mud season—a time when unpaved roads and trails become unstable and impassable as snowpack melts and soils thaw. Winter timber extraction, which typically occurs on snow-covered soil to reduce soil compaction and erosion from skid trails, could also be adversely affected by decreased snowpack.

Changes to winter snowpack are also likely to affect winter recreation. Increased snowmaking at ski resorts over the last few decades has compensated for declines in natural snow accumulation. However, opportunities for snowmaking will decrease as winter temperatures continue to rise. At the HBEF, the number of days with temperatures producing good conditions for snowmaking (i.e., daily average temperatures of 20°C or lower; Steiger and Mayer 2008) has declined significantly over the early-season period of November through January, with December showing the most significant change. Another immediate albeit complex concern is the possibility

that climate change is facilitating the northward spread of Lyme disease and the impact that this will have on mammals, including humans (Subak 2003). These examples point to the need for research on the emerging aspects of climate change that are most likely to resonate with the general public.

Climate change is just one of many factors driving vegetation change in the northern hardwood forest

Since the most recent glaciation, vegetation communities have undergone a series of shifts, responding to changes in climate and other disturbances, including disease, fire, and human use of the landscape. Although many vegetation assemblages are clearly related to the soil parent material and hydrology characteristics of the hydrobiogeological template (Leak 1978), vegetation community interactions with animals and, most noticeably, the legacies of human use (e.g., timber removal and clearing for agriculture) have generated a complex mosaic of forest communities not necessarily at steady state with respect to current climate or habitat (Vadeboncoeur et al. 2012). Human land-use history has shaped unique vegetation associations (Cogbill 2000), each of which may respond differently to climate change. Therefore, it is important to consider climate change as one of multiple drivers of vegetation composition that interact on a template of hydrobiogeochemical variation and disturbance history. The great challenge is to understand how anthropogenic disturbances such as atmospheric deposition; the introduction of exotic species, pests, and pathogens; and forest fragmentation will modify the template on which climate change is acting and will contribute to future vegetation community shifts (Iverson et al. 2011).

Over the past two centuries, hardwoods have displaced conifers such as spruce at intermediate montane elevations in northern New England (Hamburg and Cogbill 1988, Vadeboncoeur et al. 2012), whereas more recent data show a range of species shifts, from further upward encroachment of the northern hardwood zone in the Green Mountains of Vermont (Beckage et al. 2008) to apparent increases in spruce and fir at the HBEF. Hamburg and Cogbill (1988), using historical and modern data, suggested that climate change was the most likely driver of the loss of red spruce during the twentieth century as the dominant tree in midelevation forests of the HBEF region. However, climate change is not the only potential explanation for these shifts. For example, elevated acid deposition reduces cold tolerance in red spruce and has been linked to spruce decline at high altitudes (Schaberg et al. 2001) and to declines in sugar maple (Moore and Ouimet 2006) and montane paper birch (*Betula papyrifera*; Halman et al. 2011). Multiple stressors and changes in historical disturbance regimes in the coming century may lead to further declines of these species, whereas others, including beech and red oak, may increase (Mohan et al. 2009). Introduced pests and diseases, both already present (e.g., beech bark disease) and expected in the future (e.g., hemlock woolly adelgid, emerald ash borer), may alter the community dynamics or even remove species

from the forest entirely. Climate change might hasten these invasions or might mediate their effects on forest structure. For example, beech bark disease appears to have increased the susceptibility of American beech to ice storm damage while also leading to high rates of regeneration, which allows the beech to respond rapidly to increased light in gaps (Weeks et al. 2009).

Our detailed monitoring and studies of vegetation at the HBEF have implications for other long-term monitoring programs. At the HBEF, complete inventories of the 13.2-ha reference watershed (watershed 6) that have been conducted every 5 years since 1977 show marked shifts in forest composition (Schwarz et al. 2003, Van Doorn et al. 2011). However, forest aging following logging and hurricane disturbance in the early twentieth century, along with elevated nitrogen and sulfur deposition, ecosystem acidification, and beech bark disease, make it impossible to ascribe the observed changes in forest composition to any one cause in particular. The complexity of vegetation change underscores the need for long-term monitoring in a focused spatial framework. In areas of rugged topography such as the HBEF, monitoring should be focused at or near current transitions between vegetation types (e.g., the ecotone between hardwood and boreal forest zones), and it should be designed to quantify stress responses and competitive interactions among species. Monitoring must also encompass the full range of disturbance histories in an area. Whereas the HBEF represents a relatively narrow range of disturbance histories, the surrounding landscape spans a range of disturbances, including abandoned small-scale agriculture at lower elevations, heavy commercial logging, extensive fires in the more remote mountainous areas in the late nineteenth and early twentieth centuries, and modern exurban development along roads (Figure 5; Vadeboncoeur et al. 2012). There should also be a focus on monitoring regeneration (including seedlings and saplings), because these size classes will be the first to respond to change (Cleavitt et al. 2011). Monitoring should be keyed to disturbance regimes (e.g., forest gaps), which may provide opportunities for new species to colonize. Monitoring should also be sustained to distinguish stable sink populations of seedlings from advancing fronts of an invading species. The long-term monitoring of key variables must also be standardized in order to facilitate comparisons among sites to better distinguish similarities and differences in response to climate change across the landscape.

Conclusions

Whereas the scale and pervasive nature of climate change motivate macroscale approaches that can couple and depict atmospheric and ecosystem processes at regional and global scales, these methodologies may result in an incomplete or inaccurate assessment of climate change effects on ecosystem structure, function, and services at local scales. Because climate change plays out on a complex hydrobiogeochemical template (i.e., the intertwined patterns of soils, vegetation,




Figure 5. A comparison of forest composition of trees cited as boundary markers in early land surveys in central New Hampshire (580–700 m) with modern forest plots representing a range of disturbance histories at similar elevations illustrates how the local effects of climate change on vegetation are often confounded by the interactions of natural and anthropogenic factors that overshadow the effects of changes in climate on ecosystems. The data are the percentage of stems that were greater than 20 centimeters diameter at breast height. Spruce has declined regionally due to clearing, changes in climate and acid deposition, except where selected for by grazing on abandoned lands, while beech has not successfully recolonized abandoned agricultural lands. Maples are more common on the landscape than they were prior to settlement. The data are from Vadeboncoeur and colleagues (2012). Abbreviations: Be, beech; Bi, birch; F, fir; H, hemlock; ha, hectare; m, meters; M, maple; O, other; S, spruce.

and hydrologic flowpaths) and interacts with many human and natural factors at multiple temporal and spatial scales, various effects of climate change cannot be predicted from the broad constraints of temperature and precipitation alone on ecosystem properties.

We argue that long-term integrated studies, such as those conducted over the past 50 years at the HBEF, should be an essential component of climate change research and assessment. A combination of long-term and in-depth measurements is essential for understanding the interplay between climate and forest ecosystem dynamics. At the HBEF, that interplay has produced surprising effects on hydrologic variables (e.g., evapotranspiration, streamflow, soil moisture); changes in phenology and water, carbon, and nitrogen fluxes during critical seasonal transition periods; winter climate change effects on plant and animal community composition

and ecosystem services; and complex interactions with anthropogenic disturbances and land-use changes that influence plant community composition. Further research on how site-specific climate change affects multiple components of ecosystem structure and function should be useful for revealing the key determinants of plant and animal community composition, fluxes of water, element dynamics, and the human valuation and use of ecosystems in many areas of the globe. The results from these detailed studies need to be incorporated into macroscale approaches in an iterative process of modeling, experiments, and long-term monitoring at multiple scales. We suggest that the coordination of long-term research efforts and the development of common approaches will improve our understanding of and response to the overarching challenge that climate change presents to science and society.

Acknowledgments

This research was supported by grants from the National Science Foundation, primarily grant no. DEB 04-23259 to the Hubbard Brook Long Term Ecological Research Network site. The Hubbard Brook Experimental Forest is operated by the Northern Research Station of the US Department of Agriculture Forest Service, in Newtown Square, Pennsylvania, and is a Long Term Ecological Research Network site. This article is a contribution to the Hubbard Brook Ecosystem Study. The authors thank three anonymous reviewers for helpful comments.

References cited

- Albani M, Moorcroft PR, Ellison AM, Orwig DA, Foster DR. 2010. Predicting the impact of hemlock woolly adelgid on carbon dynamics of eastern United States forests. *Canadian Journal of Forest Research* 40: 119–133.
- Auclair AND, Heilman WE, Brinkman B. 2010. Predicting forest dieback in Maine, USA: A simple model based on soil frost and drought. *Canadian Journal of Forest Research* 40: 687–702.
- Beckage B, Osborne B, Gavin DG, Pucko C, Siccama T, Perkins T. 2008. A rapid upward shift of a forest ecotone during 40 years of warming in the Green Mountains of Vermont. *Proceedings of the National Academy of Sciences* 105: 4197–4202.
- Both C. 2006. Climate change and adaptation of annual cycles of migratory birds. *Journal of Ornithology* 147: 68.
- Brooks PD, Grogan P, Templer PH, Groffman PM, Åquist MG, Schimel J. 2011. Carbon and nitrogen cycling in snow-covered environments. *Geography Compass* 5: 682–699.
- Burger C, et al. 2012. Climate change, breeding date and nestling diet: How temperature differentially affects seasonal changes in pied flycatcher diet depending on habitat variation. *Journal of Animal Ecology* 81: 926–936.
- Campbell JL, Mitchell MJ, Groffman PM, Christenson LM, Hardy JP. 2005. Winter in northeastern North America: A critical period for ecological processes. *Frontiers in Ecology and the Environment* 3: 314–322.
- Campbell JL, et al. 2007. Long-Term Trends from Ecosystem Research at the Hubbard Brook Experimental Forest. US Department of Agriculture, Forest Service, Northern Research Station. General Technical Report no. NRS-17.
- Campbell JL, et al. 2009. Consequences of climate change for biogeochemical cycling in forests of northeastern North America. *Canadian Journal of Forest Research* 39: 264–284.
- Campbell JL, Ollinger SV, Flerchinger GN, Wicklein H, Hayhoe K, Bailey AS. 2010. Past and projected future changes in snowpack and soil frost at the Hubbard Brook Experimental Forest, New Hampshire, USA. *Hydrological Processes* 24: 2465–2480.
- Campbell JL, Driscoll CT, Pourmokhtarian A, Hayhoe K. 2011. Stream flow responses to past and projected future changes in climate at the Hubbard Brook Experimental Forest, New Hampshire, USA. *Water Resources Research* 47 (art. W02514). doi:10.1029/2010WR009438
- Christenson LM, Mitchell MJ, Groffman PM, Lovett GM. 2010. Winter climate change implications for decomposition in Northeastern forests: Comparisons of sugar maple litter to herbivore fecal inputs. *Global Change Biology* 16: 2589–2601.
- Cleavitt NL, Fahey TJ, Groffman PM, Hardy JP, Henry KS, Driscoll CT. 2008. Effects of soil freezing on fine roots in a northern hardwood forest. *Canadian Journal of Forest Research* 38: 82–91.
- Cleavitt NL, Fahey TJ, Battles JJ. 2011. Regeneration ecology of sugar maple (*Acer saccharum*): Seedling survival in relation to nutrition, site factors, and damage by insects and pathogens. *Canadian Journal of Forest Research* 42: 235–244.
- Cogbill CV. 2000. Vegetation of the presettlement forests of northern New England and New York. *Rhodora* 102: 250–276.
- Créte M, Ouellet J-P, Lesage L. 2001. Comparative effects on plants of caribou/reindeer, moose and white-tailed deer herbivory. *Arctic* 54: 407–417.
- Duchesne L, Houle D, Côté MA, Logan T. 2009. Modelling the effect of climate on maple syrup production in Québec, Canada. *Forest Ecology and Management* 258: 2683–2689.
- Faison EK, Motzkin G, Foster DR, McDonald JE. 2010. Moose foraging in the temperate forests of southern New England. *Northeastern Naturalist* 17: 1–8.
- Fitzhugh RD, Driscoll CT, Groffman PM, Tierney GL, Fahey TJ, Hardy JP. 2001. Effects of soil freezing disturbance on soil solution nitrogen, phosphorus, and carbon chemistry in a northern hardwood ecosystem. *Biogeochemistry* 56: 215–238.
- Fitzhugh RD, Likens GE, Driscoll CT, Mitchell MJ, Groffman PM, Fahey TJ, Hardy JP. 2003. Role of soil freezing events in interannual patterns of stream chemistry at the Hubbard Brook Experimental Forest, New Hampshire. *Environmental Science and Technology* 37: 1575–1580.
- Greene DF, Johnson EA. 1997. Secondary dispersal of tree seeds on snow. *Journal of Ecology* 85: 329–340.
- Groffman PM, Driscoll CT, Fahey TJ, Hardy JP, Fitzhugh RD, Tierney GL. 2001. Colder soils in a warmer world: A snow manipulation study in a northern hardwood forest ecosystem. *Biogeochemistry* 56: 135–150.
- Groffman PM, Driscoll CT, Eagar C, Fisk MC, Fahey TJ, Holmes RT, Likens GE, Pardo LH. 2004. A new governance structure for the Hubbard Brook Ecosystem Study. *Bulletin of the Ecological Society of America* 85: 5–6.
- Groffman PM, Hardy JP, Fisk MC, Fahey JT, Driscoll CT. 2009. Climate variation and soil carbon and nitrogen cycling processes in a northern hardwood forest. *Ecosystems* 12: 927–943.
- Groffman PM, Hardy JP, Fashu-Kanu S, Driscoll CT, Cleavitt NL, Fahey TJ, Fisk MC. 2010. Snow depth, soil freezing and nitrogen cycling in a northern hardwood forest landscape. *Biogeochemistry* 102: 223–238.
- Halman JM, Schaberg PG, Hawley GJ, Hansen CF. 2011. Potential role of soil calcium in recovery of paper birch following ice storm injury in Vermont, USA. *Forest Ecology and Management* 261: 1539–1545.
- Hamburg SP, Cogbill CV. 1988. Historical decline of red spruce populations and climatic warming. *Nature* 331: 428–431.
- Hamburg SP, Vadeboncoeur MA, Richardson AD, Bailey AS. 2012. Climate change at the ecosystem scale: A 50-year record in New Hampshire. *Climatic Change*. (14 September 2012; www.springerlink.com/content/77481321304k5311) doi:10.1007/s10584-012-0517-2
- Hayhoe K, et al. 2007. Past and future changes in climate and hydrological indicators in the US Northeast. *Climate Dynamics* 28: 381–407.
- Hentschel K, Borken W, Zuber T, Bogner C, Howe B, Matzner E. 2009. Effects of soil frost on nitrogen net mineralization, soil solution chemistry and seepage losses in a temperate forest soil. *Global Change Biology* 15: 825–836.
- Holmes RT, Sherry TW. 2001. Thirty-year bird population trends in an unfragmented temperate deciduous forest: Importance of habitat change. *The Auk* 118: 589–609.
- Holmes RT, Sherry TW, Marra PP, Petit KE. 1992. Multiple brooding and productivity of a Neotropical migrant, the black-throated blue warbler (*Dendroica caerulescens*), in an unfragmented temperate forest. *The Auk* 109: 321–333.
- Huntington TG, Richardson AD, McGuire KJ, Hayhoe K. 2009. Climate and hydrological changes in the northeastern United States: Recent trends and implications for forested and aquatic ecosystems. *Canadian Journal of Forest Research* 39: 199–212.
- Ibáñez I, Primack RB, Miller-Rushing AJ, Ellwood E, Higuchi H, Lee SD, Kobori H, Silander JA. 2010. Forecasting phenology under global warming. *Philosophical Transactions of the Royal Society B* 365: 3247–3260.
- Iverson LR, Prasad AM. 2001. Potential changes in tree species richness and forest community types following climate change. *Ecosystems* 4: 186–199.
- Iverson LR, Prasad AM, Matthews SN, Peters MP. 2011. Lessons learned while integrating habitat, dispersal, disturbance, and life-history traits into species habitat models under climate change. *Ecosystems* 14: 1005–1020.

- Jeong S-J, Medvigy D, Shevliakova E, Malyshev S. 2012. Uncertainties in terrestrial carbon budgets related to spring phenology. *Journal of Geophysical Research* 117 (art. G01030).
- Kreyling J. 2010. Winter climate change: A critical factor for temperate vegetation performance. *Ecology* 91: 1939–1948.
- Leak WB. 1978. Relationship of Species and Site Index to Habitat in the White Mountains of New Hampshire. US Department of Agriculture, Forest Service. Research Paper no. NE-397.
- Likens GE, Bormann FH. 1995. *Biogeochemistry of a Forested Ecosystem*, 2nd ed. Springer.
- Magnuson JJ, et al. 2000. Historical trends in lake and river ice cover in the Northern Hemisphere. *Science* 289: 1743–1746.
- Mitchell MJ, Likens GE. 2011. Watershed sulfur biogeochemistry: Shift from atmospheric deposition dominance to climatic regulation. *Environmental Science and Technology* 45: 5267–5271.
- Mitchell MJ, Driscoll CT, Kahl JS, Likens GE, Murdoch PS, Pardo LH. 1996. Climatic control of nitrate loss from forested watersheds in the northeast United States. *Environmental Science and Technology* 30: 2609–2612.
- Mohan JE, Cox RM, Iverson LR. 2009. Composition and carbon dynamics of forests in northeastern North America in a future, warmer world. *Canadian Journal of Forest Research* 39: 213–230.
- Moore JD, Ouimet R. 2006. Ten-year effect of dolomitic lime on the nutrition, crown vigor, and growth of sugar maple. *Canadian Journal of Forest Research* 36: 1834–1841.
- Morrison SF, Forbes GJ, Young SJ, Lusk S. 2003. Within-yard habitat use by white-tailed deer at varying winter severity. *Forest Ecology and Management* 172: 173–182.
- Muller RN, Bormann FH. 1976. Role of *Erythronium americanum* Ker. in energy flow and nutrient dynamics of a northern hardwood forest ecosystem. *Science* 193: 1126–1128.
- Paradis A, Elkinton J, Hayhoe K, Buonaccorsi J. 2008. Role of winter temperature and climate change on the survival and future range expansion of the hemlock woolly adelgid (*Adelges tsugae*) in eastern North America. *Mitigation and Adaptation Strategies for Global Change* 13: 541–554.
- Polgar CA, Primack RB. 2011. Leaf-out phenology of temperate woody plants: From trees to ecosystems. *New Phytologist* 191: 926–941.
- Richardson AD, Bailey AS, Denny EG, Martin CW, O'Keefe J. 2006. Phenology of a northern hardwood forest canopy. *Global Change Biology* 12: 1174–1188.
- Rodenhouse NL, Christenson LM, Parry D, Green LE. 2009. Climate change effects on native fauna of northeastern forests. *Canadian Journal of Forest Research* 39: 249–263.
- Roderick ML, Hobbins MT, Farquhar GD. 2009. Pan evaporation trends and the terrestrial water balance. I. Principles and observations. *Geography Compass* 3: 746–760.
- Samson C, Huot J. 1998. Movements of female black bears in relation to landscape vegetation type in southern Quebec. *Journal of Wildlife Management* 62: 718–727.
- Schaberg PG, DeHayes DH, Hawley GJ. 2001. Anthropogenic calcium depletion: A unique threat to forest ecosystem health? *Ecosystem Health* 7: 214–228.
- Schwarz PA, Fahey TJ, Dawson TE. 1997. Seasonal air and soil temperature effects on photosynthesis in red spruce (*Picea rubens*) saplings. *Tree Physiology* 17: 187–194.
- Schwarz PA, Fahey TJ, McCulloch CE. 2003. Factors controlling spatial variation of tree species abundance in a forested landscape. *Ecology* 84: 1862–1878.
- Skinner CB, DeGaetano AT, Chabot BF. 2010. Implications of twenty-first century climate change on northeastern United States maple syrup production: Impacts and adaptations. *Climatic Change* 100: 685–702.
- Steiger R, Mayer M. 2008. Snowmaking and climate change: Future options for snow production in Tyrolean ski resorts. *Mountain Research and Development* 28: 292–298.
- Subak S. 2003. Effects of climate on variability in Lyme disease incidence in the northeastern United States. *American Journal of Epidemiology* 157: 531–538.
- Templer PH, Schiller AF, Fuller NW, Soccia AM, Campbell JL, Drake JE, Kunz TH. 2012. Impact of a reduced winter snowpack on litter arthropod abundance and diversity in a northern hardwood forest ecosystem. *Biology and Fertility of Soils* 48: 413–424.
- Tierney GL, Fahey TJ, Groffman PM, Hardy JP, Fitzhugh RD, Driscoll CT. 2001. Soil freezing alters fine root dynamics in a northern hardwood forest. *Biogeochemistry* 56: 175–190.
- Vadeboncoeur MA, Hamburg SP, Cogbill CV, Sugimura WY. 2012. A comparison of presettlement and modern forest composition in central New Hampshire. *Canadian Journal of Forest Research* 42: 190–202.
- Van Doorn NS, Battles JJ, Fahey TJ, Siccama TG, Schwarz PA. 2011. Links between biomass and tree demography in a northern hardwood forest: A decade of stability and change in Hubbard Brook Valley, New Hampshire. *Canadian Journal of Forest Research* 41: 1369–1379.
- Warren DR, Likens GE, Buso DC, Kraft CE. 2008. Status and distribution of fish in an acid-impacted watershed of the northeastern United States (Hubbard Brook, NH). *Northeastern Naturalist* 15: 375–390.
- Weeks BC, Hamburg SP, Vadeboncoeur MA. 2009. Ice storm effects on the canopy structure of a northern hardwood forest after eight years. *Canadian Journal of Forest Research* 39: 1475–1483.
- Whitlaw HA, Lankester MW. 1994. The co-occurrence of moose, white-tailed deer, and *Parelaphostrongylus tenuis* in Ontario. *Canadian Journal of Zoology* 72: 819–825.
- Zak DR, Groffman PM, Pregitzer KS, Christensen S, Tiedje JM. 1990. The vernal dam: Plant microbe competition for nitrogen in northern hardwood forests. *Ecology* 71: 651–656.

Peter M. Groffman (groffmanp@caryinstitute.org) is a senior scientist and Jennifer L. Morse is a postdoctoral researcher at the Cary Institute of Ecosystem Studies, in Millbrook, New York. Lindsey E. Rustad and John L. Campbell are research ecologists with the US Department of Agriculture (USDA) Forest Service's Northern Research Station, in Durham, New Hampshire. Pamela H. Templer is an associate professor in the Department of Biology at Boston University, in Boston, Massachusetts. Lynn M. Christenson is an assistant professor in the Department of Biology at Vassar College, in Poughkeepsie, New York. Nina K. Lany is a graduate student in the Department of Biological Sciences at Dartmouth College, in Hanover, New Hampshire. Anne M. Soccia is a graduate student in the Department of Biology at Boston University, in Boston, Massachusetts. Matthew A. Vadeboncoeur is a graduate student at the Complex Systems Research Center at the Institute for the Study of Earth, Oceans, and Space at the University of New Hampshire, in Durham. Paul G. Schaberg is a research plant physiologist and Linda H. Pardo is an environmental engineer with the USDA Forest Service's Northern Research Station, in Burlington, Vermont. Geoffrey F. Wilson is the education director at the Hubbard Brook Research Foundation, in Thornton, New Hampshire. Charles T. Driscoll and Chris E. Johnson are professors in the Department of Civil and Environmental Engineering at Syracuse University, in Syracuse, New York. Timothy J. Fahey is a professor in the Department of Natural Resources at Cornell University, in Ithaca, New York. Melany C. Fisk is an associate professor in the Department of Zoology at Miami University, in Oxford, Ohio. Christine L. Goodale is an associate professor in the Department of Ecology and Evolutionary Biology Cornell University, in Ithaca, New York. Mark B. Green is an assistant professor at the Center for the Environment at Plymouth State University, in Plymouth, New Hampshire, and a research hydrologist with the USDA Forest Service's Northern Research Station, in Durham, New Hampshire. Steven P. Hamburg is the chief scientist for the Environmental Defense Fund, in Boston, Massachusetts. Myron J. Mitchell is a distinguished professor in the State University of New York's College of Environmental Science and Forestry, in Syracuse and Nicholas L. Rodenhouse is a professor in the Department of Biological Sciences at Wellesley College, in Wellesley, Massachusetts.

