

Portland State University

PDXScholar

---

Chemistry Faculty Publications and  
Presentations

Chemistry

---

5-2001

## Trimethylsulfonium Methanesulfonate

Frank R. Fronczek  
*Louisiana State University*

Rolanda J. Johnson

Robert M. Strongin  
*Portland State University, strongin@pdx.edu*

Follow this and additional works at: [https://pdxscholar.library.pdx.edu/chem\\_fac](https://pdxscholar.library.pdx.edu/chem_fac)

 Part of the [Organic Chemistry Commons](#)

Let us know how access to this document benefits you.

---

### Citation Details

Fronczek, F., Johnson, R., & Strongin, R. (2001). Trimethylsulfonium methanesulfonate. *Acta Crystallographica. Section E, Structure Reports*, 57 (Pt 5), 447-448.

This Article is brought to you for free and open access. It has been accepted for inclusion in Chemistry Faculty Publications and Presentations by an authorized administrator of PDXScholar. Please contact us if we can make this document more accessible: [pdxscholar@pdx.edu](mailto:pdxscholar@pdx.edu).

Acta Crystallographica Section E

**Structure Reports****Online**

ISSN 1600-5368

## Trimethylsulfonium methanesulfonate


Frank R. Fronczek, Rolanda J. Johnson and Robert M. Strongin

*Acta Cryst.* (2001). **E57**, o447–o448

Copyright © International Union of Crystallography

Author(s) of this paper may load this reprint on their own web site or institutional repository provided that this cover page is retained. Reproduction of this article or its storage in electronic databases other than as specified above is not permitted without prior permission in writing from the IUCr.

For further information see <http://journals.iucr.org/services/authorrights.html>



*Acta Crystallographica Section E: Structure Reports Online* is the IUCr's highly popular open-access structural journal. It provides a simple and easily accessible publication mechanism for the growing number of inorganic, metal-organic and organic crystal structure determinations. The electronic submission, validation, refereeing and publication facilities of the journal ensure very rapid and high-quality publication, whilst key indicators and validation reports provide measures of structural reliability. The average publication time is less than one month. Articles are published in a short-format style with enhanced supplementary materials. Each publication consists of a complete package – the published article, HTML and PDF supplements, CIF, structure factors, graphics, and any other submitted supplementary files.

Crystallography Journals **Online** is available from [journals.iucr.org](http://journals.iucr.org)

## Trimethylsulfonium methanesulfonate

Frank R. Fronczek,\* Rolanda J. Johnson and Robert M. Strongin

Department of Chemistry, Louisiana State University, Baton Rouge, LA 70803-1804, USA

Correspondence e-mail: fronz@chxray1.chem.lsu.edu

In the title compound,  $C_3H_9S^+ \cdot CH_3O_3S^-$ , a thermal decomposition product of dimethyl sulfoxide, both cation and anion lie on mirror planes. In the cation, the S atom lies 0.792 (2) Å out of the plane defined by the three C atoms, with S—C distances of 1.781 (2) and 1.786 (3) Å. In the anion, the S—O distances are 1.4556 (14) and 1.4646 (19) Å, and the S—C distance is 1.759 (3) Å.

Received 5 April 2001  
Accepted 20 April 2001  
Online 26 April 2001


## Key indicators

Single-crystal X-ray study  
T = 120 K  
Mean  $\sigma(S-O) = 0.002$  Å  
R factor = 0.039  
wR factor = 0.089  
Data-to-parameter ratio = 17.7

For details of how these key indicators were automatically derived from the article, see <http://journals.iucr.org/e>.

## Comment

We have been studying the structure and mechanism of formation of colored products derived from resorcinarene macrocycles upon heating in dimethyl sulfoxide (DMSO) (Lewis *et al.*, 1997, 2000; Davis *et al.*, 1999). We isolated the title compound, (I), a decomposition product formed by prolonged heating of a solution of macrocycle in DMSO, and determined its structure to ascertain its identity. While 38 salts of the trimethylsulfonium ion and 62 salts of the methylsulfonate anion are present in the Cambridge Structural Database (December 2000, 224400 entries; Allen & Kennard, 1993), the structure of the title compound has not been previously reported. Decomposition of DMSO or its complexes to form trimethylsulfonium methanesulfonate has been previously reported as a result of heating (Banci, 1967; Arsenin *et al.*, 1988) and  $\gamma$  irradiation (Gutierrez *et al.*, 1977).



Both cation and anion lie across crystallographic mirrors. In the cation, the S atom lies 0.792 (2) Å out of the plane defined by the three C atoms. Geometric parameters (Table 1) are normal. Most of the H atoms are involved in C—H $\cdots$ O hydrogen bonding (Table 2)

## Experimental

For the preparation of (I), the tetramethylresorc[4]arene 2,8,14,20-tetramethylpentacyclo[19.3.1.13,7.19,13.115,19]octacos-1(25),3,5,7(28),9,11,13(27),15,17,19(26),21,23-dodecaene-4,6,10,12,16,18,22,24-octol (250 mg, 0.46 mmol) was placed in a sealed tube along with 12 ml of DMSO and 3 ml of water. The mixture was heated at 493 K for 36 h. An aliquot was removed, and the solvent evaporated, yielding colorless crystals of the title compound.

Crystal data

$C_3H_9S^+ \cdot CH_3O_3S^-$   
 $M_r = 172.26$   
 Orthorhombic,  $Pnma$   
 $a = 12.6157(4) \text{ \AA}$   
 $b = 8.2419(4) \text{ \AA}$   
 $c = 7.5397(8) \text{ \AA}$   
 $V = 783.96(9) \text{ \AA}^3$   
 $Z = 4$   
 $D_x = 1.459 \text{ Mg m}^{-3}$

Mo  $K\alpha$  radiation  
 Cell parameters from 7642 reflections  
 $\theta = 2.5\text{--}30.0^\circ$   
 $\mu = 0.62 \text{ mm}^{-1}$   
 $T = 120 \text{ K}$   
 Plate, colorless  
 $0.12 \times 0.10 \times 0.02 \text{ mm}$

Data collection

KappaCCD diffractometer (with Oxford Cryosystems Cryostream cooler)  
 $\omega$  scans with  $\kappa$  offsets  
 Absorption correction: multi-scan (HKL SCALEPACK; Otwinowski & Minor, 1997)  
 $T_{\min} = 0.936, T_{\max} = 0.988$

7642 measured reflections  
 1222 independent reflections  
 838 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.047$   
 $\theta_{\max} = 30.1^\circ$   
 $h = -17 \rightarrow 17$   
 $k = -11 \rightarrow 11$   
 $l = -10 \rightarrow 10$

Refinement

Refinement on  $F^2$   
 $R[F^2 > 2\sigma(F^2)] = 0.039$   
 $wR(F^2) = 0.089$   
 $S = 1.02$   
 1222 reflections  
 69 parameters  
 Only coordinates of H atoms refined

$w = 1/[\sigma^2(F_o^2) + (0.0326P)^2 + 0.2622P]$   
 where  $P = (F_o^2 + 2F_c^2)/3$   
 $(\Delta/\sigma)_{\max} < 0.001$   
 $\Delta\rho_{\max} = 0.40 \text{ e \AA}^{-3}$   
 $\Delta\rho_{\min} = -0.37 \text{ e \AA}^{-3}$   
 Extinction correction: SHELXL97  
 Extinction coefficient: 0.0067 (18)

Table 1

Selected geometric parameters ( $\text{\AA}, ^\circ$ ).

S1—O1	1.4646 (19)	S2—C2	1.786 (3)
S1—O2	1.4556 (14)	S2—C3	1.781 (2)
S1—C1	1.759 (3)		
O2—S1—O1	112.20 (7)	O2—S1—C1	106.11 (8)
O2 <sup>i</sup> —S1—O2	113.45 (12)	C3—S2—C3 <sup>ii</sup>	101.31 (15)
O1—S1—C1	106.11 (13)	C3—S2—C2	102.00 (10)

Symmetry codes: (i)  $x, \frac{1}{2} - y, z$ ; (ii)  $x, \frac{3}{2} - y, z$ .

Table 2

Hydrogen-bonding geometry ( $\text{\AA}, ^\circ$ ).

$D-H \cdots A$	$D-H$	$H \cdots A$	$D \cdots A$	$D-H \cdots A$
C1—H1B $\cdots$ O2 <sup>i</sup>	0.95 (2)	2.54 (2)	3.463 (2)	163.5 (18)
C2—H2A $\cdots$ O1 <sup>ii</sup>	1.06 (3)	2.41 (3)	3.382 (4)	152 (2)
C3—H3A $\cdots$ O1 <sup>ii</sup>	0.97 (2)	2.48 (2)	3.388 (3)	156.0 (19)
C3—H3B $\cdots$ O2 <sup>iii</sup>	0.95 (2)	2.44 (2)	3.279 (3)	147.3 (19)
C3—H3C $\cdots$ O1	0.92 (3)	2.41 (3)	3.280 (2)	157.1 (19)

Symmetry codes: (i)  $\frac{3}{2} - x, -y, z - \frac{1}{2}$ ; (ii)  $1 - x, \frac{1}{2} + y, -z$ ; (iii)  $1 - x, \frac{1}{2} + y, 1 - z$ .

The coordinates of H atoms were refined, while their isotropic displacement parameters were assigned as  $U_{\text{iso}} = 1.5U_{\text{eq}}$  of the attached atom.




Figure 1

The atom-numbering scheme for (I) with ellipsoids at the 50% probability level.

Data collection: COLLECT (Nonius, 2000); cell refinement: DENZO and SCALEPACK; data reduction: DENZO and SCALEPACK (Otwinowski & Minor, 1997); program(s) used to solve structure: SIR (Altomare *et al.*, 1994); program(s) used to refine structure: SHELXL97 (Sheldrick, 1997); molecular graphics: ORTEP-3 (Farrugia, 1997); software used to prepare material for publication: SHELXL97.

The purchase of the diffractometer was made possible by grant No. LEQSF(1999–2000)-ESH-TR-13, administered by the Louisiana Board of Regents.

References

- Allen, F. H. & Kennard, O. (1993). *Chem. Des. Autom. News*, **8**, 1, 31–37.  
 Altomare, A., Casciaro, G., Giacovazzo, C. & Guagliardi, A. (1994). *J. Appl. Cryst.* **27**, 435.  
 Arsenin, K. I., Malinko, L. A., Sheka, I. A., Pishchai, I. Y., Tikhonova, R. V., Kolotio, L. V. & Antishko, A. N. (1988). *Zh. Obshch. Khim.* **58**, 2102–2109.  
 Banci, F. (1967). *Ann. Chim.* **57**, 549–554.  
 Davis, C. J., Lewis, P. T., McCarroll, M. E., Reed, M. W., Cueto, R. & Strongin, R. M. (1999). *Org. Lett.* **1**, 331–334.  
 Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.  
 Gutierrez, M. C., Barra, R., Sanz, H. & Parellada, R. (1977). *An. Quim.* **73**, 520–526.  
 Lewis, P. T., Davis, C. J., Cabell, L. A., He, M., Read, M. W., McCarroll, M. E. & Strongin, R. M. (2000). *Org. Lett.* **2**, 589–592.  
 Lewis, P. T., Davis, C. J., Saraiva, M. C., Treleavan, D., McCarley, T. & Strongin, R. M. (1997). *J. Org. Chem.* **62**, 6110–6111.  
 Nonius (2000). COLLECT. Nonius BV, Delft, The Netherlands.  
 Otwinowski, Z. & Minor, W. (1997). *Methods Enzymol.* **276**, 307–326.  
 Sheldrick, G. (1997). SHELXL97. University of Göttingen, Germany.