

6-2013

## Dynamic Modeling of Nonmedical Opioid Initiation: Epidemic and Access

Alexandra Nielsen  
*Portland State University*

Teresa D. Schmidt  
*Portland State University*

Dennis McCarty  
*Oregon Health & Science University*

Wayne W. Wakeland  
*Portland State University, wakeland@pdx.edu*

Let us know how access to this document benefits you.

Follow this and additional works at: [https://pdxscholar.library.pdx.edu/sysc\\_fac](https://pdxscholar.library.pdx.edu/sysc_fac)

 Part of the [Chemical Actions and Uses Commons](#), and the [Pharmaceutical Preparations Commons](#)

### Citation Details

Nielsen, A., Schmidt, T.D., McCarty, D., Wakeland, W. (June 2013). Dynamic modeling of nonmedical opioid initiation: Epidemic and access. Poster presentation at the College on Problems of Drug Dependents, 75th Annual Meeting, San Diego, CA.

This Poster is brought to you for free and open access. It has been accepted for inclusion in Systems Science Faculty Publications and Presentations by an authorized administrator of PDXScholar. For more information, please contact [pdxscholar@pdx.edu](mailto:pdxscholar@pdx.edu).

# Dynamic Modeling of Nonmedical Opioid Initiation: Epidemic and Access

Alexandra Nielsen BS, Portland State University, Teresa D. Schmidt MS, Portland State University, Dennis McCarty PhD, Oregon Health and Science University, Wayne Wakeland PhD, Portland State University

We report development of a systems level dynamic model of initiation and nonmedical use of pharmaceutical opioids in the US. The model calibrated to 1995-2005 National Survey of Drug Use and Health (NSDUH) data, predicts 2006-2011 data well. Preliminary findings indicate that interventions which reduce the perceived attractiveness of opioids for recreational use may be able to reduce initiation and nonmedical use most significantly, while supply restriction effected through drug take back days and prescribing changes may have more modest effects. We argue that system dynamics is an effective approach for evaluating potential interventions to this complex system where the use of pharmaceutical opioids to treat pain is fraught with undesirable distal outcomes.

## Background


The number of overdose deaths involving opioids tripled between 1999 and 2006 in the US, rising to 14,800 in 2008 (Warner et al. 2011). As evidenced by the high fraction of opioid overdose decedents without prescriptions (Hall et al. 2008), nonmedical use of pharmaceutical opioids plays a significant role in the prevalence of overdose deaths. Leftover opioid prescriptions are likely involved in much of this diversion (Compton and Volkow 2006). Studies suggest that there is a large reservoir of unused opioids stored in homes (Bates et al. 2011; Gilson 2012), and the high fraction of individuals receiving drugs for free from friends and family as reported in the NSDUH (SAMHSA 2012) is likely to be strongly correlated with the size of this reservoir.

## Methods

Drawing from empirical data and a panel of experts, a system dynamics simulation was developed to reproduce historical trends in nonmedical use of pharmaceutical opioids in the United States. Data to support assumptions and model parameters were drawn from publicly available sources in the literature. Data on initiation and nonmedical use were obtained from the NSDUH for the years 1995 to 2011. System dynamics is a simulation method in which complex relationships and feedback loops are specified and mathematically formalized. The resultant set of differential equations was calibrated to replicate opioid use data from 1995-2005, used to predict behavior from 2006 to 2011, and to evaluate policy interventions.

## Results

The model contains 5 state variables, 13 exogenous parameter variables, 21 endogenous (calculated) parameters and 10 rates of change. Three principal feedback loops (a peer initiation epidemic loop, a global availability loop, and a personal accessibility loop) contribute to the nonlinear growth patterns in nonmedical opioid initiation and use. Peer initiation is modeled as the infection of a susceptible population by peers. Global availability of opioids for nonmedical use depends on the number of current opioid users and how much free leftover medicine they obtain from prescription holders. When availability diminishes, reduced personal accessibility requires transitions



to paying for opioids. Compelling susceptible non-users to resist initiation (e.g. Gosin et al. 2003) was more effective in reducing nonmedical use than compelling prescription holders not to share their medicines (e.g. FDA 2013), and reducing global availability through prescription take back events (e.g. DEA 2012).

## Discussion

The relative impact of the three interventions simulated can not be explained by feedback alone as all three interventions are amplified by positive feedback loops. The infectivity intervention reduces the initiate rate by 25%, and this reduction is immediately amplified by feedback. The willingness to share intervention impacts both the infection loop and the global availability loop, but the 25% reduction is indirect. Reducing by 25% the number of people who are willing to share prescriptions only reduces the likelihood that a potential initiator will know at least one prescription holder by a small percentage. This likelihood is based on a binomial distribution and is large when a large fraction of the population has prescriptions. So the impact on initiation and global supply is much smaller than 25%. This smaller reduction is amplified by positive feedback, which suggests that limiting sharing is an intervention with leverage. Supply reduction through prescription take back days is the least effective strategy primarily because the reservoir of leftover medicines is so large that the reduction implemented did not cause the system to be supply constrained.

## Conclusion

The current model offers a formalized model of a common pathway to nonmedical opioid initiation and provides a tool for comparing the impact of multiple policy interventions. System dynamics modeling brings insights on the global dynamics of nonmedical opioid use and can be used to inform policy interventions to ameliorate the associated public health problems.

## References

Bates, C. et al., 2011. Overprescription of postoperative narcotics: a look at postoperative pain medication delivery, consumption and disposal in urological practice. *The Journal of urology*, 185(2), pp.551-555.  
 Compton, W.M. & Volkow, N.D., 2006. Major increases in opioid analgesic abuse in the United States: concerns and strategies. *Drug and alcohol dependence*, 81(2), pp.103-108.  
 Drug Enforcement Administration, 2012. DEA's Fifth National Prescription Drug Take-Back Day Results in Another Big Haul. *News Release*. Available at: [www.justice.gov/dea/docs/results\\_final.pdf](http://www.justice.gov/dea/docs/results_final.pdf) [Accessed May 3, 2013].  
 Food and Drug Administration, 2013. *Extended-release (ER) and long-acting (LA) opioid analgesic risk evaluation and mitigation strategy (REMS)*. Available at: <http://www.fda.gov/drugs/drugsafety/informationbydrugclass/ucm163647.htm> [Accessed May 3, 2013].  
 Gilson, A., 2012. Personal Communication: Findings of Madison WI Drug Take-Back Day.  
 Gosin, M., Marsiglia, F.F. & Hecht, M.L., 2003. Keepin'it REAL: a drug resistance curriculum tailored to the strengths and needs of pre-adolescents of the southwest. *Journal of Drug Education*, 33(2), pp.119-142.  
 Hall, A.J. et al., 2008. Patterns of abuse among unintentional pharmaceutical overdose fatalities. *JAMA: the journal of the American Medical Association*, 300(22), pp.2613-2620.  
 Substance Abuse and Mental Health Services Administration, 2012. Results from the 2011 National Survey on Drug Use and Health: Summary of National Findings. Available at: [http://www.samhsa.gov/data/2k11/WEB\\_SR\\_088/WEB\\_SR\\_088.pdf](http://www.samhsa.gov/data/2k11/WEB_SR_088/WEB_SR_088.pdf) [Accessed February 26, 2013].  
 Warner, M. et al., 2011. *Drug Poisoning Deaths in the United States, 1980-2008*, Centers for Disease Control and Prevention.

## Support

This work is supported by NIDA grant 5R21DA031361-02. The authors report no conflicts of interest.