

Portland State University

PDXScholar

Regional Research Institute

School of Social Work

4-2020

Oregon Zero Suicide Implementation Assessment Instrument, v.1.0

Karen Cellarius

Portland State University, cellark@pdx.edu

Meghan Crane

Oregon Health Authority

Follow this and additional works at: https://pdxscholar.library.pdx.edu/rri_facpubs



Part of the [Social Work Commons](#)

Let us know how access to this document benefits you.

Citation Details

Cellarius, K., Crane, M. (2019). Oregon Zero Suicide Implementation Assessment Instrument, v.1.0. Portland, OR: Portland State University.

This Report is brought to you for free and open access. It has been accepted for inclusion in Regional Research Institute by an authorized administrator of PDXScholar. Please contact us if we can make this document more accessible: pdxscholar@pdx.edu.

Oregon Zero Suicide Implementation Assessment Instrument, v.1.0

Developed by the Oregon Health Authority & Portland State University for the GLS Youth Suicide Prevention Project

Background:

This implementation self-assessment and the accompanying web survey were adapted for the Oregon Community Collaboration Initiative (OCCI) by Portland State University in collaboration with the OHA GLS Youth Suicide Prevention staff. The assessment is based on three Zero Suicide resources available at <http://zerosuicide.org/>.

- **The Organizational Self-Study** is a questionnaire about the extent to which each component of the Zero Suicide approach is in place at a single organization. Zero Suicide recommends completing this self-study at the start of an organization's Zero Suicide initiative, then every 12 months after that as a measure of fidelity to the model. **The self-study questions serve as the basis for this Oregon Zero Suicide Implementation Assessment and have been reformulated as indicators.** The response options (or anchors) for each question are included in the grid to define the level of implementation for each indicator.
- **The Data Elements Worksheet** contains primary and supplemental measures recommended for behavioral health care organizations to strive for to maintain fidelity to a comprehensive suicide care model. The supplemental measures are clinically significant but may be much harder to measure than the primary measures. Zero Suicide recommends reviewing these data elements every three months in order to determine areas for improvement. **Starting with element #3 (Identify) of this implementation assessment, these data points are requested for each relevant indicator as documentation for the rank awarded.**
- **The Work Plan Template** outlines recommended steps for implementing the seven elements of Zero Suicide. **The completion dates of specific steps in this template can be documented in the Comment section for each relevant indicator to verify any change in indicator score over time.**

OHA is using this implementation assessment to track change over time related to suicide prevention efforts among organizations participating in OHA-sponsored Zero Suicide Academies in Oregon and subsequent Zero Suicide Community of Practice Conference Calls. Funding to develop this instrument was provided by SAMHSA Garret Lee Smith Youth Suicide Prevention Grant (Grant # 1U79SM061759-01) awarded to the Oregon Health Authority.

For more information on:

--Zero Suicide, visit <http://zerosuicide.org/>

--The OCCI project, contact Megan Crane, OHA Zero Suicide Coordinator in the Oregon Health Authority's Injury and Violence Prevention Section at MEGHAN.CRANE@dhsosha.state.or.us

--The study being conducted using this instrument, contact Karen Cellarius, Senior Research Associate, Portland State University Regional Research Institute for Human Services at cellark@pdx.edu

Suggested citation:

Cellarius, K., Crane, M. (2019). Oregon Zero Suicide Implementation Assessment Instrument, v.1.0. Portland, OR: Portland State University.

Overview of the Elements of Zero Suicide

Element #1: Lead

Create a leadership-driven, safety-oriented culture committed to dramatically reducing suicide among people under care. Include suicide attempt and loss survivors in leadership and planning roles.

Element #2: Train

Develop a competent, confident and caring workforce.

Element #3: Identify

Systematically identify and assess suicide risk among people receiving care.

Element #4: Engage

Ensure every person has a suicide care management plan, or pathway to care, that is both timely and adequate to meet patient needs.

Element #5: Treat

Use effective, evidence-based treatments that directly target suicidality.

Element #6: Transition

Provide continuous contact and support, especially after acute care.

Element #7: Improve

Apply a data-driven quality improvement approach to inform system changes that will lead to improved patient outcomes and better care for those at risk.

General Scale to Implementation Ratings¹:

Anchors, or specific expectations, are included for most components following this range. For comparable pre-post ratings, use the specific definitions for each indicator on pages 5-14.

Rating	Description
1	Routine care or care as usual for this item. The organization has not yet focused specifically on developing or embedding a suicide care approach for this activity.
2	Initial actions toward improvement taken for this item. The organization has taken some preliminary or early steps to focus on improving suicide care.
3	Several steps towards improvement made for this item. The organization has made several steps towards advancing an improved suicide approach.
4	Near comprehensive practices in place for this item. The organization has significantly advanced its suicide care approach.
5	Comprehensive practices in place for this item. The organization has embedded suicide care in its approach and now relies on monitoring and maintenance to ensure sustainability and continuous quality improvement.

¹ Zero Suicide Organizational Self-Study, 1/11/17, page 2

Quick Rating Sheet for Zero Suicide Elements and their Indicators

Instructions: Choose a rating for each indicator on a scale of 1-5 (see definitions below) that best reflects the current situation at the health care entity where Zero Suicide is being implemented. When in doubt, review the specific definition and anchors detailed in the following pages. Finalize the clinic score based on a review of the specific indicators and a follow-up discussion with other on-site staff. Document your logic for the final score in the comments section under each indicator on the following pages.

Scale (For comparable pre-post ratings, use the specific definitions for each indicator on pages 5-14):

1=Routine care or care as usual. The organization has not yet focused specifically on developing or embedding a suicide care approach for this activity.

2=Initial actions toward improvement taken. The organization has taken some preliminary or early steps to focus on improving suicide care.

3=Several steps towards improvement made. The organization has made several steps towards advancing an improved suicide approach.

4=Near comprehensive practices in place. The organization has significantly advanced its suicide care approach.

5=Comprehensive practices in place. The organization has embedded suicide care in its approach and now relies on monitoring and maintenance to ensure sustainability and continuous quality improvement.

INDICATOR	Preliminary Rating	Final Rating
Element #1: Lead		
Leadership-Driven, Safety Oriented Culture		
Written Policies		
Documentation		
Training		
Staffing		
Roles for Survivors		
<i>Subtotal</i>		
Element #1 Average Score (Subtotal/6)		
Element #2: Train		
Workforce Confidence		
Non-Clinical Staff		
Clinical Staff		
<i>Subtotal</i>		
Element #2 Average Score (Subtotal/3)		
Element #3: Identify		
Screening Policies		
Screening Protocols		
Assessment Protocols		
<i>Subtotal</i>		
Element #3 Average Score (Subtotal/3)		
Element #4: Engage		
Pathway to Care		
Collaborative Safety Planning		

INDICATOR	Preliminary Rating	Final Rating
Collaborative Restriction of Access Lethal Means		
<i>Subtotal</i>		
Element #4 Average Score (Subtotal/3)		
Element #5: Treat		
Effective EBT		
<i>Subtotal</i>		
Element #5 Average Score (Subtotal/1)		
Element #6: Transition		
Continuous Contact & Support (Engagement)		
Continuous Contact and Support (Follow-up)		
<i>Subtotal</i>		
Element #6 Average Score (Subtotal/2)		
Element #7 Improve		
Approach to Reviewing Deaths		
Approach to Measuring Suicide Deaths		
Quality Improvement Activities		
<i>Subtotal</i>		
Element #7 Average Score (Subtotal/4)		

Overall average score (sum of average scores for each element/7)		
Date Completed		

Element #1: Lead

Create a leadership-driven, safety-oriented culture committed to dramatically reducing suicide among people under care. Include suicide attempt and loss survivors in leadership and planning roles.

Leadership-driven, safety-oriented culture:	Rating	1	2	3	4	5
What type of commitment has leadership made to reduce suicide and provide safer suicide care?		The organization has no processes specific to suicide prevention and care, other than what to do when someone mentions suicide during intake or a session.	The organization has 1–2 formal processes specific to suicide care.	The organization has written processes specific to suicide care. They have been developed for at least 3 different components of Zero Suicide.	The organization has processes and protocols specific to suicide care. They address at least 5 components of Zero Suicide. Staff receive training on processes as part of their orientations or when new ones developed. Processes are reviewed and modified at least annually.	Processes address all components of Zero Suicide listed above. Staff receives annual training on processes and when new ones are introduced. Processes are reviewed and modified annually and as needed.
		<u>Comment or justification for score:</u>				
Written Policies	Rating	1	2	3	4	5
Does organization have written protocols for specific components of suicide care, including (1) screening, (2) assessment, (3) lethal means restriction, (4) safety planning, and (5) suicide care management plans?		The organization has not discussed any protocols related to suicide care in the past year. No written policies exist.	The organization has discussed protocols related to suicide care in the past year, and is in the process of developing written policies.	The organization has adopted written policies for at least 2 of the 5 named components of suicide care.	The organization has adopted written policies for at least 4 of the 5 named components of suicide care, but they have <u>not</u> been discussed with staff.	The organization has written policies for all five of the named policies, <u>and</u> leadership has reviewed them verbally with staff.
		<u>Comment or justification for score:</u>				

Documentation	Rating	1	2	3	4	5
Are specific components of suicide care embedded in organization's electronic health record or easily identifiable in your written documentation (if no EHR is available), including (1) screening, (2) assessment, (3) lethal means restriction, (4) safety planning, and (5) suicide care management plans?		No suicide care components are embedded in organization's electronic health record or written documentation.	The organization has discussed embedding suicide care components into the EHR, but they are not currently active data fields.	At least 2 of the 5 named components of suicide care are embedded into the EHR or written documentation.	At least 4 of the 5 named components of suicide care are embedded into the EHR or written documentation, but they are required or routinely documented by staff.	All of the 5 named components of suicide care are embedded into the EHR or written documentation, and they are required or routinely documented by staff.
		<u>Comment or justification for score:</u>				
Training	Rating	1	2	3	4	5
Is training provided on specific components of suicide care, including (1) screening, (2) assessment, (3) lethal means restriction, (4) safety planning, and (5) suicide care management plans?		No training has been developed or provided on specific components of suicide care.	The organization is developing or choosing an existing training curricula on suicide care, and is in the process of scheduling training dates.	The organization has conducted at least one training on at least 2 of the 5 named components of suicide care.	The organization has conducted at least one training on at least 4 of the 5 named components of suicide care, <u>and</u> at least 50% of administrative and direct service staff have been trained.	The organization has conducted multiple trainings on all five of the named suicide care components, <u>and</u> 100% of current administrative and direct service staff have been trained.
		<u>Comment or justification for score:</u>				
Staffing	Rating	1	2	3	4	5
What type of formal commitment has leadership made through staffing to reduce suicide and provide safer suicide care?		The organization does not have dedicated staff to build and manage suicide care processes.	The organization has one leadership or supervisory individual who is responsible for developing suicide-related processes and care expectations. Responsibilities are diffuse. Individual does not have the authority to change policies.	The organization has assembled an implementation team that meets on an as-needed basis to discuss suicide care. The team has authority to identify and recommend changes to suicide care practices.	The organization has a formal Zero Suicide implementation team that meets regularly. The team is responsible for developing guidelines and sharing with staff.	The Zero Suicide implementation team meets regularly and is multidisciplinary. Staff members serve on the team for terms of one to two years. The team modifies processes based on data review and staff input.
		<u>Comment or justification for score:</u>				

Roles for survivors	Rating	1	2	3	4	5
<p>What is the role of suicide attempt and loss survivors in the organization's design, implementation, and improvement of suicide care policies and activities?</p>		<p>Suicide attempt or loss survivors are not explicitly involved in the development of suicide prevention activities within the organization.</p>	<p>Suicide attempt or loss survivors have ad hoc or informal roles within the organization, such as serving as volunteers or peer supports.</p>	<p>Suicide attempt or loss survivors are specifically and formally included in the organization's general approach to suicide care, but involvement is limited to one specific activity, such as leading a support group or staffing a crisis hotline. Survivors informally provide input into the organization's suicide care policies.</p>	<p>Suicide attempt and loss survivors participate as active members of decision-making teams, such as the Zero Suicide implementation team.</p>	<p>Suicide attempt and loss survivors participate in a variety of suicide prevention activities within the organization, such as sitting on decision-making teams or boards, participating in policy decisions, assisting with employee hiring and training, and participating in evaluation and quality improvement.</p>
		<p><u>Comment or justification for score:</u></p>				

Element #2: Train

Develop a competent, confident and caring workforce.

Workforce Confidence	Rating	1	2	3	4	5
How does the organization formally assess staff on their perception of their confidence, skills, and perceived support to care for individuals at risk for suicide?		There is no formal assessment of staff on their perception of confidence and skills in providing suicide care.	Clinicians who provide direct patient care are routinely asked to provide suggestions for training.	Clinical staff complete a formal assessment of skills, needs, and supports regarding suicide care. Training is tied to the results of this assessment.	A formal assessment of the perception of confidence and skills in providing suicide care is completed by all staff (clinical and non-clinical). Comprehensive organizational training plans are tied to the results.	A formal assessment of the perception of confidence and skills in providing suicide care is completed by all staff and reassessed at least every three years. Organizational training and policies are developed and enhanced in response to perceived staff weaknesses.
		<u>Comment or justification for score:</u>				
Non-clinical staff	Rating	1	2	3	4	5
What basic training on identifying people at risk for suicide or providing suicide care has been provided to NON-CLINICAL staff?		There is no organization-supported training on suicide care and no requirement for staff to complete training on suicide risk identification.	Training is available on suicide risk identification and care through the organization but not required of staff.	Training is required of select staff (e.g., crisis staff) and is available throughout the organization.	Training on suicide risk identification and care is required of all organization staff. The training used is considered a best practice and was not internally developed.	Training on suicide risk identification and care is required of all organization staff. The training used is considered a best practice. Staff repeat training at regular intervals.
		<u>Comment or justification for score:</u>				
Clinical staff	Rating	1	2	3	4	5
What advanced training on identifying people at risk for suicide, suicide assessment, risk formulation, and ongoing management has been provided to CLINICAL staff?		There is no organization-supported training on identification of people at risk for suicide, suicide assessment, risk formulation, and ongoing management, and no requirement for clinical staff to complete training on suicide.	Training is available on identification of people at risk for suicide, suicide assessment, risk formulation, and ongoing management through the organization, but it is not required of clinical staff.	Training is required of select staff (e.g., psychiatrists) and is available throughout the organization.	Training on identification of people at risk for suicide, suicide assessment, risk formulation, and ongoing management is required of all clinical staff. The training used is considered a best practice and was not internally developed.	Training on identification of people at risk for suicide, suicide assessment, risk formulation, and ongoing management is required of all clinical staff. The training used is considered a best practice. Staff repeat training at regular intervals.
		<u>Comment or justification for score:</u>				

Element #3: Identify

Systematically identify and assess suicide risk among people receiving care.

Screening Policies	Rating	1	2	3	4	5
What are the organization's policies for screening for suicide risk?		There is no systematic screening for suicide risk.	Individuals in designated higher-risk programs or categories (e.g., crisis calls) are screened.	Suicide risk is screened at intake for all individuals receiving behavioral health care.	Suicide risk is screened at intake for all individuals receiving either health or behavioral health care and is reassessed at every visit for those at risk.	Suicide risk is screened at intake for all individuals receiving health or behavioral health care and is reassessed at every visit for those at risk. Suicide risk is also screened when a patient has a change in status: transition in care level, change in setting, change to new provider, or potential new risk factors (e.g., change in life circumstances, such as divorce, unemployment, or a diagnosed illness).
<p><u>Comment or justification for score:</u> Number of clients who received a suicide screening during the reporting period/ Number of clients enrolled during the reporting period (___ / ___ = ___ %)</p>						

Screening Protocols	Rating	1	2	3	4	5
How does the organization screen for suicide risk in the people it serves?		The organization relies on the clinical judgment of its staff regarding suicide risk.	The organization developed its own suicide screening tool but not all staff are required to use it.	The organization developed its own suicide screening tool that all staff are required to use.	The organization uses a validated screening tool that all staff are required to use.	The organization uses a validated screening tool and staff receive training on its use and are required to use it.
<p><u>Comment or justification for score:</u> Screening tool used:</p>						

Assessment Protocols	Rating	1	2	3	4	5
How does the organization assess suicide risk among those who screened positive?		The policy is to send clients who have screened positive for suicide to the emergency department for clearance AND/OR there is no routine procedure for risk assessments that follow the use of a suicide screen.	Risk assessment is required after screening, but the process or tool used is up to the judgment of individual clinicians AND/OR only psychiatrists can do risk assessments.	Providers conducting risk assessments use a standardized risk assessment tool, which may have been developed in-house. All patients who screen positive for suicide have a risk assessment. Suicide risk assessments are documented in the medical records.	All individuals with risk identified, either at intake screening or at any other point during care, are assessed by clinicians who use validated instruments or established protocols and who have received training. Assessment includes both risk and protective factors.	A suicide risk assessment is completed using a validated instrument and/or established protocol that includes assessment of both risk and protective factors and risk formulation. Staff receive training on risk assessment tool and approach. Risk is reassessed and integrated into treatment sessions for every visit for individuals with risk.
<p><u>Comment or justification for score:</u> Number of clients who screened positive for suicide risk and had a comprehensive risk assessment (same day as screening) during the reporting period/ Number of clients who screened positive for suicide risk during the reporting period (___ / ___ = ___%)</p>						

Element #4: Engage

Ensure every person has a suicide care management plan, or pathway to care, that is both timely and adequate to meet patient needs.

Pathway to Care	Rating	1	2	3	4	5
Which best describes the organization's approach to caring for and tracking people at risk for suicide?		Providers use best judgment in the care of individuals with suicidal thoughts or behaviors and seek consultation if needed. There is no formal guidance related to care for individuals at risk for suicide.	When suicide risk is detected, the care plan is limited to screening and referral to a senior clinician.	All providers are expected to provide care to those at risk for suicide. The organization has guidance for care management for individuals at different risk levels, including frequency of contact, care planning, and safety planning.	Electronic or paper health records are enhanced to embed all suicide care management components listed above. Providers have clear protocols or policies for care management for individuals with suicidal thoughts or behaviors, and information sharing and collaboration among all relevant providers are documented. Staff receive guidance on and clearly understand the organization's suicide care management approach.	Individuals at risk for suicide are placed on a suicide care management plan. The organization has a consistent approach to suicide care management, which is embedded in the electronic health records and reflects all of the suicide care management components listed above. Protocols for putting someone on and taking someone off a care management plan are clear. Staff hold regular case conferences about patients who remain on suicide care management plans beyond a certain time frame, which is established by the implementation team.
		<u>Comment or justification for score:</u>				
Collaborative Safety Planning	Rating	1	2	3	4	5
What is the organization's approach to collaborative safety planning when an individual is at risk for suicide?		Safety planning is neither systematically used by nor expected of staff.	Safety plans are expected for all individuals with elevated risk, but there is no formal guidance or policy around content. There is no standardized safety plan or documentation template. Plan quality varies across providers.	Safety plans are developed for all individuals at elevated risk. Safety plans rely on formal supports or contact (e.g., call provider, call helpline). Safety plans do not incorporate individualization, such as an individual's strengths and natural supports. Plan quality varies across providers.	Safety plans are developed for all individuals at elevated risk and must include risks and triggers and concrete coping strategies. The safety plan is shared with the individual's partner or family members (with consent). All staff use the same safety plan template and receive training in how to create a collaborative safety plan.	A safety plan is developed on the same day as the patient is assessed positive for suicide risk. The safety plan is shared with the individual's partner or family members (with consent). The safety plan identifies risks and triggers and provides concrete coping strategies, prioritized from most natural to most formal or restrictive. Other clinicians involved in care or transitions are aware of the safety plan. Safety plans are reviewed and modified as needed at every visit with a person at risk.
		<u>Comment or justification for score:</u> (1) Safety planning tool or approach used by organization: <input type="checkbox"/> Stanley/Brown template <input type="checkbox"/> Other: _____. (2) How frequently is safety plan reviewed with individual? _____ (3) Number of clients with a safety plan developed on same day as screening during the reporting period / Number of clients who screened and assessed positive for suicide risk during the reporting period (___ / ___ = ___%)				

Collaborative Restriction of Access to Lethal Means	Rating	1	2	3	4	5
What is the organization's approach to lethal means reduction?		Means restriction discussions and who to ask about lethal means are up to individual clinician's clinical judgment. Means restriction counseling is rarely documented.	Means restriction is expected to be included on safety plans for all patients identified as at risk for suicide. Steps to restrict means are up to the individual clinician's judgment. The organization does not provide any training on counseling on access to lethal means	Means restriction is expected to be included on all safety plans. The organization provides training on counseling on access to lethal means. Steps to restrict means are up to the individual clinician's judgment. Family or significant others may or may not be involved in reducing access to lethal means.	Means restriction is expected to be included on all safety plans, and families are included in means restriction planning. The organization provides training on counseling on access to lethal means. The organization sets policies regarding the minimum actions for restriction of access to means.	Means restriction is expected to be included on all safety plans. Contacting family to confirm removal of lethal means is the required, standard practice. The organization provides training on counseling on access to lethal means. Policies support these practices. Means restriction recommendations and plans are reviewed regularly while the individual is at an elevated risk.
		<p><u>Comment or justification for score:</u></p> <p>Number of clients screened & assessed positive for suicide risk <u>and</u> counseled about lethal means on same day as screening) during reporting period / Number of clients who screened and assessed positive for suicide risk during reporting period (___ / ___ = ___%)</p>				

Element #5: Treat

Use effective, evidence-based treatments that directly target suicidality.

Effective, EBT	Rating	1	2	3	4	5
What is the organization's approach to treatment of suicidal thoughts and behaviors?		Clinicians rely on experience and best judgment in risk management and treatment for all mental health disorders. The organization does not use a formal model of treatment for those at risk for suicide.	The organization may use evidence-based treatments for some psychological disorders, but it does not use evidence-based treatments that specifically target suicide.	Some clinical staff have received specific training in treating suicidal thoughts and behaviors and may use this in their practices.	Individuals with suicide risk receive empirically-supported treatment specifically for suicide (CAMS, CBT-SP or DBT) in addition to evidence-based treatments for other mental health issues. The organization regularly provides all staff with access to competency-based training in empirically supported treatments targeting suicidal thoughts.	The organization has invested in evidence-based treatments for suicide care (CAMS, CBT-SP or DBT), with designated staff receiving training in these models. The organization has a model for sustaining staff training. The organization offers additional treatment modalities for those chronically or continuously screening at high risk for suicide, such as DBT groups or attempt survivor groups.
		<p><u>Comment or justification for score:</u></p> <p>Clinicians receive formal training in a specific suicide treatment model: <input type="checkbox"/> CAMS (Collaborative Assessment and Management of Suicidality) <input type="checkbox"/> C BT-SP (Cognitive Behavioral Therapy for Suicide Prevention) <input type="checkbox"/> DBT (Dialectical Behavior Therapy) <input type="checkbox"/> None of the above</p>				

Element #6: Transition

Provide continuous contact and support, especially after acute care.

Continuous contact & support (Engagement)	Rating	1	2	3	4	5
What is the organization's approach to engaging hard-to-reach individuals or those who are at risk and don't show for appointments?		There are no guidelines specific to reaching those at elevated suicide risk who don't show for scheduled appointments.	The organization requires documentation by the clinician of those individuals who have elevated suicide risk and don't show for an appointment, but the parameters and methods are up to individual clinician's judgment.	Follow-up for individuals with suicide risk who don't show for appointments includes active outreach, such as phone calls to the individual or his or her family members, until contact is made and the individual's safety is ascertained.	Follow-up for individuals with suicide risk who don't show for appointments includes active outreach, such as phone calls to the individual or his or her family members, until contact is made and the individual's safety is ascertained. Organizational protocols are in place that address follow-up after no-shows. Training for staff supports improving engagement efforts.	The organization may have an established memorandum of understanding with an outside agency to conduct follow-up calls. Follow-up and supportive contact for individuals on suicide care management plans are systematically tracked in electronic health records. Follow-up for high-risk individuals includes documented contact with the person within eight hours of the missed appointment. The organization has approaches, such as peer supports, peer-run crisis respite, home visits, or drop-in appointments, to address the needs of hard-to-reach patients.
<u>Comment or justification for score:</u>						
Continuous contact & support (Follow-up)	Rating	1	2	3	4	5
What is the organization's approach to following up on patients who have recently been discharged from acute care settings (e.g., emergency departments, inpatient psychiatric hospitals)?		There are no specific guidelines for contact of those at elevated suicide risk following discharge from acute care settings.	The organization requires follow-up for individuals with suicide risk, but the parameters and methods are up to the individual clinician's judgment.	Organizational guidelines are directed to the individual's level of risk and address one or more of the following: follow-up after crisis contact, transition from an emergency department, or transition from psychiatric hospitalization.	Organizational guidelines are directed to the individual's level of risk and address follow-up after crisis contact, non-engagement in services, transition from an emergency department, or transition from psychiatric hospitalization. Follow-up for high-risk individuals includes distance outreach, such as letters, phone calls, or e-mails.	Organizational guidelines are in place that address follow-up after crisis contact, no-shows, transition from an emergency department, or transition from psychiatric hospitalization. Follow-up for high-risk individuals includes in-person or virtual home or community visits when necessary. Follow-up and supportive contact for individuals on suicide care management plans are tracked in the electronic health record. Policies state that follow-up contact after discharge from acute settings occurs within 24 hours.
<u>Comment or justification for score:</u>						

Element #7: Improve

Apply a data-driven quality improvement approach to inform system changes that will lead to improved patient outcomes and better care for those at risk.

Approach to reviewing deaths	Rating	1	2	3	4	5
What is the organization's approach to reviewing deaths for those enrolled in care?		At best, when a suicide or adverse event happens while the client is in treatment, a team meets to discuss the case.	Root cause analysis is conducted on all suicide deaths of people in care.	Data from all root cause analyses are routinely examined to look at trends and to make changes to policies.	Root cause analysis is conducted on all suicide deaths of people in care as well as for those up to 30 days past case closed. Policies and training are updated as a result.	Root cause analysis is conducted on all suicide deaths of people in care as well as for those up to 6 months past case closed, and on all suicide attempts requiring medical attention. Policies and training are updated as a result.
		<p><u>Comment or justification for score:</u></p> <p>Date of most recent root cause analysis of a suicide death: _____. Date of most recent suicide death of (1) someone in care: _____ (2) someone who had left care less than 6 months before suicide death _____</p>				
Approach to measuring suicide deaths	Rating	1	2	3	4	5
What is the organization's approach to measuring suicide deaths?		The organization has no policy or process to measure suicide deaths for those enrolled in their care.	The organization measures the number of deaths for those who are enrolled in care based primarily on family report.	The organization has specific internal approaches to measuring and reporting on all suicide deaths for enrolled clients as well as those up to 30 days past case closed. Deaths are confirmed through coroner or medical examiner reports.	The organization annually crosswalks enrolled patients (e.g., from a claims database) against state vital statistics data or other federal data to determine the number of deaths for those enrolled in care up to <u>30 days</u> past case closed.	The organization annually crosswalks enrolled patients (e.g., from a claims database) against state vital statistics data to determine the number of deaths for those enrolled in care. The organization tracks suicide deaths among clients for up to <u>6 months</u> past case closed.
		<p><u>Comment or justification for score:</u></p> <p>Date measurement for suicide deaths was established: _____</p> <p>Date of most recent annual crosswalk of enrolled patients against vital statistics data: _____</p>				

Quality improvement activities	Rating	1	2	3	4	5
<p>What is the organization's approach to quality improvement activities related to suicide prevention?</p>		<p>The organization has no specific policies related to suicide prevention and care, and it does not focus on suicide care other than care as usual. Care is left to the judgment of the clinical provider.</p>	<p>Suicide care is discussed as part of employee training and by those in supervision in clinical settings.</p>	<p>Early discussions about using technology and/or enhanced record keeping to track and chart suicide care are underway. Suicide care management is partially embedded in an EHR or paper record.</p>	<p>Suicide care is partially embedded in an electronic health record (EHR) or paper record. Data from suicide care management plans (using EHRs or chart reviews) are examined for fidelity to organizational policies, and discussed by a team responsible for this.</p>	<p>Suicide care is entirely embedded in EHR. Data from EHR or chart reviews are routinely examined (at least every two months) by a designated team to determine that staff are adhering to suicide care policies and to assess for reductions in suicide. EHR clinical workflows or paper records are updated regularly as the team reviews data and makes changes.</p>
<p><u>Comment or justification for score:</u></p> <p>Most recent date that data from EHR or chart reviews were examined for adherence to suicide care policies _____</p>						