Portland State University

PDXScholar

Environmental Science and Management
Faculty Publications and Presentations

Environmental Science and Management

9-2015

Coastal Sediment Elevation Change Following Anthropogenic Mangrove Clearing

Heather Lyn Hayden
Portland State University

Elise F. Graneke
Portland State University, graneke@pdx.edu

Follow this and additional works at: https://pdxscholar.library.pdx.edu/esm_fac

Part of the [Biochemistry Commons](#), [Marine Biology Commons](#), [Oceanography Commons](#), and the [Terrestrial and Aquatic Ecology Commons](#)

Let us know how access to this document benefits you.

Citation Details

Hayden, H.L., Graneke, E.F., COASTAL sediment elevation change following anthropogenic mangrove clearing, *Estuarine, Coastal and Shelf Science* (2015), doi: 10.1016/j.ecss.2015.09.004.

This Post-Print is brought to you for free and open access. It has been accepted for inclusion in Environmental Science and Management Faculty Publications and Presentations by an authorized administrator of PDXScholar. Please contact us if we can make this document more accessible: pdxscholar@pdx.edu.

1 COASTAL SEDIMENT ELEVATION CHANGE FOLLOWING ANTHROPOGENIC
2 MANGROVE CLEARING
3

4
5 Heather L. Hayden¹, Elise F. Granek¹²

6 ¹Environmental Science and Management, Portland State University, Portland,
7 Oregon 97207, USA

8 ²Corresponding author: graneke@pdx.edu; 1-503-725-4241; 1719 SW 10th Ave,
9 Environmental Science and Management, Portland State University, Portland,
10 Oregon 97207, USA

11
12
13
14 Key Words: accretion, Belize, deforestation, ecosystem service, erosion,
15 mangrove
16
17

1 ABSTRACT

2
3 Coastal mangrove forests along tropical shorelines serve as an important
4 interface between land and sea. They provide a physical buffer protecting the
5 coastline from erosion and act as sediment “traps” catching terrestrial sediment,
6 thus preventing smothering of subtidal coral reefs. Coastal development that
7 removes mangrove habitat may impact adjacent nearshore coral reefs through
8 sedimentation and nutrient loading. We examined differences in sediment
9 elevation change between patches of open-coast intact and anthropogenically
10 cleared red mangroves (*Rhizophora mangle*) on the east side of Turneffe Atoll,
11 Belize, to quantify changes following mangrove clearing. Samples were collected
12 over a 24 month period at five study sites, each containing paired intact
13 (+mangrove) and cleared (-mangrove) plots. Five sediment elevation pins were
14 deployed in each plot: behind areas cleared of mangroves (-mangrove) and
15 behind adjacent intact mangroves (+mangrove). Sediment elevation increased at
16 intact mangrove sites ($M = +3.83$ mm, $SE = 0.95$) whereas cleared mangrove
17 areas suffered elevation loss ($M = -7.30$ mm, $SE = 3.38$). Mangroves inshore of
18 partial or continuous gaps in the adjacent fringing reefs had higher rates of
19 elevation loss ($M = -15.05$ mm) than mangroves inshore of continuous fringing
20 reefs ($M = -1.90$ mm). Our findings provide information on potential effects of
21 mangrove clearing and the role of offshore habitat characteristics on coastal
22 sediment trapping and maintenance of sediment elevation by mangroves. With
23 implications for coastline capacity to adjust to sea level rise, these findings are
24 relevant to management of coastal fringing mangrove forests across the
25 Caribbean.

I. INTRODUCTION

Mangrove forests serve integral ecological functions in tropical coastal zones (Mumby et al. 2004), yet damage to and loss of mangrove forests has been extensive worldwide (Crooks and Turner 1999). Mangrove forests have been rapidly cleared and converted for real estate speculation, construction of tourist resorts, agricultural land reclamation and other coastal development; shrimp aquaculture; and to a lesser extent, collection of lumber (Kaly et al. 1997, Stevenson 1997, Valiela et al. 2001). Mangrove forest loss due to anthropogenic clearing has occurred at an alarming rate (-1% annually in the 1980's [FAO 2003]; -0.66% annually between 2000-2005 [FAO 2007]) resulting in a significant decrease in global mangrove forest cover (estimated losses of up to 35% during the last two decades of the 20th century [Valiela et al. 2001]). Similarly, mainland Caribbean mangroves have suffered a 1.7% annual decline in aerial cover (Ellison and Farnsworth 1996). Some researchers project a complete loss of this ecosystem type from certain regions within the next 100 years (Duke et al. 2007).

Mangroves play a key role in ecological functioning of adjacent nearshore habitats and provide a number of ecosystem services to surrounding coastal communities. Coastal mangrove forests serve as a key habitat for ecologically and commercially important species (e.g., Mumby et al. 2004, Igulu et al. 2014) and are important for carbon storage (Blue Carbon, e.g., Donato et al. 2011). This coastal ecosystem also serves as an key physical buffer, protecting coastal zones from erosion, extensive wave action, and flooding during tropical storms and hurricanes (Field 1998, Ellison 2000, Danielsen et al. 2005, Kar 2005,

Granek and Ruttenberg 2007). As oceanic storms increase in frequency and intensity with warming sea surface temperatures (Williams 2005) and as sea level rises, the importance of mangroves is expected to increase. Finally, coastal mangroves serve as sediment “traps” accreting terrestrial sediment and preventing it from smothering subtidal coral reefs (Bird 1971, Wolanski and Ridd 1986, Augustinus 1995, Blasco et al. 1996, Woodroffe 2002, Golbuu et al. 2003, Thampanya et al. 2006, Victor et al. 2006).

These ecological functions can be disrupted when natural disturbance events cause mangrove mortality (e.g., Cahoon et al. 2003). However, the sediment trapping response of these systems to anthropogenic clearing is largely absent from the literature. Comparative research assessing differences in sediment elevation between intact and anthropogenically cleared mangrove stands is an existing data gap. We quantify how coastal development that removes mangrove habitat impacts shoreline sediment levels since sediment loss has potential implications for nutrient loading and sedimentation on adjacent nearshore habitats as well as on shoreline loss in the face of climate change. We tested the hypothesis that in coastal fringing red mangrove (*Rhizophora mangle*) ecosystems on Turneffe Atoll, Belize, sediment elevation levels decline following mangrove clearing relative to levels in adjacent intact mangroves.

II. METHODS

Study Site

The study was conducted on Turneffe Atoll off the Caribbean coast of Belize. Turneffe Atoll is a carbonate atoll with little or no allochthonous sediment

input; sediment in the mangroves is autochthonous, derived from accumulation of biogenic material, primarily mangrove subsurface peat with lenses of calcareous sand (McKee et al. 2007). The coastline of Turneffe Atoll is characterized by open coast fringing *R. mangle* (red mangrove) trees, except in locations where mangroves were removed for development and/or agriculture. The submerged vegetation in front of the study sites is dominated by *Thalassia testudinum* (turtle grass). Five locations were sampled along the east coast of Turneffe Atoll with each study site consisting of paired intact (+mangrove) and cleared (-mangrove) areas. All study sites were along the open coast and located within a 15km stretch of coastline (Figure 1 and Table 1). Wave energy is comparable within each pair. The distance between the mangrove or cleared shoreline and the fringing reef and the continuity of the fringing reef are consistent within pairs but variable among sites, leading to potential between site differences in wave energy reaching the shoreline.

The following criteria were used for site selection: 1) at least 75m along-shore length of cleared *R. mangle* adjacent to stretches of at least 100m of intact *R. mangle* habitat; and 2) >2km from major human development to reduce potential sources of anthropogenic nutrients and sediment. The cleared mangrove areas ranged from 75m to 250m in length along the shore and ranged from recent (within 2 years of study onset) to historic (40-200 years prior) clearings (Table 1). At the recently cleared sites, above ground root structure was still present, though no above ground roots were evident at the historically cleared sites. Elevations were similar within pairs at recently cleared sites at the

beginning of the study; however, at Calabash there was some seagrass growing in the cleared sediment elevation pin plot indicating lower elevation than its adjacent paired intact site.

Figure 1. Location of study area in the Caribbean Sea (left) with sites marked on Turneffe Atoll map (center): from north to south, sites include Fisherman's Cut (FC), Airport (A), Oceanic Society (OS), Calabash (C), and Ropewalk (RW); Airport study site (right) with sediment elevation pins in intact and cleared mangrove areas inshore and fringing reef offshore.

Areas with intact mangroves were characterized by monospecific stands of *R. mangle* with submerged prop roots and infrequent aerial roots colonized with sponges, epibiotic algae, tunicates and anemones. Recently cleared mangrove areas retained some structure from the submerged decaying prop roots; the soil was "muddy" with roots throughout. Historically cleared areas had

1 little to no submerged decaying prop roots, with soil consisting of coarse sand
2 with seagrass moving into former mangrove habitat.

3 Cleared areas were absent of mature trees and had sparsely dispersed
4 settlement of seedlings (~1 seedling per m²) compared to adjacent intact
5 mangrove forest with mature trees and high seedling densities (~8 seedlings per
6 m²). The cleared areas lack forest structure. The mangroves settled in cleared
7 areas were “dwarfed” (~1 m tall) relative to individuals in intact areas (up to 4 m
8 tall).

9 Additional anthropogenic disturbances impacted two sites during the study
10 period (in 2009). In the intact mangrove area at Airport (Figure 1), a section of
11 mangroves was cleared and dredging was conducted for condominium
12 development. Two sediment elevation pins were lost during the clearing.
13 Adjacent to the cleared site at Ropewalk, dredging was conducted to facilitate
14 boat access, increasing suspended sediment in the cleared area.

15

16 *Sediment Elevation Pins*

17 Tubes of PVC piping with 2.5 cm external diameter x 1.5 m length were used as
18 sediment elevation pins to quantify changes in elevation (e.g., Stokes et al. 2009).
19 Elevation changes may be due to accretion or erosion. Drivers of erosion include
20 sediment leaving the system, compaction, and shrink/swell of aerially exposed
21 sediment. Drivers of accretion include sediment trapping and subsurface expansion
22 from root growth and groundwater influx (Krauss et al. 2014, Cahoon et al. 2003). At
23 each site, five sediment elevation pins were deployed behind areas cleared of

mangroves (-mangrove) and five pins were deployed at the same distance from the seaward edge of the mangroves behind intact forest (+mangrove). At all sites except Calabash, these pins were placed during a previous study by S. Waddington and E. Granek. The protocol employed for placing the sediment elevation pins was adopted from dune studies by Moreno-Cassola (1986) and Arens and Slings (2004). The research plots are located near fishing camps, developed resorts, homes, and a Belize National Coast Guard Station. The current method of measuring mangrove sediment accretion/erosion, Surface Elevation Table (SET) (Cahoon et al. 2003) equipment, was infeasible at our study site due to the risk of equipment removal by transient fishers. Therefore we employed a low-tech, low-cost technique to measure sediment elevation change relative to a ~1-m deep benchmark.

The 1.5m long PVC piping tubes were buried in the substrate with 60cm of tube remaining above ground. Differences in rates of sediment elevation change were quantified by measuring the length of tube above the sediment at 0-, 6-, 12- and 24-months post deployment. The PVC tubes were measured to the nearest 0.1 cm from the top of the tube to the sediment on the ocean and terrestrial sides. Each pin was gently pulled upward and pushed downward to confirm that pins were secure and not floating in the hole. All pins were secure at each sampling period.

Data analysis

The five sediment elevation pins per plot were averaged and examined at 12- and 24- months post deployment. Changes in sediment elevation between intact and cleared mangrove plots and among sites were examined using a two-way ANOVA. A

multi-factor ANOVA was used to examine whether sedimentation rates differed by continuity of adjacent reef structure, site, and mangrove condition (intact vs. cleared).

III. RESULTS

Sediment elevation increased ($M = 3.83\text{mm}$, $SE = 0.95$) at the intact mangrove plots but decreased ($M = -7.30\text{ mm}$, $SE = 3.38$) at the cleared mangrove plots over the course of the study (Table 1). Between June 2008 and June 2010, sediment elevation change differed between intact and cleared areas ($F = 79.52$, $p = 0.001$), across sites ($F = 6.41$, $p < 0.001$), and the differences in elevation change between intact and cleared areas varied by site ($F = 9.57$; $p < 0.001$) (Figure 2).

In cleared areas, elevation loss rates were greater at sites with gaps in the adjacent fringing reef (Airport and Oceanic; $M = -15.05\text{mm}$) than at sites with continuous fringing reef ($M = -1.90\text{mm}$; $F = 18.59$, $p < 0.001$). In intact mangrove forest, elevation gain did not differ between sites with reef gaps ($M = 4.11\text{mm}$) and sites with continuous reef ($M = 3.65\text{mm}$). Sediment elevation gain was greater in intact than cleared areas ($F = 81$, $p < 0.001$) and the effects of reef continuity on sediment elevation gain or loss varied by mangrove condition ($F = 32.33$, $p < 0.001$).

Table 1. Fringing reef structure, anthropogenic disturbances during study period, and mean (and standard error) sediment elevation gain (+) and loss (-) rates at cleared and intact plots for each site (Year 1 = June 2008 to June 2009, Year 2 = June 2009 to June 2010).

Site	Airport	Calabash	Fisherman's	Ropewalk	Oceanic
Fringe Reef Protection	<i>Few gaps</i>	Complete	Complete	Complete	<i>Gaps</i>
Dominant sediment type	Sandy	Sandy	Detrital mud	Sandy	Sandy
Years since clearing	4	~40	2	8	~200
Anthropogenic disturbances	Dredging & clearing	None	None	Dredging 2009	None

		2009				
Mean soil accretion (mm) in intact plots	Year 1	+1.62	+5.33	+2.04	+0.73	+3.58
	Year 2	+2.08 (0.82)	+4.89 (2.23)	+4.92 (1.12)	+1.13 (0.72)	+6.14 (0.31)
Mean soil elevation loss (mm) cleared areas	Year 1	-10.7	-0.35	-2.3	-0.25	-6.32
	Year 2	-16.73 (7.39)	-2.06 (0.19)	-3.56 (0.48)	-0.09 (0.97)	-20.18 (2.3)

Figure 2. Sediment elevation change rates (in mm) between June 2008 and June 2010 at five study sites on Turneffe Atoll.

IV. DISCUSSION

Coastal mangrove forests function as important physical buffers protecting coastlines from erosion and nearshore marine ecosystems from sedimentation (Gillis et al. 2014). When mangroves are naturally or anthropogenically thinned or removed, this function is reduced or eliminated (e.g., Cahoon et al. 2003; Stokes et al. 2009). Similarly, intact mangroves on Turneffe Atoll demonstrate sediment elevation gains

1 whereas adjacent cleared areas suffered sediment elevation losses over the same time
2 period. Cleared sites with discontinuous fringing reef offshore suffered higher sediment
3 elevation loss rates than sites with continuous fringing reefs. These data add to the
4 growing body of evidence quantifying the ecological function provided by coastal
5 mangrove forest habitat to maintain soil elevation (e.g., Stokes et al. 2009, Cahoon et
6 al. 2003, Krauss et al. 2003) and facilitate coastline adaptations to sea level rise (e.g.,
7 Krauss et al. 2014).

8 Natural variability in fringing reef structure can affect sediment patterns (Table 1;
9 Gillis et al. 2014). Study sites with (50-100 m alongshore gaps in the fringing reef
10 (Airport and Oceanic), had higher sediment elevation loss rates (-16 to -20 mm) in the
11 cleared areas. Gaps in the fringing reef reduce protection from incoming wave action
12 (Kunkel et al. 2006, Sheppard 2005), which can result in higher sediment elevation loss
13 rates when mangroves are removed. On the other hand, Calabash and Fisherman's
14 have extensive and continuous reef protection and are located further from the reef
15 (Fisherman's is ~500 m and Oceanic is ~100 m inshore); these sites experienced lower
16 sediment elevation loss rates in the cleared areas (2-3.5 mm). These sites demonstrate
17 landscape-scale facilitation between reefs and mangroves (Gillis et al. 2014).

18 Anthropogenic disturbance events such as dredging can reduce sediment
19 elevation loss at cleared mangrove sites due to sediment resuspension. At Ropewalk,
20 dredging increased sediment suspension for 4 months during the summer of 2009; we
21 hypothesize that this increased sediment suspension led to increased sediment
22 deposition at the cleared area, counteracting sediment elevation loss due to absent
23 mangrove root structure (Figure 2).

Our sites on Turneffe Atoll have variable sediment dynamics due to variability in the fringing reef and the resultant wave energy in the mangroves as well as additional anthropogenic disturbances at certain sites. Though our method cannot identify the specific causes of the sediment elevation loss we report, the losses likely results from a combination of surface and subsurface subsidence from peat collapse due to decomposition of organic material (including roots) and erosion of the substrate due to the absence of root structure and cessation of mangrove organic matter deposition from loss of mangrove trees (Krauss et al. 2014, Cahoon et al. 2003). Reported rates of accretion in mangroves vary by geographic location, ranging from -38mm year^{-1} in Tauranga Harbour, New Zealand (Stokes et al. 2009) to $+10.0$ to $+11.0\text{ mm year}^{-1}$ in Cairns, Queensland, Australia (Bird and Barson 1977; Spencely 1982). Our reported rates of sediment elevation loss and gain fall within this range. However, to date, little research in mangrove ecosystems has directly compared sediment elevation changes between intact and cleared mangrove areas (but see Stokes et al. 2009). McKee et al. (2007) and McKee (2011) suggest that mangrove removal could lead to land and habitat stability losses due to a cessation of soil accretion concurrent with ongoing decomposition, compaction and erosion processes. Field data quantifying the effects of Caribbean mangrove clearing on sediment retention are absent from the current literature. Our data from sediment elevation pins provide the first quantitative evidence of sediment elevation loss in cleared areas at sites that gain sediment elevation in adjacent land where mangroves are present.

The intact mangroves on Turneffe Atoll likely protect sediment from elevation loss or erosion due to surface and subsurface physical and biological processes

including sedimentation, ground water swelling, plant litter, woody debris and root accumulation and algal mat development. These processes result in vertical sediment accretion in Caribbean mangrove ecosystems (Krauss et al. 2014; McKee 2011; Rogers et al. 2005). McKee (2011) reported low accretion rates in Belizean fringing mangroves but subsurface root growth accounting for 8.8 mm year^{-1} vertical change, with an overall elevation change of 4.1 mm year^{-1} after accounting for compaction, decomposition and hydrodynamics. Given that our accumulation rates in the intact mangrove areas were comparable, we hypothesize similar processes at work.

Our research finds that removal of mangrove vegetation leads to sediment elevation loss with potential for increased submergence and land loss. This loss is exacerbated where fringing reef is discontinuous and can vary as a result of additional disturbance events. These findings highlight the need to consider the role of mangroves in managing coastline protection with ongoing sea-level rise (Alongi 2015) and the importance of carefully considering subtidal features adjacent to sites slated for mangrove removal or transformation. Such information may prove useful in identifying areas that are most vulnerable to sediment elevation loss following mangrove removal and therefore areas of coastline most vulnerable to submergence with sea level rise. Better understanding and consideration of these processes can serve as a potential tool for determining where mangrove clearing will have the greatest impact on coastal protection generally and sea-level driven land loss specifically.

V. ACKNOWLEDGEMENTS

We thank the students from the Portland State University Belize Capstone Class from 2008, Oceanic Society for their in-kind and logistical support, Marion Usher and

Mr. Billy, our great boat tender.

VI. LITERATURE CITED

- Alongi, DM. (2015) The impact of climate change on mangrove forests. *Curr Clim Change Rep* 1:30-39
- Arens SM, Slings Q, De Vries CN (2004) Mobility of a remobilised parabolic dune in Kennemerland, The Netherlands. *Geomorphology* 59(1):175-188
- Augustinus PGEF (1995) Chapter 12 Geomorphology and Sedimentology of Mangroves. In: Perillo GME (ed) *Developments in Sedimentology*, Book Volume 53. Elsevier
- Bird ECF (1986) Mangroves and intertidal morphology in Westernport Bay, Victoria, Australia. *Marine Geology* 69:251-271
- Bird ECF, Barson MM (1977) Measurement of physiographic changes on mangrove-fringed estuaries and coastlines. *Marine Research in Indonesia*, 18:73-80
- Blasco F, Saenger P, Janodet E (1996) Mangroves as indicators of coastal change. *CATENA* 27:167-178
- Cahoon DR, Hensel P, Rybczyk J, McKee KL, Proffitt CE, Perez BC (2003) Mass tree mortality leads to mangrove peat collapse at Bay Islands, Honduras after Hurricane Mitch. *Journal of Ecology* 91:1093–1105
- Crooks S, Turner RK (1999) Integrated Coastal Management: Sustaining Estuarine Natural Resources. In: Nedwell D, Raffaelli D (eds) *Advances in Ecological Research*, Book Volume 29. Academic Press
- Danielsen F, Sørensen MK, Olwig MF, Selvam V, Parish F, Burgess ND, Hiraishi T, Karunakaran VM, Rasmussen MS, Hansen LB, Quarto A, Suryadiputra, N. (2005) The Asian tsunami: a protective role for coastal vegetation. *Science*, 310:643
- Donato D.C, Kauffman J.B, Murdiyarso D, Kurnianto S, Stidham M, Kanninen M. (2011) Mangroves among the most carbon-rich forests in the tropics. *Nature Geoscience* 4(5) 293-297.
- Duke NC, Meynecke J-O, Dittmann S, Ellison AM, Anger K, Berger U, Cannicci S, Diele K, Ewel KC, Field CD, Koedam N, Lee SY, Marchand C, Nordhaus I, Dahdouh-Guebas F (2007) A world without mangroves? *Science* 317:41-42
- Ellison AM (2000) Mangrove Restoration: Do We Know Enough? *Restoration Ecology* 8:219-229
- Ellison AM, Farnsworth EJ. (1996) Anthropogenic disturbance of Caribbean mangrove ecosystems: Past impacts, present trends and future predictions. *Biotropica* 28: 549– 565.
- FAO (Food and Agriculture Organization of the United Nations) (2003) Status and Trends in Mangrove Area Extent Worldwide. Forest Resources Division, Paris.
- FAO (Food and Agriculture Organization of the United Nations) (2007) The world's mangroves 1980–2005. FAO Forestry Paper 153. FAO, Rome.
- Field CD (1998) Rehabilitation of mangrove ecosystems: an overview. *Mar. Pollut. Bull.* 37:383–392

- Gillis LG, Bouma TJ, Jones CG, van Katwijk MM, Nagelkerke I, et al. (2014) Potential for landscape-scale positive interactions among tropical marine ecosystems. *Mar. Ecol. Prog. Ser.* 503:289–303
- Golbuu Y, Victor S, Wolanski E, Richmond RH (2003) Trapping of fine sediment in a semi-enclosed bay, Palau, Micronesia. *Estuarine, Coastal and Shelf Science* 57:941-949
- Granek EF, Ruttenberg BI (2007) Protective capacity of mangroves during tropical storms: a case study from 'Wilma' and 'Gamma'. *Mar Ecol Prog Ser* 343:101–105
- Greipsson S, Davy AJ (1996) Sand accretion and salinity as constraints on the establishment of *Leymus arenarius* for land reclamation in Iceland. *Annals of Botany* 78: 611-618
- Hugenholtz CH, Wolfe SA, Walker IJ, Moorman BJ (2009) Spatial and temporal patterns of aeolian sediment transport on an inland parabolic dune, Bigstick Sand Hills, Saskatchewan, Canada. *Geomorphology* 105(1):158-170
- Igulu MM, Nagelkerken I, Dorenbosch M, Grol MG, Harborne AR, Kimirei IA, Mgaya YD (2014) Mangrove habitat use by juvenile reef fish: meta-analysis reveals that tidal regime matters more than biogeographic region. *PloS one*, 9(12), e114715.
- Kaly UL, Eugelink G, Robertson AI (1997) Soil conditions in damaged North Queensland mangroves. *Estuaries* 20:291 - 300
- Kar R, Kar RK (2005) Mangroves can check the wrath of tsunami. *Current Sci* (Bangalore) 88:675
- Krauss KW, Allen JA, Cahoon DR. (2003) Differential rates of vertical accretion and elevation change among aerial root types in Micronesian mangrove forests. *Estuarine, Coastal, and Shelf Science* 56: 251-259
- Krauss KW, McKee KL, Lovelock CE, Cahoon DR, Saintilan N, Reef R, Chen L (2014) How mangrove forests adjust to rising sea level. *New phytologist*, 202(1), 19-34
- Kukel CM, Hallberg RW, Oppenheimer M (2006) Coral reefs reduce tsunami impact in model simulations. *Geophysical Research Letters* 33:23
- Levin N, Kidron GJ, Ben-Dor E (2006) The spatial and temporal variability of sand erosion across a stabilizing coastal dune field. *Sedimentology* 53(4): 697-715
- McKee KL (2011) Biophysical controls on accretion and elevation change in Caribbean mangrove ecosystems. *Estuarine, Coastal and Shelf Science* 1(4):475-483
- McKee KL, Cahoon DR, Feller IC (2007) Caribbean mangroves adjust to rising sea level through biotic controls on change in soil elevation. *Global Ecology and Biogeography* 16(5):545-556
- Moreno-Casasola P (1986) Sand movement as a factor in the distribution of plant communities in a coastal dune system. *Plant Ecology* 65: 67-76
- Mumby PJ, Edwards AJ, Arias-Gonzalez JE, Lindeman KC, Blackwell PG, Gall A, Gorczynska MI, Harborne AR, Pescod CL, Renken H, Wabnitz CC, Llewellyn G (2004) Mangroves enhance the biomass of coral reef fish communities in the Caribbean. *Nature* 427:533-536
- Rogers K, Saintilan N, Cahoon DR, (2005) Surface elevation dynamics in aregenerating mangrove forest at Homebush Bay, Australia. *Wetl. Ecol. Manage.* 13, 587–598
- Sheppard C, Dixon DJ, Gourlay M, Sheppard A, Payet R (2005) Coral mortality increases wave energy reaching shores protected by reef flats: examples from the Seychelles. *Estuarine, Coastal and Shelf Science* 64(2):223-234

- 1 Spenceley, AP (1982) Sedimentation patterns in a mangal on Magnetic Island near
2 Townsville, North Queensland, Australia. *Singapore Journal of Tropical Geography*
3 3:100-107
- 4 Stevenson NJ, Lewis RR, Burbridge PR (1999) Disused Shrimp Ponds and Mangrove
5 Rehabilitation. In: Streever W (ed) *An International Perspective on Wetland*
6 *Rehabilitation*. Springer Netherlands
- 7 Stokes DJ, Healy TR, and Cooke PJ. 2009. Surface elevation changes and sediment
8 characteristics of intertidal surfaces undergoing mangrove expansion and
9 mangrove removal, Waikaraka Estuary, Tauranga Harbour, New Zealand.
10 *International Journal of Ecology and Development* 12: 88-106.
- 11 Thampanya U, Vermaat JE, Sinsakul S, Panapitukkul N (2006) Coastal erosion and
12 mangrove progradation of Southern Thailand. *Estuarine, Coastal and Shelf Science*
13 68:75-85
- 14 Valiela I, Bowen JL, York JK (2001) Mangrove Forests: One of the World's Threatened
15 Major Tropical Environments. *BioScience* 51:807-815
- 16 Victor S, Neth L, Golbuu Y, Wolanski E, Richmond RH (2006) Sedimentation in
17 mangroves and coral reefs in a wet tropical island, Pohnpei, Micronesia. *Estuarine,*
18 *Coastal and Shelf Science* 66:409-416
- 19 Williams N (2005) Tsunami insight to mangrove value. *Current Biology* 15:R73
- 20 Wolanski E, Ridd P (1986) Tidal mixing and trapping in mangrove swamps. *Estuarine,*
21 *Coastal and Shelf Science* 23:759-771
- 22 Woodruffe CD (2002) *Coasts: form, process and evolution*. Cambridge University Press,
23 Cambridge, England

Highlights:

- We compared sediment elevation changes in adjacent cleared and intact mangroves.
- Sediment elevation increased in intact areas and decreased in cleared areas.
- Cleared sites with discontinuous fringing reef experienced greater elevation loss.
- Condition of mangroves and of offshore habitat affect mangrove sediment elevation.