

Portland State University

PDXScholar

Mathematics and Statistics Faculty
Publications and Presentations

Fariborz Maseeh Department of Mathematics
and Statistics

2015

Periodic State Revivals in Commensurate Waveguide Arrays

Jovan Petrovic
Vinca Institute of Nuclear Sciences

J. J. P. Veerman
Portland State University, veerman@pdx.edu

Follow this and additional works at: https://pdxscholar.library.pdx.edu/mth_fac



Part of the [Non-linear Dynamics Commons](#), and the [Nuclear Commons](#)

Let us know how access to this document benefits you.

Citation Details

Petrovic, Jovan and Veerman, J. J. P., "Periodic State Revivals in Commensurate Waveguide Arrays" (2015). *Mathematics and Statistics Faculty Publications and Presentations*. 143.
https://pdxscholar.library.pdx.edu/mth_fac/143

This Poster is brought to you for free and open access. It has been accepted for inclusion in Mathematics and Statistics Faculty Publications and Presentations by an authorized administrator of PDXScholar. Please contact us if we can make this document more accessible: pdxscholar@pdx.edu.

Periodic state revivals in commensurate waveguide arrays

Jovana Petrovic¹ and Peter J. J. Veerman^{2,3}

¹*Vinca Institute of Nuclear Sciences, Mike Alasa 12-14, 11000 Belgrade, Serbia*

²*Fariborz Maseeh Dept. of Math. and Stat., Portland State Univ., Portland, OR, USA*

³*CCQCN, Dept of Physics, University of Crete, 71003 Heraklion, Greece*

E-mail: jovanap@vin.bg.ac.rs

Emerging optical and quantum computers require hardware capable of coherent transport of and operations on quantum states. Here, we investigate finite optical waveguide arrays with linear coupling as means of efficient and compact coherent state transfer. Coherent transfer with periodic state revivals is enabled by engineering coupling coefficients between neighbouring waveguides to yield commensurate eigenvalue spectrum. Particular cases of finite arrays have been actively studied to achieve the perfect state transfer by mirroring the input into the output state [1, 2]. We explore a much wider scope of coherent propagation and revivals of both the state amplitude and phase.

We analytically solve the inverse eigenvalue problem to find the corresponding array coupling coefficients and use them to construct optical waveguide arrays that support full state revivals. We present analytical solutions for general arrays with 4 and 5 waveguides and for symmetric arrays with 7 and 9 waveguides [3]. These solutions include previously proposed families of solutions based on equidistance between eigenvalues.

Applications of analytic solutions are numerically demonstrated on experimentally accessible optical waveguide arrays, Fig. 1. Coupling coefficients are controlled by controlling inter-waveguide separations. It is shown that the same array can perform coherent transport of a vector state and different coupling functions if different input ports are excited [4]. It is further shown that the coupler output is sensitive to fabrication imperfections and that the accuracy required is nearly within the reach of the state-of-the-art laser writing techniques [5]. The proposed WGAs can also be of interest as directional 1xN couplers in optical circuits, input and output couplers in integrated multipath interferometers and simulators of atomic angular momenta.




Fig. 1 Numerical simulation of a commensurate 5-waveguide array with a) the input state $(1, 0, 0, 0)$ and arbitrary length, b) the input state $(0, 1, 0, 0)$ and the length that renders equal 1x4 coupler.

Besides the mathematical challenge to find analytical solutions arrays with larger and even number of waveguides, there is a considerable interest in coherent transfer through two dimensional arrays (optical lattices) and closed chains (multicore fibres), which are outlined as compelling subjects for future work.

[1] R. Gordon, "Harmonic oscillation in a spatially finite array waveguide," *Opt. Lett.* 29, 2752-2754 (2004)

[2] I. D. Chremmos and N. K. Efremidis, "A note on perfect revivals infinite waveguide arrays," *Opt. Comms.* 285, 4364-4367 (2012)

[3] P. J. J. Veerman and J. Petrovic, "Full State Revivals in Optical Waveguide Arrays with Commensurate Eigenspectra", <http://arxiv.org/abs/1507.04154> (2015)

[4] J. Petrovic, "Multiport waveguide couplers with periodic energy exchange," *Opt. Lett.* 40, 139-142 (2015)

[5] A. Perez-Leija, R. Keil, A. Kay, H. Moya-Cessa, S. Nolte, L.-C. Kwek, B. M. Rodriguez-Lara, A. Szameit, and D. N. Christodoulides, *Phys. Rev. A* 87, 012309 (2013)