

Portland State University

PDXScholar

Student Research Symposium

Student Research Symposium 2019

May 7th, 11:00 AM - 1:00 PM

Arbuscular Mycorrhizal Fungi Associations Across Multiple Saltmarsh Plant Species Mitigating the Impacts of Sea-Level Rise

Rayna Koberstein
Portland State University

Catherine E. de Rivera
Portland State University

Follow this and additional works at: <https://pdxscholar.library.pdx.edu/studentsymposium>



Part of the [Environmental Sciences Commons](#)

Let us know how access to this document benefits you.

Koberstein, Rayna and de Rivera, Catherine E., "Arbuscular Mycorrhizal Fungi Associations Across Multiple Saltmarsh Plant Species Mitigating the Impacts of Sea-Level Rise" (2019). *Student Research Symposium*. 21.

<https://pdxscholar.library.pdx.edu/studentsymposium/2019/Posters/21>

This Poster is brought to you for free and open access. It has been accepted for inclusion in Student Research Symposium by an authorized administrator of PDXScholar. Please contact us if we can make this document more accessible: pdxscholar@pdx.edu.



Arbuscular Mycorrhizal Fungi Associations Across Multiple Saltmarsh Plant Species Mitigating the Impacts of Sea-Level Rise

Rayna E. Koberstein*, Catherine E. de Rivera

Department of Environmental Science and Management, Portland State University

Background and Goals

- Salt marshes provide important habitat and ecosystem functions and services.
- Sea level rise will increase the elevation and rate of terrestrial salt water inundation, thereby reducing the ecosystem services provided by salt marshes.
- Marsh managers need information about what plants may be most successful in marsh restoration and mitigation of sea level rise.
- Halophytic plants within salt marshes can benefit from symbiotic arbuscular mycorrhizal fungi (AMF) facilitations through increased inundation tolerance, greater nutrient availability and uptake and alleviation from saline stress.
- This study aims to identify which Oregon salt marsh plants have AMF to help with future maintenance and restoration of marshes.




Research Question

- What plant species host AMF, and what is the pattern of distribution across the low and high marsh areas?

Expected Results

- There may be higher rates of AMF in dominant plants in the low marsh and lower rates of AMF in plants which lack biological saline and hydrophytic adaptations.
- Soil parameters in samples containing AMF will have lower saline content, moisture and a neutral pH. The proportion of AMF spores will be homogeneous across all samples.

What is Arbuscular Mycorrhizal Fungi (AMF)?



Hyphae
Arbuscule
Vesicle


- **Arbuscular Mycorrhizal Fungi:** A fungi which receives carbon nutrition from the roots of its associated plant. In return, it can increase nutrient uptake and inundation resilience to the plant.
- **Hyphae:** The AMF's filaments that absorb the mineral nutrients from soil and transports them to the host plant.
- **Vesicle:** Storage structure for nutrients collected by hyphae.
- **Arbuscule:** Sites of exchange within hyphae for phosphorus, carbon, water, and other nutrients (9).

AMF connects plant roots with soil; increases root surface area; allows fungus to uptake nutrients to share with plants.



Methods

- Extracted AMF from eight monoculture plots from four plant species in high and low marsh plots.
- Dyed extracted roots with Trypan blue before mounting them on slides.
- Will calculate AMF spores in soils at each site.
- Soil samples will be analyzed for percent water, soil salinity and pH.



Outcomes


- Determining the presence of AMF across plant species will help future restoration efforts build resilience against sea level rise.

Acknowledgements

- Special thanks to Dr. Catherine de Rivera, Dr. Sarah Eppley, Dr. Martin Lafrenz, Elinore Webb, Vanessa Robertson-Rojas, Katie North, Ajelabi Olamide Oladayo, Korin Kiser, the Bushby Foundation and the PSU McNair Scholars team.



Jaumea
(*Jaumea carnosa*)



Lyngbye's sedge
(*Carex lyngbyei*)



Pickleweed
(*Salicornia virginica*)



Salt Grass
(*Distichlis spicata*)

References

1. Ailangharzad, Nasser & N. Rastin, Saleh & Towfighi, Hasan & Alizadeh, A. (2001). Occurrence of arbuscular mycorrhizal fungi in saline soils of the Tabriz Plain of Iran in relation to some physical and chemical properties of soil. *Mycorrhiza*, 11, 119-122.
2. Bathe, Kelly E., et al. "Seasonal Dynamics of Arbuscular Mycorrhizal Fungi in Differing Wetland Habitats." *Mycorrhiza*, vol. 14, no. 5, Nov. 2004, pp. 329-337.
3. Cavalho, I.M., Correia, P.M., Rye, R.L., and Martins-Loucao, M.A. (2003) Spatial variability of arbuscular mycorrhizal fungal spores in two natural plant communities. *Plant Soil* 251: 227-236.
4. Daleo, Pedro, et al. "Mycorrhizal Fungi Determine Salt Marsh Plant Zonation Depending on Nutrient Supply." *Journal of Ecology*, vol. 96, no. 3, Nov. 2008, pp. 431-437.
5. Eppley, S.M. 2005. Gender-specific selection during early life-history stages in the dioecious grass *Distichlis spicata*. *Ecology* 82: 2002-2031.
6. Eppley, S.M., CA Mercer, C. Haaning, CB Graves. 2009. Sex-specific variation in the interaction between *Distichlis spicata* (Poaceae) and mycorrhizal fungi. *Am J Bot* 96:1967-1973.
7. Ewlin, H. R. Kapoor, & Giri. 2009. Arbuscular mycorrhizal fungi in alleviation of salt stress: a review. *Ann Bot* 104:1263-1280.
8. Frenkel, Robert L., and Janet C. Morlan. "Can We Restore Our Salt Marshes? Lessons from the Salmon River, Oregon." *Biological Conservation*, vol. 63, no. 1, 1991, p. 104.
9. H. Hoefnagels, Mariëtte & Broome, Stephen & Shafer, Steven. (1993). Vesicular-Arbuscular Mycorrhizae in Salt Marshes in North Carolina. *Estuaries and Coasts*. 16: 851-858. 10.2307/1352444.
9. Marschner H, B Dell. 1994. Nutrient uptake in mycorrhizal symbiosis. *Plant Soil* 159:89-102.
10. Middleton, Beth A., Proffitt, C. Edward, et al. "Seed Flootation and Germination of Salt Marsh Plants: The Effects of Stratification, Salinity, and/or Inundation Regime." *Aquatic Botany*, vol. 81, no. 3, 2009, pp. 40-46.
9. Porcel R, R Arca, JM Ruiz-Lozano. 2012. Salinity stress alleviation using arbuscular mycorrhizal fungi: a review. *Agron Sustain Dev* 32:181-200.
10. Reuss-Schmidt, Kassandria, et al. "Effects of Sex and Mycorrhizal Fungi on Gas Exchange in the Dioecious Salt Marsh Grass *Distichlis spicata*." *International Journal of Plant Sciences*, vol. 176, no. 2, 2015, pp. 141-149.
11. Rosta, Juan L., et al. "Spatial Distribution of Arbuscular Mycorrhizal Fungi in the Rhizosphere of the Salt Marsh *Plantainia Cochimoides*, Along a Salinity Gradient." *Arid Land Research and Management*, vol. 22, no. 4, July 2008, pp. 310-315.
9. Schmitz, D., Dannenberg, G., Handberg, B., Klingner, A., and Bathe, H. (1991) Quantification of vesicular-arbuscular mycorrhiza by biochemical parameters. *J Plant Physiol* 139: 106-114.
10. Smith, S.E., & Smith, 2011. Roles of arbuscular mycorrhizas in plant nutrition and growth: new paradigms from cellular to ecosystem scales. *Annual Review Plant Biology* 62:227-250.
11. Xu, Zhongyong, et al. "Arbuscular Mycorrhizal Fungi in Wetland Habitats and Their Application in Constructed Wetland: A Review." *Pedosphere*, vol. 26, no. 5, 2016, pp. 592-617.

