

3-2018

Draft Genome Sequence of a *Dictyoglomus* sp. from an Enrichment Culture of a New Zealand Geothermal Spring

Anna-Louise Reysenbach

Portland State University, reysenbacha@pdx.edu

John A. Donaho

Oak Ridge National Laboratory, Oak Ridge, Tennessee, USA

John F. Kelley

Portland State University

Emily St. John

Portland State University


Christina Turner

Portland State University

See next page for additional authors

Let us know how access to this document benefits you.

Follow this and additional works at: https://pdxscholar.library.pdx.edu/bio_fac

 Part of the [Biology Commons](#), and the [Genetics and Genomics Commons](#)

Citation Details

Reysenbach, A. L., Donaho, J. A., Kelley, J. F., John, E. S., Turner, C., Podar, M., & Stott, M. B. (2018). Draft Genome Sequence of a *Dictyoglomus* sp. from an Enrichment Culture of a New Zealand Geothermal Spring. *Genome announcements*, 6(11), e00150-18.

This Article is brought to you for free and open access. It has been accepted for inclusion in Biology Faculty Publications and Presentations by an authorized administrator of PDXScholar. For more information, please contact pdxscholar@pdx.edu.

Authors

Anna-Louise Reysenbach, John A. Donaho, John F. Kelley, Emily St. John, Christina Turner, Mircea Podar, and Matthew B. Stott



Draft Genome Sequence of a *Dictyoglomus* sp. from an Enrichment Culture of a New Zealand Geothermal Spring

Anna-Louise Reysenbach,^a John A. Donaho,^a John F. Kelley,^a Emily St. John,^a Christina Turner,^a Mircea Podar,^b Matthew B. Stott^c

^aBiology Department, Portland State University, Portland, Oregon, USA

^bOak Ridge National Laboratory, Oak Ridge, Tennessee, USA

^cSchool of Biological Sciences, University of Canterbury, Christchurch, New Zealand

ABSTRACT A draft genome of a novel *Dictyoglomus* sp., NZ13-RE01, was obtained from a New Zealand hot spring enrichment culture. The 1,927,012-bp genome is similar in both size and G+C content to other *Dictyoglomus* spp. Like its relatives, *Dictyoglomus* sp. NZ13-RE01 encodes many genes involved in complex carbohydrate metabolism.

Dictyoglomus spp. form a distinct bacterial phylum, with only two described species with genome sequences, *D. turgidum* (1) and *D. thermophilum* (2, 3). *Dictyoglomus* spp. have been detected in thermophilic environments, such as terrestrial hot springs (2, 4–8) and paper pulp factory effluent (9), and they have an unusual morphology consisting of large spherical bodies assembled from bundles of filamentous cells (2, 5, 9). *Dictyoglomus* spp. are of industrial interest due to their capability to ferment a wide range of complex carbohydrates, including starch, cellulose, xylan, and pectin (2, 8, 10, 11). Due to the low number of representative genomes, their phylogenetic placement within the domain *Bacteria* is not certain as *Dictyoglomus* spp. are, although they appear to be affiliated most closely with a cluster containing *Coprothermobacter*, *Synergistes*, *Acetothermia*, and *Thermotogales* (1, 12, 13).

We obtained a genome of a novel *Dictyoglomus* sp., NZ13-RE01, from a New Zealand hot spring enrichment culture (Hell's Gate, Tikitere, New Zealand, 38°03'47"S, 176°21'39"E, pH 6.0, 74°C). Enrichments were incubated at 80°C for 4 days in a modified anaerobic DSMZ medium (no. 88) containing yeast extract (0.5 g/liter) and tryptone (0.5 g/liter). Spherical bodies were apparent in the enrichment cultures by phase-contrast microscopy. DNA was extracted using the Qiagen DNeasy blood and tissue kit. Metagenome Nextera DNA libraries were sequenced on the Illumina MiSeq platform. Adapters and low-quality reads were trimmed using Trimmomatic (14), reads were assembled with IDBA-UD version 1.1.0 (15, 16), and contigs ≥ 1 kb were binned using MaxBin version 1.4.5 (17). To optimize the assembly, the reads were mapped back to the *Dictyoglomus* sp. NZ13-RE01 draft genome using Bowtie2 version 2.2.5 (18) and SAMtools version 1.2 (19, 20) and reassembled with IDBA-UD. The genome was further curated using emergent self-organizing maps (21). Open reading frames were annotated using the Rapid Annotations using Subsystems Technology (RAST) server (22–24), the NCBI Prokaryotic Genome Annotation Pipeline (PGAP) (25), the Clusters of Orthologous Groups of proteins (COG) database (26), and the dbCAN database (27). tRNAs were predicted with tRNAscan-SE version 2.0 (28).

Based on CheckM (29) analysis, the *Dictyoglomus* sp. NZ13-RE01 genome is about 100% complete with no contamination. The 1,927,012-bp genome consists of 34 contigs, with a 33% G+C content, 1,870 predicted protein-coding genes, and 48 tRNAs. The *Dictyoglomus* sp. NZ13-RE01 genome has an average nucleotide identity (ANI) score of 74% and an average amino acid identity (AAI) score of 65% compared to the

Received 3 February 2018 Accepted 13 February 2018 Published 15 March 2018

Citation Reysenbach A-L, Donaho JA, Kelley JF, St John E, Turner C, Podar M, Stott MB. 2018. Draft genome sequence of a *Dictyoglomus* sp. from an enrichment culture of a New Zealand geothermal spring. Genome Announc 6: e00150-18. <https://doi.org/10.1128/genomeA.00150-18>.

Copyright © 2018 Reysenbach et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Anna-Louise Reysenbach, reysenbacha@pdx.edu.

genomes of both *D. turgidum* and *D. thermophilum* (30, 31). Further, Circos synteny plots (32) show that the *Dictyoglomus* sp. NZ13-RE01 genome does not share the highly syntenic genome arrangement found between *D. turgidum* and *D. thermophilum* (1). Considering the dissimilarity in genome nucleotide and amino acid sequences, and the low synteny in genome arrangement between NZ13-RE01 and either of the described *Dictyoglomus* spp., it is likely that NZ13-RE01 represents a new species within the genus *Dictyoglomus*.

The *Dictyoglomus* sp. NZ13-RE01 genome encodes an extensive suite of carbohydrate metabolism genes (63 glycosyl hydrolases, 10 carbohydrate esterases, and 28 glycosyltransferases), including α -amylases, α -xylosidases, a chitinase, endo-1,4- β xylanases, and a β -mannanase. Like other *Dictyoglomus* spp., the genome has a reverse gyrase. The sporulation gene, *spoVS*, was also present and has been proposed to play a role in morphological changes in *Dictyoglomus* spp. (1).

Accession number(s). The nucleotide genome sequence reported here has been deposited in DDBJ/ENA/GenBank under the accession no. [NIRF0000000](https://doi.org/10.3389/fmicb.2016.01979). The version described in this paper is the first version, NIRF01000000.

ACKNOWLEDGMENTS

We thank Yitai Liu for assistance with the enrichment cultures and Kristen Brileya and Jennifer Meneghin for assistance in the initial stages of the project. We thank Haley Nasman for her assistance in metagenomic analysis. We also thank the Tikitere Trust (Whakapoungakau 24) for its continued support and for sampling access in the Hell's Gate geothermal area.

This work was funded by the National Science Foundation (grant no. DEB 1134877 to A.-L.R. and M.P.) and by the Geothermal Resources of New Zealand research program at GNS Science (to M.B.S.).

REFERENCES

1. Brumm PJ, Gowda K, Robb FT, Mead DA. 2016. The complete genome sequence of hyperthermophile *Dictyoglomus turgidum* DSM 6724TM reveals a specialized carbohydrate fermentor. *Front Microbiol* 7. <https://doi.org/10.3389/fmicb.2016.01979>.
2. Saiki T, Kobayashi Y, Kawagoe K, Beppu T. 1985. *Dictyoglomus thermophilum* gen. nov., sp. nov., a chemoorganotrophic, anaerobic, thermophilic bacterium. *Int J Syst Evol Microbiol* 35:253–259. <https://doi.org/10.1099/00207713-35-3-253>.
3. Coil DA, Badger JH, Forberger HC, Riggs F, Madupu R, Fedorova N, Ward N, Robb FT, Eisen JA. 2014. Complete genome sequence of the extreme thermophile *Dictyoglomus thermophilum* H-6-12. *Genome Announc* 2(1): e00109–14. <https://doi.org/10.1128/genomeA.00109-14>.
4. Patel BK, Morgan HW, Wiegel J, Daniel RM. 1987. Isolation of an extremely thermophilic chemoorganotrophic anaerobe similar to *Dictyoglomus thermophilum* from New Zealand hot springs. *Arch Microbiol* 147:21–24. <https://doi.org/10.1007/BF00492899>.
5. Hoppert M, Valdez M, Enseleit M, Theilmann W, Valerius O, Braus GH, Föst C, Liebl W. 2012. Structure-functional analysis of the *Dictyoglomus* cell envelope. *Syst Appl Microbiol* 35:279–290. <https://doi.org/10.1016/j.syapm.2012.06.004>.
6. Kublanov IV, Perevalova AA, Slobodkina GB, Lebedinsky AV, Bidzhieva SK, Kolganova TV, Kaliberda EN, Rumsh LD, Haertlé T, Bonch-Osmolovskaya EA. 2009. Biodiversity of the thermophilic prokaryotes with hydrolytic activities in hot springs of Uzon Caldera, Kamchatka (Russia). *Appl Environ Microbiol* 75:286–291. <https://doi.org/10.1128/AEM.00607-08>.
7. Svetlichny VA, Svetlichnaya TP. 1988. *Dictyoglomus turgidus* sp. nov., a new extremely thermophilic eubacterium isolated from hot springs of the Uzon volcano caldera. *Mikrobiologi* 57:435–441.
8. Mathrani IM, Ahring BK. 1992. Thermophilic and alkalophilic xylanases from several *Dictyoglomus* isolates. *Appl Microbiol Biotechnol* 38:23–27. <https://doi.org/10.1007/BF00169413>.
9. Mathrani IM, Ahring BK. 1991. Isolation and characterization of a strictly xylan-degrading *Dictyoglomus* from a man-made, thermophilic anaerobic environment. *Arch Microbiol* 157:13–17. <https://doi.org/10.1007/BF00245328>.
10. Brumm P, Hermanson S, Hochstein B, Boyum J, Hermersmann N, Gowda K, Mead D. 2011. Mining *Dictyoglomus turgidum* for enzymatically active carbohydrases. *Appl Biochem Biotechnol* 163:205–214. <https://doi.org/10.1007/s12010-010-9029-6>.
11. Kobayashi Y, Motoike M, Fukuzumi S, Ohshima T, Saiki T, Beppu T. 1988. Heat-stable amylase complex produced by a strictly anaerobic and extremely thermophilic bacterium, *Dictyoglomus thermophilum*. *Agric Biol Chem* 52:615–616. <https://doi.org/10.1271/abb1961.52.615>.
12. Love CA, Patel BKC, Ludwig W, Stackebrandt E. 1993. The phylogenetic position of *Dictyoglomus thermophilum* based on 16S rRNA sequence analysis. *FEMS Microbiol Lett* 107:317–320. <https://doi.org/10.1111/j.1574-6968.1993.tb06050.x>.
13. Nishida H, Beppu T, Ueda K. 2011. Whole-genome comparison clarifies close phylogenetic relationships between the phyla *Dictyoglomi* and *Thermotogae*. *Genomics* 98:370–375. <https://doi.org/10.1016/j.ygeno.2011.08.001>.
14. Bolger AM, Lohse M, Usadel B. 2014. Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics* 30:2114–2120. <https://doi.org/10.1093/bioinformatics/btu170>.
15. Peng Y, Leung HCM, Yiu S-M, Chin FYL. 2010. IDBA—a practical iterative de Bruijn graph de novo assembler, p. 426–440. *In Annual International Conference on Research in Computational Molecular Biology*, Springer, New York, NY. https://doi.org/10.1007/978-3-642-12683-3_28.
16. Peng Y, Leung HCM, Yiu SM, Chin FYL. 2012. IDBA-UD: a de novo assembler for single-cell and metagenomic sequencing data with highly uneven depth. *Bioinformatics* 28:1420–1428. <https://doi.org/10.1093/bioinformatics/bts174>.
17. Wu Y-W, Tang Y-H, Tringe SG, Simmons BA, Singer SW. 2014. MaxBin: an automated binning method to recover individual genomes from metagenomes using an expectation-maximization algorithm. *Microbiome* 2:26. <https://doi.org/10.1186/2049-2618-2-26>.
18. Langmead B, Trapnell C, Pop M, Salzberg SL. 2009. Ultrafast and memory-efficient alignment of short DNA sequences to the human genome. *Genome Biol* 10:R25. <https://doi.org/10.1186/gb-2009-10-3-r25>.
19. Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis G, Durbin R. 1000 Genome Project Data Processing Subgroup.

2009. The Sequence Alignment/Map format and SAMtools. *Bioinformatics* 25:2078–2079. <https://doi.org/10.1093/bioinformatics/btp352>.
20. Li H. 2011. A statistical framework for SNP calling, mutation discovery, association mapping and population genetical parameter estimation from sequencing data. *Bioinformatics* 27:2987–2993. <https://doi.org/10.1093/bioinformatics/btr509>.
21. Ultsch A, Mörchen F. 2005. ESOM-Maps: tools for clustering, visualization, and classification with Emergent SOM. Technical Report, no 46 Department of Mathematics and Computer Science, University of Marburg, Marburg, Germany.
22. Brettin T, Davis JJ, Disz T, Edwards RA, Gerdes S, Olsen GJ, Olson R, Overbeek R, Parrello B, Pusch GD, Shukla M, Thomason JA, III, Stevens R, Vonstein V, Wattam AR, Xia F. 2015. RASTtk: a modular and extensible implementation of the RAST algorithm for building custom annotation pipelines and annotating batches of genomes. *Sci Rep* 5:8365. <https://doi.org/10.1038/srep08365>.
23. Overbeek R, Olson R, Pusch GD, Olsen GJ, Davis JJ, Disz T, Edwards RA, Gerdes S, Parrello B, Shukla M, Vonstein V, Wattam AR, Xia F, Stevens R. 2014. The SEED and the Rapid Annotation of microbial genomes using Subsystems Technology (RAST). *Nucleic Acids Res* 42:D206–D214. <https://doi.org/10.1093/nar/gkt1226>.
24. Aziz RK, Bartels D, Best AA, DeJongh M, Disz T, Edwards RA, Formsma K, Gerdes S, Glass EM, Kubal M, Meyer F, Olsen GJ, Olson R, Osterman AL, Overbeek RA, McNeil LK, Paarmann D, Paczian T, Parrello B, Pusch GD, Reich C, Stevens R, Vassieva O, Vonstein V, Wilke A, Zagnitko O. 2008. The RAST server: Rapid Annotations using Subsystems Technology. *BMC Genomics* 9:75. <https://doi.org/10.1186/1471-2164-9-75>.
25. Tatusova T, DiCuccio M, Badretdin A, Chetvernin V, Nawrocki EP, Zaslavsky L, Lomsadze A, Pruitt KD, Borodovsky M, Ostell J. 2016. NCBI prokaryotic genome annotation pipeline. *Nucleic Acids Res* 44: 6614–6624. <https://doi.org/10.1093/nar/gkw569>.
26. Tatusov RL, Galperin MY, Natale DA, Koonin EV. 2000. The COG database: a tool for genome-scale analysis of protein functions and evolution. *Nucleic Acids Res* 28:33–36. <https://doi.org/10.1093/nar/28.1.33>.
27. Yin Y, Mao X, Yang J, Chen X, Mao F, Xu Y. 2012. dbCAN: a Web resource for automated carbohydrate-active enzyme annotation. *Nucleic Acids Res* 40:W445–W451. <https://doi.org/10.1093/nar/gks479>.
28. Lowe TM, Chan PP. 2016. tRNAscan-SE on-line: integrating search and context for analysis of transfer RNA genes. *Nucleic Acids Res* 44: W54–W57. <https://doi.org/10.1093/nar/gkw413>.
29. Parks DH, Imelfort M, Skennerton CT, Hugenholtz P, Tyson GW. 2015. CheckM: assessing the quality of microbial genomes recovered from isolates, single cells, and metagenomes. *Genome Res* 25:1043–1055. <https://doi.org/10.1101/gr.186072.114>.
30. Rodriguez-R LM, Konstantinidis KT. 2016. The enveomics collection: a toolbox for specialized analyses of microbial genomes and metagenomes. *PeerJ* 4:e1900v1. <https://doi.org/10.7287/peerj.preprints.1900v1>.
31. Konstantinidis KT, Rosselló-Móra R, Amann R. 2017. Uncultivated microbes in need of their own taxonomy. *ISME J* 11:2399–2406. <https://doi.org/10.1038/ismej.2017.113>.
32. Krzywinski M, Schein J, Birol I, Connors J, Gascoyne R, Horsman D, Jones SJ, Marra MA. 2009. Circos: an information aesthetic for comparative genomics. *Genome Res* 19:1639–1645. <https://doi.org/10.1101/gr.092759.109>.