

Portland State University

PDXScholar

Psychology Faculty Publications and
Presentations

Psychology

6-26-2022

Family-Supportive Supervisor Behaviors and Psychological Distress: A Secondary Analysis across Four Occupational Populations

Philip G. Bouleh

Portland State University, pbouleh@pdx.edu

Shalene Joyce Allen

Portland State University, s_allen_92@hotmail.com

Leslie Hammer

Portland State University, hammerl@pdx.edu

Follow this and additional works at: https://pdxscholar.library.pdx.edu/psy_fac

Let us know how access to this document benefits you.

Citation Details

Bouleh, P. G., Allen, S. J., & Hammer, L. B. (2022). Family-Supportive Supervisor Behaviors and Psychological Distress: A Secondary Analysis across Four Occupational Populations. *International Journal of Environmental Research and Public Health*, 19(13), 7845.

This Article is brought to you for free and open access. It has been accepted for inclusion in Psychology Faculty Publications and Presentations by an authorized administrator of PDXScholar. Please contact us if we can make this document more accessible: pdxscholar@pdx.edu.



Article

Family-Supportive Supervisor Behaviors and Psychological Distress: A Secondary Analysis across Four Occupational Populations

Philip G. Bouleh ¹, Shalene J. Allen ¹ and Leslie B. Hammer ^{2,*}

¹ Department of Psychology, Portland State University, Portland, OR 97207, USA; pbouleh@pdx.edu (P.G.B.); shallen@pdx.edu (S.J.A.)

² Oregon Institute of Occupational Health Sciences, Oregon Health & Science University, Portland, OR 97239, USA

* Correspondence: hammerl@ohsu.edu

Abstract: This study assessed the associations of employee's perceptions of family-supportive supervisor behaviors (FSSB) and their psychological distress across four occupational populations ($n = 3778$): information technology; healthcare; military-connected Veterans; and National Guard service members. Data were gathered and analyzed from four larger archival datasets to compare differences in these relationships. Results revealed significant negative relationships between employee reports of FSSB and their psychological distress within occupations, as expected. Furthermore, results revealed significant differences across occupational populations for employee reports of both FSSB and psychological distress. Hierarchical moderated multiple regression analyses were conducted to examine the extent of these mean differences across groups. Results revealed significant differences among these four groups such that the military-connected Veteran employees demonstrated significantly stronger associations of FSSB, and psychological distress compared to the other three occupations of information technology, healthcare, and National Guard service members. These findings suggest the importance of FSSB to worker psychological health across a variety of occupational populations, specifically noting the importance and presence of FSSB for Veteran employees' psychological distress in civilian workplaces. Practical implications include the need for training leaders on how to better support employees' work and non-work lives, mental health, and well-being.

Keywords: psychological distress; FSSB; occupational stress; mental health; social support



Citation: Bouleh, P.G.; Allen, S.J.; Hammer, L.B. Family-Supportive Supervisor Behaviors and Psychological Distress: A Secondary Analysis across Four Occupational Populations. *Int. J. Environ. Res. Public Health* **2022**, *19*, 7845. <https://doi.org/10.3390/ijerph19137845>

Academic Editors: Michael Kenneth Lemke, Adam Hege and Brian Whitaker

Received: 11 May 2022

Accepted: 21 June 2022

Published: 26 June 2022

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2022 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

1.1. Psychological Distress and Work-Family Concerns

With both work-related challenges and environmental disasters, the psychological health and well-being of employees and their families is of growing concern in today's society [1]. In fact, many employees have recently faced significant impacts of distress as result of the COVID pandemic, ultimately affecting the way they are able to perform at work [2]. As such, there is a great need to understand how we can protect and support workers' mental health and well-being [3], specifically, by examining factors that can improve their reports of psychological distress. For instance, mounting evidence suggests employees have experienced challenges managing their work and family responsibilities due to the growing incorporation of a 24/7 economy and the demands placed upon single parents and dual-earner couples who may be caring for children and/or elderly family members [4–7]. Such work-family conflict emerges when employees struggle to simultaneously fulfill their overlapping responsibilities between work and home life, and in turn problematically sacrifice the energy and time they invest in one domain for the other [8]. As a consequence, employees who experience greater levels of work–family conflict and decreased levels of organizational support are at greater risk of having increased negative affective experiences, such as anxiety and distress, as well as diminished organizational effectiveness [9,10]. For example, mothers who manage multiple roles composed of higher childcare responsibility

in addition to professional work-related duties experience greater levels of psychological distress, depression and absenteeism, and lower levels of well-being [11,12]. This is especially relevant given that psychosocial occupational stressors (e.g., high work demands, long work hours, and low job control) have been consistently linked to poor psychological and physical health outcomes (e.g., cardiovascular disease, mortality, type 2 diabetes, depression) [13,14]. In consideration of the consequential relationship between work-family conflict and employee health and well-being, it is important to examine workplace conditions that can impact and alleviate the conflict between work–non-work domains and distress. For example, supervisor support for family is one factor that has been found to improve work–non-work lives of employees.

1.2. Family-Supportive Supervisor Behaviors (FSSB)

Social support has been extensively studied in the organizational science and occupational health literatures and is generally found to mitigate the bi-directional interference between work and family [15–20]. In particular, the provision of social support in the workplace via supervisors denotes a vehicle, which can provide important resources to employees [6]. FSSB is a more content-specific formulation of supervisor support that consists of four behavioral dimensions that can be implemented by supervisors to assist in balancing work-family demands [19,21]. The four dimensions comprising FSSB are emotional support, instrumental support, role modeling behaviors, and creative work-family management—all of which are characterized by a distinct set of leadership behaviors that improve the experience between the work and non-work lives of employees. For example, leaders express *emotional support* when they engage in authentic interpersonal interactions with their employees while creating a safe space to discuss and sympathize with their personal and family commitments. *Role modeling* behaviors are manifested as the exemplification of work–family balance in the daily life of a leader. *Instrumental support* consists of a manager proactively helping an employee manage work and family challenges, such as adjusting an employee’s schedule in response to their expressed work and family scheduling conflicts. Finally, *creative work-family management* is the proactive and innovative process, by which leadership collaborates with employees to restructure work for the purpose of maximizing the benefits experienced at work and at home [19,21]. Employee reports of FSSB have been correlated with important health, family, and workplace outcomes [22–24].

Training leaders on the importance of FSSB and in particular, specific behaviors to enact, has been shown to improve employee well-being outcomes [25,26]. For instance, research suggests that workplace contextual factors such as family-supportive organizational culture, work-family infrastructures, as well as individual-level factors (e.g., family life stage) contribute to FSSB [27]. This suggests that certain supervisor populations may be more or less likely to enact, and certain worker populations may be more or less responsive to, FSSB. Furthermore, certain occupations may be more or less conducive to FSSB. For example, information technology occupations are more conducive to work from home practices compared to traditional healthcare settings, and therefore, may have higher levels of FSSB enacted and received by supervisors and employees due to the nature of the work.

1.3. Current Study

In alignment with prior research [28], this study utilized a novel common measures methodological approach to investigate a repository of cross-sectional data from several large-scale studies for the purpose of examining the relationship between employee reports of FSSB and their reports of psychological distress among workers in various occupations. The goal of this study is to understand the relationship between supervisor support (i.e., FSSB) and psychological distress across diverse organizational contexts. Since FSSB is influenced by individual and contextual factors [27], this study provides insight into how different populations of workers perceive the presence of FSSB in their particular workplace, and how it relates to their psychological well-being.

Given FSSB principles are based in social support theory and literature [29], we anticipate that increased employee perceptions of FSSB will be related to decreases in their reports of psychological distress and improved employee psychological health and well-being. Social connections and connectedness are key levers in impacting psychological and physical health [30,31]. Therefore, it is important to understand not only the relationship between FSSB and psychological distress but discerning how this relationship may differ across occupational populations may point researchers and practitioners to workplace conditions that may be altered to facilitate higher levels of FSSB, which in turn, could help reduce psychological distress. As Straub [27] suggests, antecedents of FSSB include workplace contextual variables such as family-supportive organizational culture, top management openness, the organizational reward system, and access to work family infrastructure. While these contextual variables may explain differences in occupations that can impact levels of FSSB, when examining different occupational populations Straub [27] also suggests that individual factors such as life course stage, family life stage, social identification, and gender roles may also impact the enactment of FSSB. Thus, we aim to shed light on and explore how employee reports of FSSB and their psychological distress differs by occupational population. We examined the following hypothesis and research questions:

Hypothesis 1: *There will be a significant negative association between employee reports of FSSB and psychological distress across and within occupational populations.*

Research Question 1a: Is there a significant difference in employee reports of FSSB by occupational population?

Research Question 1b: Is there a significant difference in employee reports of psychological distress by occupational population?

Research Question 2: Does the relationship between employee reports of FSSB and psychological distress differ by occupational population?

As emphasized by Hanson and colleagues [28], the common measures methodology presents several empirical advantages given that it employs the same set of standardized measures both across and within populations. Additionally, population-based studies tend to be more expensive, may be limited to broader occupational categories, and have been lacking in their generalizability to U.S. populations given the majority have been conducted in Europe [28]. Therefore, the implementation of a common measures approach in the current study, enables a robust evaluation of the relationship between employee reports of FSSB and their psychological distress, specifically, examining data from four U.S. worker populations.

To the best of our knowledge, two prior studies utilized the common measures approach [28,32]. However, this is the first study to conduct a cross-sectional comparison utilizing baseline data to specifically examine the relationship between employee perceptions of FSSB and their psychological distress across four occupational samples. Our samples consist of military-connected and civilian samples, which lends to a unique assessment of the associations between perceptions of FSSB and psychological distress from National Guard service members, Veterans, healthcare, and information technology employees. This study additionally provides a benchmark for the reported rates of FSSB and psychological distress across such populations. In doing so, we provide further evidence for the generalizability of FSSB to enhance the psychological health and mental well-being of employees across multiple occupational roles, while also providing evidence for mental health prevention and intervention strategies for organizations, supervisors, and employees.

2. Materials and Methods

2.1. Overview of Study Populations

Demographic information across occupational populations revealed employees were primarily white (74.6%), female (53.7%), with an approximate age of 40 years old ($M = 39.85$, $SD = 11.10$). Below we briefly describe each study population (also see Table 1 for study and population overview). See Table 2 for a complete overview of sample characteristics and descriptive information within and across occupational populations.

Table 1. Study Breakdown and Population Overview.

Reference Number	Citation	Study Name	Study Design	Sample(s)	Method
[33]	Hammer et al., 2017	Study for Employment Retention of Veterans (SERVe)	Randomized controlled trial	Post-9/11 veterans working in civilian organizations working at least 20 h per week ($n = 512$)	A Veteran Supervisor Support Training intervention was implemented to supervisors across 42 civilian organizations to determine overall effectiveness for improving the lives of veteran employees with data collected at baseline, 3-month, and 9-months.
[22]	Hammer et al., 2021	Military Employee Sleep & Health Study (MESH)	Cluster randomized controlled trial (cRCT)	Full-time National Guard service members from Army and Air branches ($n = 919$)	Large-scale Intervention study with training content focused on FSSB and Supervisor Support for sleep health in tandem with sleep tracking and individualized sleep health feedback for employees and supervisors, with data collected at baseline, 4-month, and 9-months.
[34]	Bray et al., 2013	Work, Family, and Health Network Study (WFHS)	Group-randomized field experiment	Low and high-wage employees across two large companies from information technology ($n = 823$) and healthcare care ($n = 1524$) working 22 h or more per week.	Intensive intervention study aimed at enhancing employees' control over their work time by implementing participatory work redesign activities with training supervisors on support for employee's family lives, with data collected at baseline, 6-, 12-, and 18-months.

Table 2. Means, Standard Deviations, and Percentages for Employee Study Demographic and Work Characteristics by Occupational Group.

Variable	Combined Sample ($n = 3778$) $M (SD) \%$	Information Technology ($n = 823$) $M (SD) \%$	Healthcare ($n = 1524$) $M (SD) \%$	Veteran ($n = 512$) $M (SD) \%$	Service Member ($n = 919$) $M (SD) \%$
Female	53.7%	39.1%	91.8%	16.4%	24.2%
Male	46.3%	60.9%	8.2%	83.6%	75.6%
Age	39.85 (11.10)	45.66 (9.20)	38.46 (12.58)	39.00 (9.39)	37.39 (8.92)
Race/Ethnicity					
White	74.6%	71.2%	70.1%	80.6%	82.0%
Latinx or Hispanic	8.9%	6.6%	13.4%	1.0%	8.0%
Black or African American	6.9%	2.9%	13.9%	1.8%	1.4%
Asian or Asian Indian	6.6%	20.9%	3.4%	2.0%	1.9%
American Indian or Alaska Native	0.7%	0.1%	0.6%	1.6%	0.8%
Native Hawaiian or Pacific Islander	1.0%	1.7%	0.7%	0.2%	1.2%
More Than One Race or Other Race	8.0%	3.0%	11.2%	12.9%	4.7%
Married or Cohabiting	72.4%	79.3%	62.9%	77.7%	79.0%
Number of children	1.09 (1.20)	0.99 (1.08)	1.03 (1.18)	0.94 (1.13)	1.33 (1.30)
Some college, technical school, or degree	80.5%	96.2%	61.7%	94.5%	89.9%
Hours worked per week	40.90 (7.28)	45.42 (5.69)	36.88 (7.29)	42.36 (6.54)	42.72 (5.23)
Years at current job or tenure	7.55 (7.80)	13.34 (9.22)	5.93 (6.53)	8.05 (5.97)	4.88 (5.75)

Note: Data represents Baseline information.

Study for Employment Retention of Veterans (SERVe). SERVe was a randomized control trial (RCT) designed to evaluate the impact of a Veteran Supportive Supervisor Training on the health and well-being of Veteran employees in civilian organizations. The sample consisted of 512 baseline participants who were post-911 Veterans, employed across 35 organizations in the Pacific Northwest, indicating they were primarily white (80.6%),

males (83.6%), with an approximate age of 39 years old ($M = 39.00, SD = 9.39$). See Table 2 for demographic information. For a detailed explanation of study methodology see [33].

Military Employee Sleep and Health Study (MESH). MESH was a cluster RCT designed to improve the sleep, health and well-being of 919 baseline participants who were full-time military service members of the National Guard. Service Members indicated they were primarily white (82.0%), males (75.6%), with an approximate age of 37 years old ($M = 37.39, SD = 8.92$). See Table 2 for demographics. For a detailed description of study methods see Hammer and colleagues [22].

Work, Family, and Health Network Study (WFHS). WFHS is a National Institutes of Health (NIH, Kansas City, MO, USA) and Center for Disease Control and Prevention (CDC, Atlanta, GA, USA) funded study evaluating a work-family support intervention designed to improve the health and well-being of managers, employees, and their families. The study consisted of two baseline employee samples, namely 823 information technology (IT) employees working at a Fortune 500 firm, and 1524 healthcare employees working in long-term care positions within nursing homes. See Bray and colleagues [34] for an in-depth description of study methods.

Information Technology Sample. Of the 823 IT employee sample collected from the WFHS, participants reported they were primarily white (71.2%), males (60.9%), with an approximate age of 46 years old ($M = 45.66, SD = 9.20$). See Table 2 for a breakdown of demographic and population characteristic information.

Healthcare Sample. Of the 1524 healthcare employees working in long-term care positions who participated in the WFHS, participants primarily indicated they were white (70.1%), females (91.8%), with an approximate age of 38 years old ($M = 38.46, SD = 12.58$). See Table 2 for detailed characteristic and demographic information.

2.2. Common Measures across Study Populations

Family-Supportive Supervisor Behaviors (FSSB). Employees rated the extent to which they agreed that their direct supervisor exhibited family-supportive supervisor behaviors with the following four items on a 5-point scale (1 = *strongly disagree*, 5 = *strongly agree*): “Your supervisor makes you feel comfortable talking to him/her about your conflicts between work and non-work”, “Your supervisor demonstrates effective behaviors in how to juggle work and non-work issues”, “Your supervisor works effectively with employees to creatively solve conflicts between work and non-work”, and “Your supervisor organizes the work in your department or unit to jointly benefit employees and the company” [21]. See Table 3 for alpha coefficients.

Table 3. Means, Standard Deviations, and Reliabilities for FSSB and Psychological Distress Measures.

Employee Group	FSSB				Psychological Distress			
	M	SD	N	α	M	SD	N	α
Information Technology	3.83	1.32	821	0.88	1.81	0.54	823	0.77
Healthcare	3.65	1.10	1515	0.89	1.98	0.72	1522	0.83
Veteran	3.81	0.94	508	0.93	2.11	0.75	509	0.90
Service Member	4.10	0.94	916	0.95	1.62	0.65	900	0.87
Combined Sample	3.84	0.90	3752	0.91	1.87	0.73	3757	0.85

Note: Data represents Baseline information.

Psychological Distress scale-K6. Employees rated how often they experienced non-specific psychological distress in the last month with the following six items on a 5-point scale (1 = *None of the time*, 5 = *All of the time*): “Feel so depressed that nothing could cheer you up?”, “Feel hopeless?”, “Feel restless or fidgety?”, “Feel that everything was an effort?”, “Feel worthless?”, “Feel nervous?” [35]. See Table 3 for alpha coefficients.

2.3. Analyses

Means, standard deviations, correlations, and reliabilities for study variables were computed to create individual profiles for study samples across four populations (IT, healthcare, Veteran, service member). Initial reliabilities were computed using Cronbach's alpha across integrated measures. Pearson correlations were computed to determine if there was a significant association between employee perceptions of their supervisors' FSSB and employee self-reported psychological distress across occupational populations. A one-way analysis of variance (ANOVA) was conducted to determine the differences between means of the four occupational groups for FSSB and psychological distress measures. Additionally, to assess differences of the association of employee perceptions of FSSB and psychological distress between occupational groups, hierarchical moderated multiple regression analysis was implemented.

Data were analyzed in SPSS Version 27 for descriptives, correlations, reliabilities and for ANOVA analyses. Furthermore, data were analyzed in Mplus Version 8 [35] to assess the moderated regression models. Alpha significance was set at $p = 0.05$ for two-tailed tests for determining statistical significance. Descriptive information on demographic and work characteristics across the four study samples ($n = 3778$) is found in Table 2.

2.4. Measurement Integration

Baseline item responses across studies (WFHS, SERVe, and MESH) were combined to create composite scale scores for the 4-item FSSB-SF measure [21] and the 6-item psychological distress measure [35]. To standardize item and scale integration across populations, data discrepancies were corrected before running analyses. For example, the FSSB measure, retrieved from the WFHN study, originally sampled participants on a 5-item scale, thus, composite scores for the integrated data excluded item "Respondent and supervisor talk effectively to solve conflicts". After the FSSB scale was re-scaled to the 4-item measure, items were reversed coded and composite scores were obtained to standardize measures across groups. To integrate data for the K6 psychological distress measure item responses from the WFHN study were reverse coded for all 6-items before obtaining composite scores. In addition, item responses were originally scaled from 1 to 4 for participant data on the K6 measure from SERVe. Thus, for consistency, all 4 item response options were inflated to standardize composite scores. New scale scores were then computed for SERVe participant responses on the K6 measure prior to integration. The measures, including item responses, composites, and response options for FSSB and psychological distress were then integrated into one composite measure of FSSB and psychological distress across all study samples.

3. Results

3.1. Association of FSSB and Psychological Distress

A Pearson correlation coefficient was computed to examine the relationship between employee reports of FSSB, $M = 3.84$, $SD = 0.90$ and their self-reports of psychological distress, $M = 1.87$, $SD = 0.73$. Results indicated that there was a significant negative correlation between employee perceptions of FSSB and their psychological distress, $r = -0.20$, $p < 0.01$, 95% CI $[-0.23, -0.17]$ across all occupational populations. Approximately 4% of the variance was shared between FSSB and psychological distress in the combined sample, $r^2 = 0.04$. Pearson correlation coefficients obtained similar findings within occupational groups: IT ($r = -0.12$, $p < 0.01$, 95% CI $[-0.19, -0.05]$), healthcare workers ($r = -0.13$, $p < 0.01$, 95% CI $[-0.18, -0.08]$), Veteran employees ($r = -0.21$, $p < 0.01$, 95% CI $[-0.29, -0.13]$), and service member employees ($r = -0.18$, $p < 0.01$, 95% CI $[-0.24, -0.12]$). Thus, Hypothesis 1 was confirmed; there was a significant negative association between FSSB and psychological distress across and within occupational populations.

3.2. Mean Difference Comparisons

A one-way analysis of variance tested whether employees from the four occupational groups (IT, healthcare, Veteran, service member) differed significantly on their self-reported

perceptions of FSSB (RQ2a). Service members reported higher FSSB ($M = 4.10$, $SD = 0.94$), compared to IT employees ($M = 3.83$, $SD = 0.82$), Veterans ($M = 3.81$, $SD = 0.93$), and employees in healthcare ($M = 3.69$, $SD = 0.88$). Results indicated that these means differed significantly, $F(3,3748) = 41.54$, $p < 0.001$. The proportion of variance in FSSB accounted for by type of occupational group was approximately 3%, $\eta^2 = 0.03$. The Tukey post hoc tests indicated that reports of FSSB across service member employees and healthcare employees differed significantly from Veterans and information technology employees ($p < 0.05$).

Similarly, a one-way analysis of variance was used to test whether employees from the four occupational groups differed significantly on their self-reported psychological distress levels (RQ2b). Veteran employees reported higher psychological distress ($M = 2.11$, $SD = 1.00$), compared to healthcare employees ($M = 1.98$, $SD = 0.72$), employees in IT ($M = 1.81$, $SD = 0.54$), and service member employees ($M = 1.62$, $SD = 0.65$). Results indicated that these means differed significantly, $F(3,3756) = 70.66$, $p < 0.001$. The proportion of variance in psychological distress accounted for by type of occupational group was approximately 5%, $\eta^2 = 0.05$. The Tukey post hoc tests indicated that reports of psychological distress across all four occupational categories differed significantly from each of the groups ($p < 0.05$).

3.3. Hierarchical Moderated Multiple Regression

Finally, to determine if the relationship between employee perceptions of FSSB and psychological distress differed across employee occupational populations (RQ3), hierarchical moderated regression analyses were conducted to examine if any one or more of these group memberships impacted the strength of the relationship between these two variables. Psychological distress was modeled as the outcome and FSSB was included as the predictor for the model. The moderator was dummy coded into four dummy variables: P1 = IT; P2 = healthcare; P3 = Veteran; P4 = service member; representing four categories with the Veteran sample serving as the reference group.

Interaction terms were added into the second model and effects were constrained across the groups to assess and compare the full vs. reduced model results for the reference group (i.e., Veteran sample) compared to the other three occupational categories. Results of the categorical moderation revealed the relationship between FSSB and psychological distress was significantly different for the Veteran sample compared to the other three occupational groups, as indicated by the significant $\Delta R^2 = 0.002$, ($\Delta\chi^2 = 9.04$, $df = 3$, $p = 0.029$), such that the negative relationship between employee reports of FSSB and psychological distress for those employees in the Veteran sample was significantly stronger compared to the other three samples. Results suggest Veteran employees reporting low FSSB from their civilian supervisors have significantly stronger reports of psychological distress compared to the other three samples, lending support to the importance of supervisors' support for work–non-work lives of Veteran employees especially. Results of this analysis appear in Table 4. Figure 1 shows the relationship of these results across the association of FSSB and psychological distress for the Veteran sample compared to the other three groups. Additionally, when excluding the Veteran sample from model comparisons, there were no significant differences in the strength of the relationship between FSSB and psychological distress across the other three occupational populations samples.

Table 4. Multiple moderated regression results.

Variable	B	SE	β	R^2	ΔR^2
<i>Model 1</i>				0.081 **	
IT	−0.30 **	0.05	−0.17 **		
Healthcare	−0.15 **	0.05	−0.10 **		
SM	−0.45 **	0.05	−0.27 **		
FSSB	−0.13 **	0.01	−0.17 **		
Constant	2.11 **	0.04	2.88 **		

Table 4. Cont.

Variable	B	SE	β	R ²	ΔR^2
<i>Model 2</i>				0.083 **	0.002 *
IT	−0.70 **	0.20	−0.40 **		
Healthcare	−0.55 **	0.20	−0.37 **		
SM	−0.86 **	0.20	−0.50 **		
FSSB	−0.22 **	0.05	−0.28 **		
IT X FSSB	0.11 *	0.05	0.24 *		
Healthcare X FSSB	0.11 *	0.05	0.27 *		
SM X FSSB	0.11 *	0.05	0.26 *		
Constant	2.11 **	0.04	2.88 **		
<i>Model 3</i>				0.076 **	0.000
IT	−0.20	0.13	−0.11		
SM	−0.40 **	0.14	−0.23		
FSSB	−0.14 **	0.02	−0.17		
IT X FSSB	0.00	0.03	0.01		
SM X FSSB	0.01	0.03	0.03		
Constant	2.00 **	0.02	2.73 **		

Note: IT = Information Technology, SM = Service Member sample. FSSB was centered. For Models 1 and 2 the veteran sample was modeled as the reference category. The Healthcare group was modeled as the reference category in Model 3. * $p < 0.05$. ** $p < 0.01$. Ns range = 3752–3778.




Figure 1. Model Comparisons of FSSB and Employee Psychological Distress of the Veteran sample compared to the other three occupations (IT, Healthcare, Service Member populations), graphically revealing the negative relationship between employee reports of FSSB and psychological distress for Veteran employees compared to the other three occupations (IT, Healthcare, Service Member).

4. Discussion

This study used a common measures methodological approach to explore the relationship between employee perceptions of FSSB and their psychological distress both within and across four occupations. Our findings revealed a consistently significant, negative relationship between employee perceptions of FSSB and their psychological distress within and between samples of healthcare, IT, service member and Veteran employees. These findings provide evidence for the importance of FSSB in relation to employee psychological distress across diverse occupational roles and environments. Furthermore, mean comparison findings revealed healthcare workers’ perceptions of FSSB were lower when compared with IT, Veteran, and service member populations, consequently signaling the need to intervene in this occupation distinctly, consistent with findings on high employee burnout and psychological distress among healthcare workers during the coronavirus pandemic (COVID-19) [36,37]. Additionally, results highlight the unique needs of supervisors in civilian organizations offering support to Veteran employees, as demonstrated by the findings that Veteran employees who reported lower FSSB from their supervisors also

reported higher psychological distress compared to the other three samples. This research has important implications for future occupational health research and leader-centered workplace interventions.

Consistent with previous findings, this study reasserts the negative association of FSSB with markers of poor mental health in employees [25,38,39], while simultaneously calling for further investigation of the moderators (e.g., occupational context, population characteristics) of the psychological benefits of workplace leadership support. For example, FSSB has been previously shown to increase psychological well-being in healthcare employees [39] and diminish psychological distress in IT employees working less hours and under low-strain conditions [40]. Furthermore, FSSB was found to be negatively associated with psychological distress in an active-duty military population [41] and supervisor support has been shown to reduce negative emotions in Veterans screening positive for post-traumatic stress disorder (PTSD) [42]. High-risk military populations, such as Veterans and active-duty military personnel, typically report more adverse mental health outcomes when compared to their civilian counterparts [43] and may therefore require more leadership support that is sympathetic to their work and non-work needs in civilian contexts. Thus, this may in part explain why the relationship between FSSB and psychological distress was strongest among Veterans in this study, with the addition of the military's influence on Veterans' strong predisposition towards leadership structures [44]. However, Veterans also reported the highest rates of psychological distress compared to the other three groups—and while prior evidence has shown Veterans experience immense benefits across work, and the physical and psychosocial outcomes from having more supportive leaders in the workplace [45,46], the nature and severity of the psychological distress experienced by Veteran employees may require additional intervention [47]. For example, combat-exposed Veterans report relatively high rates of PTSD [48], a high comorbid disorder with depression [49] that is known to diminish the beneficial effects of social support on psychological distress [50].

It is also important to note the service member sample did not demonstrate comparable rates of psychological distress to Veterans even while reporting the highest FSSB levels. The lower rates of psychological distress in the service member sample may be best accounted for by differences in combat or trauma exposure, branch of service, and history of deployment [51,52] while their higher FSSB scores may be associated with their ongoing engagement in reservist activities in contrast from military separation that is experienced by Veterans.

As suggested by Straub [27], our study reveals the degree of benefit afforded by FSSB for psychological health. For example, a rigorous RCT demonstrated a larger psychological benefit of FSSB-centered training interventions for employees with increased elder and childcare responsibilities off the job [25]. Furthermore, another study demonstrated that an IT setting characterized by lower job demands and higher job control contributed to greater declines in employee psychological distress over 18 months when compared to a healthcare setting with higher job demands and lower control [40]. This suggests the differences in reported rates of FSSB and their relationship with improved psychological health may in part be impacted by occupational factors such as the structure/nature of work, as demonstrated in this study. For instance, our healthcare sample, which is predominantly female (91.8%), reported the lowest rate of FSSB ($M = 3.69$, $SD = 0.88$) in contrast with the predominantly male sample (75.6%) of military service members who reported the highest levels of FSSB ($M = 4.10$, $SD = 0.94$). In addition, the healthcare sample had the second highest scores of psychological distress, coming only second to Veterans. As a possible explanation for the health care sample's lowest reports of FSSBs—gender role ideologies may predispose women to greater household and family caretaking responsibilities [27], which in turn create a greater need for FSSB and the more sensitized perception of the lack of FSSB in a predominantly female-led work context. Furthermore, the healthcare sample reported the lowest rate of marriage or cohabitation while having the second highest average number of children, suggesting the increased likelihood of being a single

caretaker. Consequently, FSSB may be expressly critical for psychological well-being within occupational populations that are female-led and composed of single parent households, especially in workplaces characterized by high-strain (i.e., long hours and low control), such as those seen in healthcare settings [40].

4.1. Implications

A growing body of evidence derived from rigorous RCTs suggests that training supervisors in workplace support interventions (e.g., FSSB training's) provides an effective means for supporting employee psychological health and well-being outcomes [22,24,33,34,46,53,54]. This research suggests the important role that the workplace can play in improving the mental health of employee populations, and highlights the potential of FSSB training interventions for ameliorating the negative effects of psychological distress. Furthermore, this research suggests that not only do employee reports of FSSB and psychological distress differ by occupational population, but that the relationship between FSSB and psychological distress differs by occupation. This is not surprising given the extensive research on the psychosocial stressors such as demands, control, and social support at work as significantly impacting psychological health and well-being of workers [14]. Thus, our findings suggest that continued work at mitigating the impact of such psychosocial stressors is an important strategy, regardless of the occupational population.

In addition, employees may benefit from supervisors who enact FSSB as the conditions of organizations and the workforce environment may have more recent challenges brought about by COVID-19. Furthermore, there has been an increase in the number of workers working from home (WFH), with an estimated 42% of the U.S. workforce being driven to adapt to the new conditions associated with full-time WFH [55]. For example, during the peak of the pandemic, schools and daycares closed doors, childcare increased [56,57] and parents were faced with having to manage greater responsibilities within the home environment, creating conflicts between competing work and family demands [57,58]. And while WFH is related to increased control over work and positive outcomes for many, a growing percentage of parents reported deteriorating mental health for themselves and worsening behavioral health for their children [59]. Among the various known risks of WFH stated above, are those psychosocial in nature (e.g., social isolation, work-home lines blurred) [60], which can exasperate work-family demands [61], thereby requiring greater work-home boundary management support from employers [62]. On the other hand, frontline workers (e.g., healthcare employees) have also confronted their own set of work-related mental health challenges [37], while under the pressures of increased risk of infection [63] and fears of transmitting COVID-19 to family members [64]. As such, leaders in organizations across the U.S. need innovative solutions and resources for maneuvering through the urgent workplace demands presented by COVID-19 while mitigating risks to the well-being of employees and their families [62]. Providing more leadership support around such work-life challenges has been called for by the popular press and has been shown to be an effective way of reducing distress and improving well-being among workers [23,25]. It is well-established that social isolation and decreased social connectedness are related to increased psychological distress [65]. In fact, Holt-Lunstad [65] called for prioritization of social factors when addressing public health during, and in recovery from, the pandemic. Such a call is consistent with the focus of this study, which examines work as one of the missing links in improving mental health [66]. Furthermore, supervisors should be trained to look for signs of increased psychological distress during times of global crisis, and proactively destigmatize the self-disclosure of higher rates of anxiety or depression by their employees. This can in turn create a more transparent and authentic work environment where signs of reduced mental health are not viewed as a threat to employment, but as an opportunity to build trust with supervisors who can intervene by encouraging access to mental health resources.

4.2. Limitations

Limitations of this study include being focused on only four specific occupational populations and therefore, generalizing to other occupations may be limited. However, we did analyze occupations that were conceptually different from one another (e.g., desk jobs in information technology compared to high-risk occupations, such as the military) to enhance generalizability of findings. Furthermore, common method bias is a limitation given combined data were based on baseline cross-sectional samples. However, evidence of differences across populations suggests that this is not a critical limitation. Finally, we do have a methodological limitation relevant to reverse scoring items for analysis. For instance, for validation and standardization purposes across transformations, items were reversed coded for both FSSB and the K6 psychological distress scale, as such, a possible consequence contributing to the reduction in reliability. Lastly, it is important to note that the data analyzed in this study were gathered during the course of four separate studies occurring at different time points. Therefore, the differentiation in timing and research methodologies between studies should be considered when interpreting this study's results.

4.3. Future Directions

Future leadership support interventions would benefit from a greater understanding of population-dependent contextual and individual factors that encourage supervisors to engage in FSSB with respect to specific employee characteristics and needs. How does one intervention fair in one population versus another, and what accounts for its differential effects on psychological well-being? For example, Kossek [25] found an "organizational job resource-enhancing intervention" that is predicated on FSSB was more effective for diminishing perceived stress than psychological distress, and suggested introducing an intervention that more precisely targets symptoms of psychological distress relevant for individuals caring for children and elders. Similarly, Veterans with high levels of psychological distress may benefit from interventions that encourage supervisors to recognize and ask Veterans about their mental health over and above focusing on work and family. Equally so, more research is needed on the effects of supervisor interventions for supervisors themselves. For example, one study demonstrated that a supervisor intervention may be detrimental to a supervisor's work-to-family conflict and organizational commitment while simultaneously diminishing employee psychological distress [67], yet this has not been investigated extensively. As such, further research on the specific cultural and contextual factors of occupations such as healthcare, military service members, Veteran, and IT employee populations may shed light on the unique attributes that influence FSSB in their respective settings. Future research should also investigate how specific employee mental health symptoms (e.g., depression, PTSD, anxiety symptoms) differentially moderate the effectiveness of leadership support in various populations, and whether additional clinical interventions may help maximize the benefits of supervisor support in similar higher-risk employee populations.

5. Conclusions

This study provides evidence for the importance of FSSB in relation to psychological distress within and across multiple occupational populations, especially for high-risk populations such as Veteran employees working in a civilian context. Furthermore, this study presents a benchmark for the association between FSSB and psychological distress in a large, combined sample of IT, healthcare, Veteran, and service member employees ($n = 3778$). Our findings suggest the importance for training supervisors on how to enact FSSB, which may have an impact on lowering psychological distress and improving employee mental health in various occupational contexts. Most importantly, our results draw attention to the indispensable nuances of individual and occupational characteristics that may affect the relationship between supervisor support for family and employee psychological distress. This study has implications for helping shape future workplace supervisor support research and interventions and the importance of taking workplace context into consideration in

the developments of such interventions. Our study suggests that workplace efforts to train supervisors on how to be more supportive of work–non-work lives of employees with an emphasis on support for the psychological health and well-being of employees may be a critical strategy for improving mental health and well-being of workers going forward.

Author Contributions: Conceptualization, S.J.A., P.G.B. and L.B.H.; methodology, P.G.B., S.J.A. and L.B.H.; statistical analysis, S.J.A.; writing—review and editing, P.G.B., S.J.A. and L.B.H. All authors have read and agreed to the published version of the manuscript.

Funding: *MESH:* The U.S. Army Medical Research Acquisition Activity, 820 Chandler Street, Fort Detrick MD 21702-5014 is the awarding and administering acquisition office. This work was supported by Office of the Assistant Secretary of Defense for Health Affairs, through the Psychological Health and Traumatic Brain Injury Research Program–Comprehensive Universal Prevention/Health Promotion Interventions Award, under Award No. W81XWH-16-1-0720. Opinions, interpretations, conclusions, and recommendations are those of the author and are not necessarily endorsed by the Department of Defense. *SERVE:* The U.S. Army Medical Research Acquisition Activity, 820 Chandler Street, Fort Detrick, Maryland 21702-5014, is the awarding and administering acquisition office. This work was supported by the Office of the Assistant Secretary of Defense for Health Affairs, through the United States Army Medical Research and Development Command (USAMRMC) Broad Agency Announcement under Award W81XWH13-2-0020. Opinions, interpretations, conclusions, and recommendations are those of the authors and are not necessarily endorsed by the Department of Defense. *WFHN:* This research was conducted as part of the Work, Family and Health Network (www.WorkFamilyHealthNetwork.org; accessed on 12 August 2021), which is funded by a cooperative agreement through the National Institutes of Health and the Centers for Disease Control and Prevention: Eunice Kennedy Shriver National Institute of Child Health and Human Development (Grant # U01HD051217, U01HD051218, U01HD051256, U01HD051276); National Institute on Aging (Grant # U01AG027669); the National Heart, Lung, and Blood Institute (R01HL107240); Office of Behavioral and Social Sciences Research; and National Institute for Occupational Safety and Health (Grant # U01OH008788, U01HD059773). Grants from the William T. Grant Foundation, Alfred P. Sloan Foundation, and the Administration for Children and Families have provided additional funding. The contents of this publication are solely the responsibility of the authors and do not necessarily represent the official views of these institutes and offices. Special acknowledgment goes to Extramural Staff Science Collaborator, Rosalind King, and Lynne Casper, for design of the original Workplace, Family, Health and Well-Being Network Initiative. The findings and conclusions in this manuscript are those of the authors and do not necessarily represent the views of these institutes and offices. This work was also supported by the Oregon Institute of Occupational Health Sciences at Oregon Health & Science University via funds from the Division of Consumer and Business Services of the State of Oregon (ORS 656.630). Portions of this research were also supported by the Grant # T03OH008435 awarded to Portland State University, funded by the Centers for Disease Control and Prevention, National Institute for Occupational Safety and Health. Its contents are solely the responsibility of the authors and do not necessarily represent the official views of NIOSH, CDC or HHS.

Institutional Review Board Statement: Ethical review and approval were waived for this study due to IRB approval already confirmed and approved for each of the four studies at various institutions as listed in the funding section above.

Informed Consent Statement: Informed consent was obtained from all subjects involved in the study.

Data Availability Statement: Data available upon request. Permission was available for re-analysis of data across studies. Data use security protocols were signed before permission was granted.

Acknowledgments: Authors would like to acknowledge the contribution of Todd E. Bodner, professor and quantitative methodologist from Portland State University for his technical expertise, advice, and support in the methodology and statistical analysis of this study.

Conflicts of Interest: Leslie B. Hammer has a financial interest in Work Life Help, LLC., a company that may have a commercial interest in the results of this research and technology. This potential conflict of interest has been reviewed and managed by OHSU.

References

- Reinert, M.; Fritze, D.; Nguyen, T. *The State of Mental Health in America 2022*; Mental Health in America: Alexandria, VA, USA, 2021.
- Kumar, P.; Kumar, N.; Aggarwal, P.; Yeap, J.A.L. Working in Lockdown: The Relationship between COVID-19 Induced Work Stressors, Job Performance, Distress, and Life Satisfaction. *Curr. Psychol.* **2021**, *40*, 6308–6323. [[CrossRef](#)] [[PubMed](#)]
- Hamouche, S. COVID-19 and Employees' Mental Health: Stressors, Moderators and Agenda for Organizational Actions. *Emerald Open Res.* **2020**, *2*, 15. [[CrossRef](#)]
- Hammer, L.B.; Allen, E.; Grigsby, T.D. Work–Family Conflict in Dual-Earner Couples: Within-Individual and Crossover Effects of Work and Family. *J. Vocat. Behav.* **1997**, *50*, 185–203. [[CrossRef](#)]
- Hammer, L.B.; Bauer, T.N.; Grandey, A.A. Work-Family Conflict and Work-Related Withdrawal Behaviors. *J. Bus. Psychol.* **2003**, *17*, 419–436. [[CrossRef](#)]
- Hammer, L.B.; Zimmerman, K.L. Quality of Work Life. In *APA Handbook of Industrial and Organizational Psychology*; APA Handbooks in Psychology®; American Psychological Association: Washington, DC, USA, 2011; Volume 3, pp. 399–431. ISBN 1-4338-0734-3.
- Montez, J.K.; Sabbath, E.; Glymour, M.M.; Berkman, L.F. Trends in Work–Family Context among U.S. Women by Education Level, 1976 to 2011. *Policy Rev.* **2014**, *33*, 629–648. [[CrossRef](#)]
- Greenhaus, J.H.; Beutell, N.J. Sources of Conflict between Work and Family Roles. *Acad. Manag. Rev.* **1985**, *10*, 76–88. [[CrossRef](#)]
- Amstad, F.T.; Meier, L.L.; Fasel, U.; Elfering, A.; Semmer, N.K. A Meta-Analysis of Work–Family Conflict and Various Outcomes with a Special Emphasis on Cross-Domain versus Matching-Domain Relations. *J. Occup. Health Psychol.* **2011**, *16*, 151–169. [[CrossRef](#)]
- Eby, L.T.; Maher, C.P.; Butts, M.M. The Intersection of Work and Family Life: The Role of Affect. *Annu. Rev. Psychol.* **2010**, *61*, 599–622. [[CrossRef](#)]
- Hammer, L.B.; Neal, M.B. Working Sandwiched-Generation Caregivers: Prevalence, Characteristics, and Outcomes. *Psychol. Manag. J.* **2008**, *11*, 93–112. [[CrossRef](#)]
- Ozer, E.M. The Impact of Childcare Responsibility and Self-Efficacy on the Psychological Health of Professional Working Mothers. *Psychol. Women Q.* **1995**, *19*, 315–335. [[CrossRef](#)]
- Ganster, D.C.; Rosen, C.C. Work Stress and Employee Health: A Multidisciplinary Review. *J. Manag.* **2013**, *39*, 1085–1122. [[CrossRef](#)]
- Karasek, R.A. Job Demands, Job Decision Latitude, and Mental Strain: Implications for Job Redesign. *Adm. Sci. Q.* **1979**, *24*, 285–308. [[CrossRef](#)]
- Bavik, Y.L.; Shaw, J.D.; Wang, X.-H. (Frank) Social Support: Multidisciplinary Review, Synthesis, and Future Agenda. *Acad. Manag. Ann.* **2020**, *14*, 726–758. [[CrossRef](#)]
- Chang, A.; Chen, S.-C.; Chi, S.-C.S. Role Salience and Support as Moderators of Demand/Conflict Relationships in China. *Eur. J. Work Organ. Psychol.* **2014**, *23*, 859–874. [[CrossRef](#)]
- Ferguson, M.; Carlson, D.; Kacmar, K.M.; Halbesleben, J.R.B. The Supportive Spouse at Work: Does Being Work-Linked Help? *J. Occup. Health Psychol.* **2016**, *21*, 37–50. [[CrossRef](#)]
- Griggs, T.L.; Casper, W.J.; Eby, L.T. Work, Family and Community Support as Predictors of Work–Family Conflict: A Study of Low-Income Workers. *J. Vocat. Behav.* **2013**, *82*, 59–68. [[CrossRef](#)]
- Hammer, L.B.; Kossek, E.E.; Yragui, N.L.; Bodner, T.E.; Hanson, G.C. Development and Validation of a Multidimensional Measure of Family Supportive Supervisor Behaviors (FSSB). *J. Manag.* **2009**, *35*, 837–856. [[CrossRef](#)]
- Kossek, E.E.; Pichler, S.; Bodner, T.; Hammer, L.B. Workplace Social Support and Work–Family Conflict: A Meta-Analysis Clarifying the Influence of General and Work–Family-Specific Supervisor and Organizational Support. *Pers. Psychol.* **2011**, *64*, 289–313. [[CrossRef](#)]
- Hammer, L.B.; Ernst Kossek, E.; Bodner, T.; Crain, T. Measurement Development and Validation of the Family Supportive Supervisor Behavior Short-Form (FSSB-SF). *J. Occup. Health Psychol.* **2013**, *18*, 285–296. [[CrossRef](#)]
- Hammer, L.B.; Brady, J.M.; Brossoit, R.M.; Mohr, C.D.; Bodner, T.E.; Crain, T.L.; Brockwood, K.J. Effects of a Total Worker Health® Leadership Intervention on Employee Well-Being and Functional Impairment. *J. Occup. Health Psychol.* **2021**, *26*, 582–598. [[CrossRef](#)]
- Hammer, L.B.; Perry, M.L. Reducing Work–Life Stress: The Place for Integrated Interventions. In *Total Worker Health*; American Psychological Association: Washington, DC, USA, 2019; pp. 263–278. ISBN 1433830256.
- Brady, J.M.; Hammer, L.B.; Mohr, C.D.; Bodner, T.E. Supportive Supervisor Training Improves Family Relationships among Employee and Spouse Dyads. *J. Occup. Health Psychol.* **2021**, *26*, 31–48. [[CrossRef](#)] [[PubMed](#)]
- Kossek, E.E.; Thompson, R.J.; Lawson, K.M.; Bodner, T.; Perrigino, M.B.; Hammer, L.B.; Buxton, O.M.; Almeida, D.M.; Moen, P.; Hurtado, D.A.; et al. Caring for the Elderly at Work and Home: Can a Randomized Organizational Intervention Improve Psychological Health? *J. Occup. Health Psychol.* **2019**, *24*, 36–54. [[CrossRef](#)]
- Hammer, L.B. The Interplay of Workplace Redesign and Public Policy in the 21st Century. *Am. J. Public Health* **2021**, *111*, 1784–1786. [[CrossRef](#)] [[PubMed](#)]
- Straub, C. Antecedents and Organizational Consequences of Family Supportive Supervisor Behavior: A Multilevel Conceptual Framework for Research. *Hum. Resour. Manag. Rev.* **2012**, *22*, 15–26. [[CrossRef](#)]

28. Hanson, G.C.; Rameshbabu, A.; Bodner, T.E.; Hammer, L.B.; Rohlman, D.S.; Olson, R.; Wipfli, B.; Kuehl, K.; Perrin, N.A.; Alley, L.; et al. A Comparison of Safety, Health, and Well-Being Risk Factors across Five Occupational Samples. *Front. Public Health* **2021**, *9*, 614725. [[CrossRef](#)]
29. Cohen, S.; Wills, T.A. Stress, Social Support, and the Buffering Hypothesis. *Psychol. Bull.* **1985**, *98*, 310–357. [[CrossRef](#)]
30. Holt-Lunstad, J.; Smith, T.B.; Layton, J.B. Social Relationships and Mortality Risk: A Meta-Analytic Review. *PLoS Med.* **2010**, *7*, e1000316. [[CrossRef](#)]
31. Joiner, T.E., Jr.; Van Orden, K.A.; Witte, T.K.; Rudd, M.D. *The Interpersonal Theory of Suicide: Guidance for Working with Suicidal Clients*; The interpersonal theory of suicide: Guidance for working with suicidal clients; American Psychological Association: Washington, DC, USA, 2009; p. 246. ISBN 978-1-4338-0426-7.
32. Duffany, K.O.; Finegood, D.T.; Matthews, D.; McKee, M.; Venkat Narayan, K.M.; Puska, P.; Siegel, K.; Stevens, D.; Wong, F.; Woodward, M.; et al. Community Interventions for Health (CIH): A Novel Approach to Tackling the Worldwide Epidemic of Chronic Diseases. *CVD Prev. Control.* **2011**, *6*, 47–56. [[CrossRef](#)]
33. Hammer, L.B.; Wan, W.H.; Brockwood, K.J.; Mohr, C.D.; Carlson, K.F. Military, Work, and Health Characteristics of Separated and Active Service Members from the Study for Employment Retention of Veterans (SERVe). *Mil. Psychol.* **2017**, *29*, 491–512. [[CrossRef](#)]
34. Bray, J.W.; Kelly, E.L.; Hammer, L.B.; Almeida, D.M.; Dearing, J.W.; King, R.B.; Buxton, O.M. An Integrative, Multilevel, and Transdisciplinary Research Approach to Challenges of Work, Family, and Health. In *Methods Report*; RTI Press: Research Triangle Park, NC, USA, 2013; pp. 1–38. [[CrossRef](#)]
35. Kessler, R.C.; Andrews, G.; Colpe, L.J.; Hiripi, E.; Mroczek, D.K.; Normand, S.-L.T.; Walters, E.E.; Zaslavsky, A.M. Short Screening Scales to Monitor Population Prevalences and Trends in Non-Specific Psychological Distress. *Psychol. Med.* **2002**, *32*, 959–976. [[CrossRef](#)]
36. Norman, S.B.; Feingold, J.H.; Kaye-Kauderer, H.; Kaplan, C.A.; Hurtado, A.; Kachadourian, L.; Feder, A.; Murrrough, J.W.; Charney, D.; Southwick, S.M.; et al. Moral Distress in Frontline Healthcare Workers in the Initial Epicenter of the COVID-19 Pandemic in the United States: Relationship to PTSD Symptoms, Burnout, and Psychosocial Functioning. *Depress. Anxiety* **2021**, *38*, 1007–1017. [[CrossRef](#)] [[PubMed](#)]
37. Pearman, A.; Hughes, M.L.; Smith, E.L.; Neupert, S.D. Mental Health Challenges of United States Healthcare Professionals during COVID-19. *Front. Psychol.* **2020**, *11*, 2065. [[CrossRef](#)] [[PubMed](#)]
38. Crain, T.L.; Stevens, S. Family-Supportive Supervisor Behaviors: A Review and Recommendations for Research and Practice. *J. Organ. Behav.* **2018**, *39*, 869–888. [[CrossRef](#)]
39. Şahin, S.; Adegbite, W.M.; Tiryaki Şen, H. How Do Family Supportive Supervisors Affect Nurses' Thriving: A Research before and during COVID-19 Pandemic. *Arch. Psychiatr. Nurs.* **2021**, *35*, 602–609. [[CrossRef](#)]
40. Fan, W.; Moen, P.; Kelly, E.L.; Hammer, L.B.; Berkman, L.F. Job Strain, Time Strain, and Well-Being: A Longitudinal, Person-Centered Approach in Two Industries. *J. Vocat. Behav.* **2019**, *110*, 102–116. [[CrossRef](#)] [[PubMed](#)]
41. Huffman, A.H.; Olson, K.J. Gender Differences in Perceptions of Resources and Turnover Intentions of Work-Linked Couples in Masculine Occupations. *Stress Health* **2017**, *33*, 309–321. [[CrossRef](#)] [[PubMed](#)]
42. Mohr, C.D.; Hammer, L.B.; Brady, J.M.; Perry, M.L.; Bodner, T. Can Supervisor Support Improve Daily Employee Well-Being? Evidence of Supervisor Training Effectiveness in a Study of Veteran Employee Emotions. *J. Occup. Organ. Psychol.* **2021**, *94*, 400–426. [[CrossRef](#)]
43. Hoglund, M.W.; Schwartz, R.M. Mental Health in Deployed and Nondeployed Veteran Men and Women in Comparison with Their Civilian Counterparts. *Mil. Med.* **2014**, *179*, 19–25. [[CrossRef](#)]
44. Wong, L.; Bliese, P.; McGurk, D. Military Leadership: A Context Specific Review. *Leadersh. Q.* **2003**, *14*, 657–692. [[CrossRef](#)]
45. Hammer, L.B.; Brockwood, K.; Haverly, S.N. The Key Role of Supervisors for Supporting Veterans in the Workplace. In *Military Veteran Employment*; Oxford University Press: New York, NY, USA, 2021; ISBN 978-0-19-064298-3.
46. Hammer, L.B.; Wan, W.H.; Brockwood, K.J.; Bodner, T.; Mohr, C.D. Supervisor Support Training Effects on Veteran Health and Work Outcomes in the Civilian Workplace. *J. Appl. Psychol.* **2019**, *104*, 52–69. [[CrossRef](#)]
47. Smith, R.T.; True, G. Warring Identities: Identity Conflict and the Mental Distress of American Veterans of the Wars in Iraq and Afghanistan. *Soc. Ment. Health* **2014**, *4*, 147–161. [[CrossRef](#)]
48. Hoge, C.W.; Warner, C.H. Estimating Ptsd Prevalence in Us Veterans: Considering Combat Exposure, Ptsd Checklist Cutpoints, and DSM-5. *J. Clin. Psychiatry* **2014**, *75*, 6657. [[CrossRef](#)] [[PubMed](#)]
49. Chan, D.; Cheadle, A.D.; Reiber, G.; Unützer, J.; Chaney, E.F. Health Care Utilization and Its Costs for Depressed Veterans with and without Comorbid PTSD Symptoms. *Psychiatr. Serv.* **2009**, *60*, 1612–1617. [[CrossRef](#)] [[PubMed](#)]
50. Brancu, M.; Thompson, N.L.; Beckham, J.C.; Green, K.T.; Calhoun, P.S.; Elbogen, E.B.; Robbins, A.T.; Fairbank, J.A.; Wagner, H.R. The Impact of Social Support on Psychological Distress for U.S. Afghanistan/Iraq Era Veterans with PTSD and Other Psychiatric Diagnoses. *Psychiatry Res.* **2014**, *217*, 86–92. [[CrossRef](#)]
51. Baker, D.G.; Heppner, P.; Afari, N.; Nunnink, S.; Kilmer, M.; Simmons, A.; Harder, L.; Bosse, B. Trauma Exposure, Branch of Service, and Physical Injury in Relation to Mental Health among U.S. Veterans Returning from Iraq and Afghanistan. *Mil. Med.* **2009**, *174*, 6. [[CrossRef](#)]
52. Perconte, T.; Wilson, A.T. Psychological and War Stress Symptoms among Deployed and Non-Deployed Reservists Following the Persian Gulf War. *Mil. Med.* **1993**, *158*, 6. [[CrossRef](#)]

53. Hammer, L.B.; Brady, J.M.; Perry, M.L. Training Supervisors to Support Veterans at Work: Effects on Supervisor Attitudes and Employee Sleep and Stress. *J. Occup. Organ. Psychol.* **2020**, *93*, 273–301. [[CrossRef](#)]
54. Hammer, L.B.; Truxillo, D.M.; Bodner, T.; Rineer, J.; Pytlovany, A.C.; Richman, A. Effects of a Workplace Intervention Targeting Psychosocial Risk Factors on Safety and Health Outcomes. *Biomed. Res. Int.* **2015**, *2015*, 836967. [[CrossRef](#)]
55. Bloom, N. *How Working from Home Works Out*; Stanford Institute for Economic Policy Research: Stanford, CA, USA, 2020; pp. 1–8.
56. Fisher, J.; Languilaire, J.-C.; Lawthom, R.; Nieuwenhuis, R.; Petts, R.J.; Runswick-Cole, K.; Yerkes, M.A. Community, Work, and Family in Times of COVID-19. *Community Work. Fam.* **2020**, *23*, 247–252. [[CrossRef](#)]
57. Zamarro, G.; Prados, M.J. Gender Differences in Couples' Division of Childcare, Work and Mental Health during COVID-19. *Rev. Econ. Househ.* **2021**, *19*, 11–40. [[CrossRef](#)]
58. Power, K. The COVID-19 Pandemic Has Increased the Care Burden of Women and Families. *Sustain. Sci. Pract. Policy* **2020**, *16*, 67–73. [[CrossRef](#)]
59. Patrick, S.W.; Henkhaus, L.E.; Zickafoose, J.S.; Lovell, K.; Halvorson, A.; Loch, S.; Letterie, M.; Davis, M.M. Well-Being of Parents and Children during the COVID-19 Pandemic: A National Survey. *Pediatrics* **2020**, *146*, e2020016824. [[CrossRef](#)] [[PubMed](#)]
60. Bouziri, H.; Smith, D.R.M.; Descatha, A.; Dab, W.; Jean, K. Working from Home in the Time of COVID-19: How to Best Preserve Occupational Health? *Occup. Env. Med.* **2020**, *77*, 509–510. [[CrossRef](#)] [[PubMed](#)]
61. Oakman, J.; Kinsman, N.; Stuckey, R.; Graham, M.; Weale, V. A Rapid Review of Mental and Physical Health Effects of Working at Home: How Do We Optimise Health? *BMC Public Health* **2020**, *20*, 1825. [[CrossRef](#)] [[PubMed](#)]
62. Kniffin, K.M.; Narayanan, J.; Anseel, F.; Antonakis, J.; Ashford, S.P.; Bakker, A.B.; Bamberger, P.; Bapuji, H.; Bhawe, D.P.; Choi, V.K.; et al. COVID-19 and the Workplace: Implications, Issues, and Insights for Future Research and Action. *Am. Psychol.* **2021**, *76*, 63–77. [[CrossRef](#)]
63. Do, D.P.; Frank, R.U.S. Frontline Workers and COVID-19 Inequities. *Prev. Med.* **2021**, *153*, 106833. [[CrossRef](#)]
64. Shechter, A.; Diaz, F.; Moise, N.; Anstey, D.E.; Ye, S.; Agarwal, S.; Birk, J.L.; Brodie, D.; Cannone, D.E.; Chang, B.; et al. Psychological Distress, Coping Behaviors, and Preferences for Support among New York Healthcare Workers during the COVID-19 Pandemic. *Gen. Hosp. Psychiatry* **2020**, *66*, 1–8. [[CrossRef](#)]
65. Holt-Lunstad, J. Social Connection as a Public Health Issue: The Evidence and a Systemic Framework for Prioritizing the “Social” in Social Determinants of Health. *Annu. Rev. Public Health* **2022**, *43*, 193–213. [[CrossRef](#)]
66. Dimoff, J.K.; Kelloway, E.K. With a Little Help from My Boss: The Impact of Workplace Mental Health Training on Leader Behaviors and Employee Resource Utilization. *J. Occup. Health Psychol.* **2019**, *24*, 4–19. [[CrossRef](#)]
67. Perry, M.L.; El-Askari, L.M.; Hammer, L.B.; Brown, N.D. Securing Your Own Mask before Assisting Others: Effects of a Supervisor Training Intervention on Supervisors and Employees. *Occup. Health Sci.* **2020**, *4*, 417–443. [[CrossRef](#)]