

Portland State University

PDXScholar

---

Mathematics and Statistics Faculty  
Publications and Presentations

Fariborz Maseeh Department of Mathematics  
and Statistics

---

3-17-2022

# Dependence Comparisons of Order Statistics in the Proportional Hazards Model

Subhash C. Kochar

Portland State University, [kochar@pdx.edu](mailto:kochar@pdx.edu)

Follow this and additional works at: [https://pdxscholar.library.pdx.edu/mth\\_fac](https://pdxscholar.library.pdx.edu/mth_fac)



Part of the [Physical Sciences and Mathematics Commons](#)

Let us know how access to this document benefits you.

---

## Citation Details

Kochar, Subhash C., (2022). [PRE-PRINT] "Dependence Comparisons of Order Statistics in the Proportional Hazards Model," To be published in *Probability in Engineering and Information Sciences*.

This Pre-Print is brought to you for free and open access. It has been accepted for inclusion in Mathematics and Statistics Faculty Publications and Presentations by an authorized administrator of PDXScholar. Please contact us if we can make this document more accessible: [pdxscholar@pdx.edu](mailto:pdxscholar@pdx.edu).

# Dependence comparisons of order statistics in the proportional hazards model

Subhash Kochar

Fariborz Maseeh Department of Mathematics and Statistics

Portland State University, Portland, OR, USA

Email:kochar@pdx.edu

March 17, 2022

## Abstract

Let  $X_1, \dots, X_n$  be mutually independent exponential random variables with distinct hazard rates  $\lambda_1, \dots, \lambda_n$  and let  $Y_1, \dots, Y_n$  be a random sample from the exponential distribution with hazard rate  $\bar{\lambda} = \sum_{i=1}^n \lambda_i/n$ . Also let  $X_{1:n} < \dots < X_{n:n}$  and  $Y_{1:n} < \dots < Y_{n:n}$  be their associated order statistics. It is proved that for  $1 \leq i < j \leq n$ , the generalized spacing  $X_{j:n} - X_{i:n}$  is more dispersed than  $Y_{j:n} - Y_{i:n}$  according to dispersive ordering and for  $2 \leq i \leq n$ , the dependence of  $X_{i:n}$  on  $X_{1:n}$  is less than that of  $Y_{i:n}$  on  $Y_{1:n}$ , in the sense of the more stochastically increasing ordering. This dependence result is also extended to the PHR model. This extends the earlier work of *Genest, Kochar and Xu* [*Journal of Multivariate Analysis* **100** (2009), 1587-1592] who proved this result for  $i = n$ .

**Key Words** Dispersive ordering; Exponential distribution; Kendall's tau; Stochastically increasing; Monotone regression dependence; Concordance ordering; generalized spacing.

# 1 Introduction

Several notions of monotone dependence have been introduced and studied in the literature. Researchers have also developed the corresponding dependence (partial) orderings which compare the degree of (monotone) dependence within the components of different random vectors of the same length. For details, see the pioneering paper of *Lehmann* (1966) and Chapter 5 of *Barlow and Proschan* (1981) for different notions of positive dependence, and that of *Kimeldorf and Sampson* (1989) for a unified treatment of families, orderings and measures of monotone dependence. For more detailed discussion of these concepts see Chapter 2 of *Joe* (1997) and Chapter 5 of *Nelsen* (1999).

Let  $\{X_1, \dots, X_n\}$  be a set of continuous random variables. Many authors have investigated the nature of the dependence that may exist between two order statistics  $X_{i:n}$  and  $X_{j:n}$  for  $1 \leq i < j \leq n$  under different distributional scenarios. It seems natural to expect some degree of positive dependence between them. It is well known that the order statistics based on independent (but not necessarily identically distributed) random variables are associated (cf, *Barlow and Proschan*, 1981, p.32). This yields many useful product inequalities for order statistics of independent random variables, and in particular,  $Cov(X_{i:n}, X_{j:n}) \geq 0$  for all  $i$  and  $j$ , which was initially proved by *Bickel* (1967) when the  $X$ 's are independent and identically distributed (i.i.d.).

*Boland et al.* (1996) studied the problem of dependence among order statistics in detail and discussed different types of dependence that hold between them. It is shown in that paper that in the case of i.i.d. observations, any pair of order statistics is  $TP_2$  dependent (also called monotone likelihood ratio dependent) which is the strongest type of dependence in the hierarchy of various dependence criteria as described in *Barlow and Proschan* (1981). But this is not the case in the non-iid case. However, as is shown in *Boland et al.* (1996) that, in general, whereas  $X_{j:n}$  may not be stochastically increasing in  $X_{i:n}$  for  $i < j$ ,  $X_{j:n}$  is always right tail increasing

(RTI) in  $X_{i:n}$ . The reader is referred to Chapter 5 of *Barlow and Proschan* (1981) for basic definitions and relations among various types of dependence.

It is also of interest to compare the strength of dependence that may exist between two pairs of order statistics. When the parent distribution from which the random sample is drawn has an increasing hazard rate and a decreasing reverse hazard rate, *Tukey* (1958) showed that

$$\text{Cov}(X_{i':n}, X_{j':n}) \leq \text{Cov}(X_{i:n}, X_{j:n}) \quad (1.1)$$

for either  $i = i'$  and  $j \leq j'$ ; or  $j = j'$  and  $i' \leq i$ . *Kim and David* (1990) proved that if both the hazard and the reverse hazard rates of the  $X_i$ 's are increasing, then inequality (1.1) remains valid when  $i = i'$  and  $j \leq j'$ ; However, the inequality (1.1) is reversed when  $j = j'$  and  $i' \leq i$ .

Let  $X_1, \dots, X_n$  be mutually independent exponentials with distinct hazard rates  $\lambda_1, \dots, \lambda_n$  and let  $Y_1, \dots, Y_n$  form a random sample from the exponential distribution with hazard rate  $\bar{\lambda} = (\sum_{i=1}^n X_i)/n$ . *Sathe* (1988) proved that for any  $j \in \{2, \dots, n\}$

$$\text{corr}(X_{1:n}, X_{j:n}) \leq \text{corr}(Y_{1:n}, Y_{j:n}). \quad (1.2)$$

Although this observation is interesting, it merely compares the relative degree of *linear* association within the two pairs. It is now widely recognized, however, that margin-free measures of association are more appropriate than Pearson's correlation, because they are based on the unique underlying copula which governs the dependence between the components of a continuous random pair.

In this paper a long standing open problem that for  $1 \leq i < j \leq n$ , the generalized spacing  $X_{j:n} - X_{i:n}$  is more dispersed than  $Y_{j:n} - Y_{i:n}$  according to dispersive ordering, has been solved (cf. *Xu and Balakrishnan*, 2012). This result is used to solve a related open problem that for  $2 \leq i \leq n$ , the dependence of  $X_{i:n}$  on  $X_{1:n}$  is less than that of  $Y_{i:n}$  on  $Y_{1:n}$ , in the sense of the more stochastically increasing ordering. This dependence result is also extended to the PHR model. This extends the earlier work of *Genest, Kochar and Xu* (2009) who proved this result for  $i = n$ . Section 2 is

on some preliminaries where various definitions and notations are given. The main results of this paper are given in Chapter 3.

## 2 Preliminaries

For  $i = 1, 2$ , let  $(X_i, Y_i)$  be a pair of continuous random variables with joint cumulative function  $H_i$  and margins  $F_i, G_i$ . Let

$$C_i(u, v) = H_i\{F_i^{-1}(u), G_i^{-1}(v)\}, \quad u, v \in (0, 1)$$

be the unique copula associated with  $H_i$ . In other words,  $C_i$  is the distribution of the pair  $(U_i, V_i) \equiv (F_i(X_i), G_i(Y_i))$  whose margins are uniform on the interval  $(0, 1)$ . See, e.g., Chapter 1 of *Nelsen* (1999) for details.

The most well understood partial order to compare the strength of dependence within two random vectors is that of *positive quadrant dependence* (PQD) as defined below.

**Definition 2.1** *A copula  $C_1$  is said to be less dependent than copula  $C_2$  in the positive quadrant dependence ordering (PQD), denoted  $(X_1, Y_1) \prec_{\text{PQD}} (X_2, Y_2)$ , if and only if*

$$C_1(u, v) \leq C_2(u, v), \quad u, v \in (0, 1).$$

This condition implies that  $\kappa(S_1, T_1) \leq \kappa(S_2, T_2)$  for all concordance measures meeting the axioms of *Scarsini* (1984) like Kendall's  $\tau$  and Spearman's  $\rho$ . See *Tchen* (1980) for details.

*Lehmann* (1966) in his seminal work introduced the notion of *monotone regression dependence* (MRD) which is also known as *stochastic increasingness* (SI) in the literature.

**Definition 2.2** *Let  $(X, Y)$  be a bivariate random vector with joint distribution function  $H$ . The variable  $Y$  is said to be stochastically increasing (SI) in  $X$  if for all  $(x_1, x_2) \in \mathbb{R}^2$ ,*

$$x_1 < x_2 \Rightarrow P(Y \leq y | X = x_2) \leq P(Y \leq y | X = x_1), \quad \text{for all } y \in \mathbb{R}, \quad (2.1)$$

If we denote by  $H_x$  the distribution function of the conditional distribution of  $Y$  given  $X = x$ , then (2.1) can be rewritten as

$$x_1 < x_2 \Rightarrow H_{x_2} \circ H_{x_1}^{-1}(u) \leq u, \quad \text{for } 0 \leq u \leq 1. \quad (2.2)$$

Note that in case  $X$  and  $Y$  are independent,  $H_{x_2} \circ H_{x_1}^{-1}(u) = u$ , for  $0 \leq u \leq 1$  and for all  $(x_1, x_2)$ . The SI property is not symmetric in  $X$  and  $Y$ . It is a very strong notion of positive dependence and many of the other notions of positive dependence follow from it.

Denoting by  $\xi_p = F^{-1}(p)$ , the  $p$ th quantile of the marginal distribution of  $X$ , we see that (2.2) will hold if and only if for all  $0 \leq u \leq 1$ ,

$$0 \leq p < q \leq 1 \Rightarrow H_{\xi_q} \circ H_{\xi_p}^{-1}(u) \leq u.$$

In his book, *Joe* (1997) mentions a number of bivariate stochastic ordering relations. One such notion is that of greater monotone regression (or more SI) dependence, originally considered by *Yanagimoto and Okamoto* (1969) and later extended and further investigated by *Schriever* (1987), *Capéraà and Genest* (1990), and *Avérous, Genest and Kocher* (2005), among others.

As discussed in the books by *Joe* (1997) and *Nelsen* (1999) a reasonable way to compare the relative degree of dependence between two random vectors is through their copulas. We will discuss here the concept of more SI, a partial order to compare the strength of dependence that may exist between two bivariate random vectors in the sense of monotone regression dependence (stochastic increasingness).

Suppose we have two pairs of continuous random variables  $(X_1, Y_1)$  and  $(X_2, Y_2)$  with joint cumulative distribution functions  $H_i$  and marginals  $F_i$  and  $G_i$  for  $i = 1, 2$ .

**Definition 2.3**  $Y_2$  is said to be more stochastically increasing in  $X_2$  than  $Y_1$  is in  $X_1$ , denoted by  $(Y_1|X_1) \prec_{\text{SI}} (Y_2|X_2)$  or  $H_1 \prec_{\text{SI}} H_2$ , if

$$0 < p \leq q < 1 \implies H_{2, \xi_{2q}} \circ H_{2, \xi_{2p}}^{-1}(u) \leq H_{1, \xi_{1q}} \circ H_{1, \xi_{1p}}^{-1}(u), \quad (2.3)$$

for all  $u \in (0, 1)$ , where for  $i = 1, 2$ ,  $H_{i,s}$  denotes the conditional distribution of  $Y_i$  given  $X_i = s$ , and  $\xi_{ip} = F_i^{-1}(p)$  stands for the  $p$ th quantile of the marginal distribution of  $X_i$ .

Obviously, (2.3) implies that  $Y_2$  is stochastically increasing in  $X_2$  if  $X_1$  and  $Y_1$  are independent. It also implies that if  $Y_1$  is stochastically increasing in  $X_1$ , then so is  $Y_2$  in  $X_2$ ; and conversely, if  $Y_2$  is stochastically decreasing in  $X_2$ , then so is  $Y_1$  in  $X_1$ . The above definition of more SI is copula based and

$$H_1 \prec_{\text{SI}} H_2 \Rightarrow H_1 \prec_{\text{PQD}} H_2.$$

*Avérous, Genest and Kochar* (2005) have shown in their paper that in the case of i.i.d. observations, the copula of any pair of order statistics is independent of the distribution of the parent observations as long as they are continuous. In a sense, their copula has a distribution-free property. But this is not the case if the observations are not i.i.d.

The natural question is to see if we can extend the result given in (1.2) to a copula based positive dependence ordering. *Genest, Kochar and Xu* (2009) proved that under the given conditions,

$$(X_{n:n}|X_{1:n}) \prec_{\text{SI}} (Y_{n:n}|Y_{1:n}).$$

It has been an open problem to see whether this result can be generalized from the the largest order statistics to other order statistics. We prove in this paper that for  $2 \leq i \leq n$ ,

$$(X_{i:n}|X_{1:n}) \prec_{\text{SI}} (Y_{i:n}|Y_{1:n}).$$

This implies in particular that

$$\kappa(X_{i:n}, X_{1:n}) \leq \kappa(Y_{i:n}, Y_{1:n}),$$

where  $\kappa(S, T)$  represents any concordance measure between random variables  $S$  and  $T$  in the sense of *Scarsini* (1984), e.g., Spearman's rho or Kendall's tau. A related work to this problem is *Dolati, Genest and Kochar* (2008).

### 3 Main results

The proof of our main result relies heavily on the notion of dispersive ordering between two random variables  $X$  and  $Y$ , and properties thereof. For completeness, the definition of this concept is recalled below.

**Definition 3.1** *A random variable  $X$  with distribution function  $F$  is said to be less dispersed than another variable  $Y$  with distribution  $G$ , written as  $X \leq_{disp} Y$  or  $F \leq_{disp} G$ , if and only if*

$$F^{-1}(v) - F^{-1}(u) \leq G^{-1}(u) - G^{-1}(v)$$

for all  $0 < u \leq v < 1$ .

The dispersive order is closely related to the star-ordering which is a partial order to compare the relative aging or skewness and is defined as below.

**Definition 3.2** *A random variable  $X$  with distribution function  $F$  is said to be star ordered with respect to another random variable  $Y$  with distribution  $G$ , written as  $X \leq_* Y$  or  $F \leq_* G$ , if and only if*

$$\frac{G^{-1} \circ F(x)}{x} \text{ is increasing in } x.$$

For nonnegative random variables, the star order is related to the dispersive order by

$$X \leq_* Y \Leftrightarrow \log X \leq_{disp} \log Y$$

The proof of the next lemma, which is a refined version of a result by *Deshpande and Kochar* (1983), on relation between the star-order and the dispersive order, can be found in *Kochar and Xu* (2012).

**Lemma 3.1** *Let  $X$  and  $Y$  be two random variables. If  $X \leq_* Y$  and  $X \leq_{st} Y$ , then  $X \leq_{disp} Y$ .*

The following result of *Khaledi and Kochar* (2005) will be used to prove our main theorem.

**Lemma 3.2** (*Khaledi and Kochar, 2005*) *Let  $X_i$  and  $Y_i$  be independent random variables with distribution functions  $F_i$  and  $G_i$ , respectively for  $i = 1, 2$ . Then*

$$X_2 \leq_{disp} X_1 \text{ and } Y_1 \leq_{disp} Y_2 \Rightarrow (X_2 + Y_2)|X_2 \prec_{SI} (X_1 + Y_1)|X_1$$

The next theorem on dispersive ordering between general spacings which is also of independent interest, will be used to prove our main result.

**Theorem 3.1** *Let  $X_1, \dots, X_n$  be mutually independent exponential random variables with distinct hazard rates  $\lambda_1, \dots, \lambda_n$  and let  $Y_1, \dots, Y_n$  be a random sample from the exponential distribution with hazard rate  $\bar{\lambda}$ . Then for  $1 \leq i < j \leq n$ ,*

$$(Y_{j:n} - Y_{i:n}) \leq_{disp} (X_{j:n} - X_{i:n}) \quad (3.1)$$

**Proof.** *Yu* (2021) proved in Corollary 1 of his paper that for  $1 \leq i < j \leq n$ ,

$$(Y_{j:n} - Y_{i:n}) \leq_* (X_{j:n} - X_{i:n}) \quad (3.2)$$

and *Kochar and Rojo* (1996) proved in their Corollary 2.1 that

$$(Y_{j:n} - Y_{i:n}) \leq_{st} (X_{j:n} - X_{i:n}) \quad (3.3)$$

Using (3.2) and (3.3), we get (3.1) using Lemma 3.1. ■

*Xu and Balakrishnan* (2012) proved a special case of the above result when  $i = n$ .

Now we give the main result of this paper.

**Theorem 3.2** *Let  $X_1, \dots, X_n$  be mutually independent exponentials with distinct hazard rates  $\lambda_1, \dots, \lambda_n$  and let  $Y_1, \dots, Y_n$  form a random sample from the exponential distribution with hazard rate  $\bar{\lambda} = (\sum_{i=1}^n \lambda_i)/n$ . Then for  $i \in \{2, \dots, n\}$ ,*

$$(X_{i:n}|X_{1:n}) \prec_{SI} (Y_{i:n}|Y_{1:n}).$$

**Proof.** It follows from Theorem 3.1 above that for  $2 \leq i \leq n$ ,

$$(Y_{i:n} - Y_{1:n}) \leq_{disp} (X_{i:n} - X_{1:n}). \quad (3.4)$$

*Kochar and Korwar (1996)* proved that  $(X_{i:n} - X_{1:n})$  is independent of  $X_{1:n}$  and  $X_{1:n} \stackrel{st}{=} Y_{1:n}$ . Similarly,  $(Y_{i:n} - Y_{1:n})$  is independent of  $Y_{1:n}$ . Now we can express  $X_{i:n}$  and  $Y_{i:n}$  as

$$X_{i:n} = (X_{i:n} - X_{1:n}) + X_{1:n} \text{ and } Y_{i:n} = (Y_{i:n} - Y_{1:n}) + Y_{1:n}$$

Using (3.4) and the above two equations, it follows from Lemma 3.2 that

$$(X_{i:n}|X_{1:n}) \prec_{SI} (Y_{i:n}|Y_{1:n}), \text{ for } 2 \leq i \leq n.$$

This proves the required result. ■

**Remark 3.1** . Since the copula of the order statistics of a random sample has the distribution-free property, it is not required that the common hazard rate of  $Y$ 's is necessarily  $\bar{\lambda}$  in the above theorems. In fact, the  $Y$ 's could be a random sample from *any* continuous distribution.

The above theorem can be extended to the PHR model using the technique used in *Genest, Kochar and Xu (2009)*.

**Theorem 3.3** Let  $X_1, \dots, X_n$  be independent continuous random variables following the PHR model with  $\lambda_1, \dots, \lambda_n$  as the proportionality parameters. Let  $Y_1, \dots, Y_n$  be i.i.d. continuous random variables with common survival function  $\bar{F}^{\bar{\lambda}}$ , then

$$(X_{i:n}|X_{1:n}) \prec_{SI} (Y_{i:n}|Y_{1:n}) \text{ for } 2 \leq i \leq n. \quad (3.5)$$

**Proof.** Let  $R(t) = -\log\{\bar{F}(t)\}$  in the PHR model. Make the monotone transformations  $X_i^* = R(X_i)$  and  $Y_i^* = -\log\{\bar{F}(Y_i)\}$ . Then the transformed variable  $X_i^*$  has exponential distribution with hazard rate  $\lambda_i$ ,  $i = 1, \dots, n$  and  $Y_1^*, \dots, Y_n^*$  is a random

sample from an exponential distribution with parameter  $\bar{\lambda}$ . Let  $X_{(1)}^* < \dots < X_{(n)}^*$  and  $Y_{(1)}^* < \dots < Y_{(n)}^*$  be the order statistics corresponding to the new sets of variables.

In view of their invariance by monotone increasing transformations of the margins, the copulas associated with the pairs  $(X_{1:n}, X_{n:n})$  and  $(X_{1:n}^*, X_{n:n}^*)$  coincide. Similarly, the pairs  $(Y_{1:n}, Y_{n:n})$  and  $(Y_{1:n}^*, Y_{n:n}^*)$  have the same copula.

Also, since the more SI dependence ordering is copula-based,

$$(X_{n:n}|X_{1:n}) \prec_{\text{SI}} (Y_{n:n}|Y_{1:n}) \Leftrightarrow (X_{n:n}^*|X_{1:n}^*) \prec_{\text{SI}} (Y_{n:n}^*|Y_{1:n}^*)$$

■

Under the conditions of Theorem 3.3, an upper bound on  $\kappa(X_{1:n}, X_{i:n})$  is given by  $\kappa(Y_{1:n}, Y_{i:n})$ . *Avérous, Genest and Kochar* (2005) obtained an analytic expression for computing the exact values of the Kendall's  $\tau$  for any pair of order statistics from a random sample from a continuous distribution which in our case reduces to

$$\tau(Y_{1:n}, Y_{i:n}) = 1 - \frac{2(n-1)}{2n-1} \binom{n-2}{i-2} \sum_{s=0}^{n-i} \binom{n}{s} / \binom{2n-2}{n-i+s}. \quad (3.6)$$

## 4 Acknowledgments

The authors are grateful to the editors and two anonymous referees for their constructive comments and suggestions which led to this improved version of the paper.

## References

- [1] Avérous, J., Genest, C. and Kochar, S. C. (2005). On dependence structure of order statistics. *Journal of Multivariate Analysis* **94**, 159-171.
- [2] Barlow, R. E. and Proschan, F. (1981) *Statistical Theory of Reliability and Life Testing*. To Begin with, Silver Spring, Maryland.
- [3] Bickel, P. J. (1967). Some contributions to the theory of order statistics, in: L.M. LeCam, J. Neyman (Eds.), *Fifth Berkeley Symposium on Mathematics*

- and Statistics, Vol. 1, University of California Press, Berkeley, CA, 1967, pp. 575–591.
- [4] Boland ,P. J., Hollander, M., Joag-Dev, K. and Kochar, S. C. (1996). Bivariate dependence properties of order statistics. *Journal of Multivariate Analysis* **56**, 75-89.
- [5] Capéraà, P. and Genest, C. (1990). Concepts de dépendance et ordres stochastiques pour des lois bidimensionnelles. *Canadian Journal of Statistics* **18**, 315–326.
- [6] Chen, Y. P. (2007). A note on the relationship between Spearman’s  $\rho$  and Kendall’s  $\tau$  for extreme order statistics. *Journal of Statistical Planning and Inference* **137**, 2165–2171.
- [7] Deshpandé, J. V. and Kochar, S. C. (1983). Dispersive ordering is the same as tail ordering. *Advances in Applied Probability* **15**, 686–687.
- [8] Dolati, A., Genest , C. and Kochar, S. C. (2008). On the dependence between the extreme order statistics in the proportional hazards model. *Journal of Multivariate Analysis* **99**, 777-786.
- [9] Genest , C., Kochar, S. and Xu, M. (2009). On the range of heterogeneous samples. *Journal of Multivariate Analysis* **100**, 1587-1592.
- [10] Joe,H. (1997). *Multivariate Models and Dependence Concepts*. Chapman & Hall, London, 1997.
- [11] Khaledi, B. and Kochar, S C. Dependence orderings for generalized order statistics. *Statistics and Probability Letters* **73**, 357–367.
- [12] Kim, S. H. and David, H. A. (1990). On the dependence structure of order statistics and concomitants of order statistics. *Journal of Statistical Planning and Inference* **24**, 363-368.

- [13] Kimeldorf, G. and Sampson, A. R. (1989). A framework for positive dependence. *Annals of Institute of Statistical Mathematics* **41**, 31–45.
- [14] Kochar, S. C. and Korwar, R. (1996). Stochastic orders for spacings of heterogeneous exponential random variables. *Journal of Multivariate Analysis* **57**, 69–83.
- [15] Kochar, S.C. and Rojo, J. (1996). Some new results on stochastic comparisons of spacings from heterogeneous exponential distributions. *Journal of Multivariate Analysis* **59**, 272–281.
- [16] Kochar, S.C. and Xu, M. (2012). Some unified results on comparing linear combinations of independent gamma random variables. *Probability in the Engineering and Informational Sciences* **26**, 393–404.
- [17] Lehmann, E. L. (1966). Some concepts of dependence. *Annals of Mathematical Statistics* **37**, 1137–1153.
- [18] Nelsen, R. B. (1999). *An Introduction to Copulas*. Lecture Notes in Statistics No 139, Springer, New York, 1999.
- [19] Sathe, Y. S. (1988). On the correlation coefficient between the first and the  $r$ -th smallest order statistics based on  $n$  independent exponential random variables. *Communications in Statistics Theory and Methods* **17**, 3295–3299.
- [20] Scarsini, M. (1984). On measures of concordance. *Stochastica* **8**, 201–218.
- [21] Schmitz, V. (2004). Revealing the dependence structure between  $X_{(1)}$  and  $X_{(n)}$ . *Probability in Engineering and Information Sciences*. **123**, 41–47.
- [22] Schriever, B. F. (1987). An ordering for positive dependence. *Annals of Statistics* **15**, 1208–1214.
- [23] Shaked, M. and Shanthikumar, J. G. (2007). *Stochastic Orders and Their Applications*. Springer, New York.

- [24] Tchen, A. H. (1980). Inequalities for distributions with given marginals *Annals of Probability* **8**, 814–827.
- [25] Tukey, J. W. (1958). A problem of Berkson, and minimum variance orderly estimators,. *Annals of Mathematical Statistics* **29**, 588–592.
- [26] Xu, M. and Balakrishnan, N. (2012). On the sample ranges from heterogeneous exponential variables. *Journal of Multivariate Analysis* **109**, 1-9.
- [27] Yanagimoto, T. and Okamoto, M. (1969). Partial orderings of permutations and monotonicity of a rank correlation statistic. *Annals of Institute of Statistical Mathematics* **21**, 489-506. .
- [28] Yu, Y. (2021). On stochastic comparisons of order statistics from heterogeneous exponential samples. *Probability in Engineering and Information Sciences*. **35**, 532-537.