

4-1-2021

Events Associated with DNA Replication Disruption Are Not Observed in Hydrogen Peroxide-Treated *Escherichia Coli*

Chettar A. Hoff
Portland State University

Sierra S. Schmidt
Portland State University

Brandy J. Hackert
Portland State University

Travis K. Worley
Portland State University

Justin Courcelle
Portland State University

Follow this and additional works at: https://pdxscholar.library.pdx.edu/bio_fac



See next page for additional authors
Part of the [Biology Commons](#)

Let us know how access to this document benefits you.

Citation Details

Hoff, C. A., Schmidt, S. S., Hackert, B. J., Worley, T. K., Courcelle, J., & Courcelle, C. T. (2021). Events associated with DNA replication disruption are not observed in hydrogen peroxide-treated *Escherichia coli*. *G3 Genes/Genomes/Genetics*, 11(4), jkab044. <https://doi.org/10.1093/g3journal/jkab044>

This Article is brought to you for free and open access. It has been accepted for inclusion in Biology Faculty Publications and Presentations by an authorized administrator of PDXScholar. Please contact us if we can make this document more accessible: pdxscholar@pdx.edu.

Authors

Chettar A. Hoff, Sierra S. Schmidt, Brandy J. Hackert, Travis K. Worley, Justin Courcelle, and Charmain Courcelle

Events associated with DNA replication disruption are not observed in hydrogen peroxide-treated *Escherichia coli*

Chettar A. Hoff,[†] Sierra S. Schmidt ,[†] Brandy J. Hackert, Travis K. Worley, Justin Courcelle ,* and Charmain T. Courcelle

Department of Biology, Portland State University, Portland, OR 97201, USA

[†]These authors contributed equally to this work.

*Corresponding author: Department of Biology, Portland State University, Portland, OR 97201, USA. justc@pdx.edu.

Abstract

UV irradiation induces pyrimidine dimers that block polymerases and disrupt the replisome. Restoring replication depends on the *recF* pathway proteins which process and maintain the replication fork DNA to allow the lesion to be repaired before replication resumes. Oxidative DNA lesions, such as those induced by hydrogen peroxide (H₂O₂), are often thought to require similar processing events, yet far less is known about how cells process oxidative damage during replication. Here we show that replication is not disrupted by H₂O₂-induced DNA damage *in vivo*. Following an initial inhibition, replication resumes in the absence of either lesion removal or RecF-processing. Restoring DNA synthesis depends on the presence of manganese in the medium, which we show is required for replication, but not repair to occur. The results demonstrate that replication is enzymatically inactivated, rather than physically disrupted by H₂O₂-induced DNA damage; indicate that inactivation is likely caused by oxidation of an iron-dependent replication or replication-associated protein that requires manganese to restore activity and synthesis; and address a long standing paradox as to why oxidative glycosylase mutants are defective in repair, yet not hypersensitive to H₂O₂. The oxygen-sensitive pausing may represent an adaptation that prevents replication from occurring under potentially lethal or mutagenic conditions.

Keywords: RecF; glycosylases; DNA replication; DNA repair; manganese

Introduction

All cells must accurately duplicate their genomes to reproduce. However, DNA damage can block the replication machinery and prevent it from completing this task, resulting in mutations if the wrong base is incorporated, rearrangements if replication resumes from the wrong site, or cell lethality if the block to replication cannot be overcome (reviewed in (Courcelle and Hanawalt 2003)). In order to understand how the cell maintains its genome, it is important to characterize how replication accurately processes DNA damage in these situations.

DNA damage generated by UV irradiation has often been used as a model to study how cells respond to and recover from DNA damage (Courcelle et al. 2003). Irradiation with 254-nm light induces pyrimidine dimers that block DNA polymerases and disrupt replisome progression (Setlow et al. 1963; Chan et al. 1985; Courcelle et al. 1997). We use the term disruption to distinguish it from replisome pausing or inactivation. These latter terms imply that the replisome remains structurally intact, and could simply resume once a missing precursor or cofactor is resupplied. In *Escherichia coli*, the disruption of replication at UV-induced lesions involves uncoupling of the leading and lagging strand, exposure of the nascent DNA to exonucleolytic degradation, and the dissociation of several components of the replisome, including Pol III, the beta clamp, and the tau complex, whereas both helicase and

primase remain associated with the branched replication fork DNA (Courcelle and Hanawalt 1999; Pages and Fuchs 2003; Jeiranian et al. 2013). The integrity of the replication fork DNA and its fork structure remains intact throughout the recovery process, as no collapsed forks or broken linear DNA are observed (Courcelle et al., 2003; Chow and Courcelle, 2004). Restoring replication and the replisome after disruption requires enzymatic processing by several RecF pathway gene products which recruit RecA and function to maintain and process the replication fork DNA so that the lesion can be repaired by nucleotide excision repair (Courcelle et al. 1997; Courcelle and Hanawalt, 1999; Courcelle et al. 1999, 2003; Chow and Courcelle 2004; Courcelle et al. 2006; Bichara et al. 2007). In the absence of enzymatic processing by RecF, the DNA at the disrupted fork is extensively degraded and replication fails to resume (Courcelle et al. 1997, 1999, 2003). The recovery of replication is also heavily dependent on the removal of the UV lesions by the UvrABC excinuclease complex which incises a 12-bp region surrounding the damaged bases (Setlow et al. 1963; Courcelle et al., 1997, 1999). UvrD helicase, DNA polymerase I, and ligase then remove, resynthesize, and join the missing bases to complete repair (Caron et al. 1985; Van Houten 1990). In *uvrA*, *uvrB*, or *uvrC* mutants, these lesions are not removed, the replication forks remain blocked, and the resumption of DNA replication is severely impaired

Received: December 18, 2020. Accepted: February 05, 2021

© The Author(s) 2021. Published by Oxford University Press on behalf of Genetics Society of America.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted reuse, distribution, and reproduction in any medium, provided the original work is properly cited.

(Setlow *et al.* 1963; Howard-Flanders *et al.* 1969; Courcelle *et al.* 1999, 2003). In these repair-defective cells, high frequencies of chromosomal exchanges and extensive cell death are observed (Howard-Flanders and Theriot 1966; Rupp and Howard-Flanders 1968; Rupp *et al.* 1971; Courcelle and Hanawalt 2001; Courcelle *et al.* 2001; Bichara *et al.* 2007). In repair-proficient cells, these recombination events are efficiently suppressed, survival is greatly enhanced, and a robust recovery of replication is observed, indicating that the normal mechanism of recovery is integrated with lesion repair (Howard-Flanders *et al.* 1969; Rupp *et al.* 1971; Courcelle and Hanawalt 2001; Courcelle *et al.* 2001; Bichara *et al.* 2007).

In addition to UV, other environmental agents interact with and damage DNA, including ionizing radiation and a variety of chemicals (Singer and Kusmierek 1982; Hutchinson 1985). Of particular importance is the damage generated by both endogenous and exogenous sources of reactive oxygen species. DNA damage induced by reactive oxygen species is thought to be a primary source of the mutations in the etiology of cancer and aging in humans and is used as defense by our immune systems to kill invading pathogens (Fridovich 1978; Ames 1983; Cross *et al.* 1987; Nathan and Cunningham-Bussell 2013). Reactive oxygen species, such as that formed by H₂O₂ treatment, induce a broad spectrum of distinct base modifications in DNA (reviewed in Wallace 2002). These include lesions that have been characterized *in vitro* to maintain their coding specificities, such as dihydrothymine; that mispair during replication or transcription, such as 8-oxo-guanine; and others that block some DNA and RNA polymerases and are potentially lethal, such as thymine glycol (Ide *et al.* 1991; Evans *et al.* 1993; Efrati *et al.* 1999). Similar to UV damage, oxidized base damage is also repaired by excision and resynthesis of damaged bases. However, in this case, the excision is catalyzed by a lesion-specific DNA glycosylase that cleaves the glycosidic bond and releases the damaged bases. Most oxidative glycosylases also have an associated AP-lyase activity that incises the sugar-phosphate backbone (Baillly and Verly 1987; Melamede *et al.* 1994; Bhagwat and Gerlt 1996; Jiang *et al.* 1997). The apurinic or apyrimidinic (AP) site is then either incised or processed by an AP endonuclease to generate a clean 3'-OH that can be extended and joined by DNA polymerase I and ligase to restore the template's integrity (reviewed in Wallace, 1994).

Little is known about how replication recovers after DNA damage induced by oxygen free radical species. It is often assumed that the processing and recovery of replication in the presence of oxidative DNA damage will mimic those observed in the presence of UV-induced damage. However, mutants lacking the DNA glycosylases responsible for repairing oxidized-base damage are as resistant as wild-type cells when treated with H₂O₂ (Laspia and Wallace 1988; Asad *et al.* 1995; Saito *et al.* 1997; Schalow *et al.* 2011). Similarly, mutants deficient in nucleotide excision repair are also resistant to H₂O₂ (Imlay and Linn 1987; Asad *et al.* 1995; Schalow *et al.* 2011). Recently, we also observed that the recovery of replication after oxidative challenge had a unique requirement for manganese, something that is not observed after UV-induced damage (Hutfilz *et al.* 2019). These observations raise the possibility that the mechanism by which replication deals with H₂O₂-induced DNA damage is distinct from that seen after UV. Here we considered this possibility, and found that oxidative lesions induced by H₂O₂ do not disrupt replication, and that the resumption of DNA synthesis does not require any enzymatic processing by RecF, implying that the replisome

remains intact but is simply inactive due to the lack of metal cofactors.

Materials and methods

Bacterial strains and plasmids

SR108 (*thyA36 deoC2* derivative of W3110), HL921 (SR108 Δ (*srIR-recA*)306::Tn10), HL922 (SR108 *recB21recC22 argA81*::Tn10), HL952 (SR108 *uvrA*::Tn10), CL008 (SR108 *recG258*::Tn5), CL544 (SR108 *recR6212*::*cat*), CL577 (SR108 *ruvC53eda-51*::Tn10), CL579 (SR108 *recF6206*::*tet^R*), and CL915 (SR108 *recN*::*cat*) have been previously described (Mellon and Hanawalt 1989; Courcelle *et al.* 1997, 1999, 2003; Donaldson *et al.* 2004, 2006; Cole *et al.* 2018). CL1746 (SR108 *nth*::*kan^R nei*::*cat*) was constructed by P1 transduction of the *nei*::*cat* allele from CL1005 (SR108 *nei*::*cat* (Schalow *et al.* 2011)) into CL1006 (SR108 *nth*::*kan^R*) (Schalow *et al.* 2011). CL1941 (SR108 *nth*::*kan^R nei*::*cat fpg*::*tet^R*) was constructed by P1 transduction of the *fpg*::*tet^R* allele from CL1778 (SR108 *fpg*::*tet^R* (Schalow *et al.* 2011)) into CL1746. CL1155 (DY329 *xthA*::*cat*) was constructed by gene replacement using the recombineering strain DY329 (Yu *et al.* 2000). The *cat* cassette from pPCR-Script Cam was amplified using the primers 5'GTCTCTTTTAAATCAACGGCCTGCGGCCAGACCTACTGTGACGGAAGATCACTTCG and 5'CGGTTTTTCCA TGCTGCGGATTTTCATAGTCGATGCCGGTACCAGAATAGACATAAGCG. The PCR product was transformed into DY329 to generate CL1155, selecting for chloramphenicol resistance. The gene replacement was then moved into SR108 by standard P1 transduction to generate CL1168 (SR108 *xthA*::*cat*). All strains used in this work are summarized in Supplementary Table S1.

pBR322 is a medium copy number, ColE1-based, 4.4-kb plasmid (Promega).

H₂O₂ survival assays

Fresh overnight cultures were diluted 1:100 in Davis medium supplemented with 0.4% glucose, 0.2% casamino acids and 10 μ g/ml thymine (DGcthy), grown at 37°C to an OD₆₀₀ of 0.3, and then treated with 10 mM H₂O₂. At the times indicated, 0.1-ml aliquots of each culture were removed and serially diluted in 10-fold increments into DGcthy medium. Triplicate 10- μ l aliquots of each dilution were then spotted on Luria-Bertani (LB) plates supplemented with 10 μ g/ml thymine (LBthy). Viable colonies were counted following overnight incubation at 37°C.

UV survival assays

UV irradiation used a 15-W germicidal lamp (254 nm) at an incident dose of 0.9 J/m²/s. Fresh overnight cultures were diluted 1:100 in DGcthy medium and grown at 37°C to an OD₆₀₀ of 0.3. At this time, 0.1-ml aliquots of each culture were removed and serially diluted in 10-fold increments into DGcthy medium. Triplicate 10- μ l aliquots of each dilution were then spotted on LBthy plates and irradiated with increasing doses of UV as indicated. Viable colonies were counted following overnight incubation at 37°C.

Lesion frequency

For UV irradiation, fresh overnight cultures were diluted 1:100 and grown at 37°C in DGcthy medium to an OD₆₀₀ of 0.3. Where indicated 200 μ M manganese (II) chloride (MnCl₂·4H₂O) was added to the medium as well. At this time, cultures were irradiated with an incident dose of 50 J/m² and then returned to 37°C to allow recovery. For H₂O₂ challenge, fresh overnight cultures were diluted 1:100 in DGcthy medium supplemented with MnCl₂

as indicated, grown at 37°C to an OD₆₀₀ of 0.3, and then treated with 10 mM H₂O₂ for 5 min at 37°C. Cells were filtered on 0.45-μm membranes (Fisherbrand) to remove excess H₂O₂ from the medium, resuspended in fresh DGcthy medium supplemented with manganese as appropriate to initial growth conditions and returned to 37°C for the duration of the time course. For both treatments, a 0.75-ml aliquot was transferred at the times indicated to an equal volume of 2× 200 mM NaCl, 20 mM Tris, pH 8.0, 40 mM EDTA, pH 8.0 (NET). Cells were pelleted and resuspended in 0.14-ml of lysis buffer (1 mg/ml lysozyme, 0.5 mg/ml RNaseA in 10 mM Tris, pH 8.0, 1 mM EDTA, pH 8.0) and incubated for 30 min at 37°C. Then, 0.01-ml 10 mg/ml ProteinaseK and 0.01-ml 20% Sarkosyl were added to the samples and incubation was continued for 30 min at 37°C. The samples were then extracted once with four volumes of phenol:chloroform, followed by two volumes of chloroform, and dialyzed against 200-ml 1 mM Tris, pH 8.0, 1 mM EDTA, pH 8.0 for 30 min using 47-mm Millipore 0.025-μm pore disks.

For UV-irradiated samples, 15 μl of each DNA sample was treated with reaction buffer (12.5 mM sodium phosphate [pH 6.8], 5 mM EDTA [pH 8.0], 50 mM NaCl, 0.5 mM dithiothreitol, 0.005% Triton X-100, 0.1 mg/ml bovine serum albumin) supplemented with either no enzyme or 2 U T4 endonuclease V (T4 Endo V; Trevigen) for 1 h at 37°C. For H₂O₂-treated samples, 15 μl of each DNA sample was treated with reaction buffer (30 mM EDTA [pH 8.0], 22.5 mM NaCl, 5 mM Tris [pH 8.0]) supplemented with either no enzyme or 0.53 μM Fpg glycosylase and 2.7 μM Endo III for 1 h at 37°C. Enzyme preparations were titrated using purified undamaged genomic DNA as a template. The highest enzyme concentration that did not exhibit nonspecific activity on undamaged DNA was used. For the preparations in our lab, this corresponded to 2 U T4 Endo V, 0.53 μM Fpg glycosylase and 2.7 μM Endo III. Treated samples were then electrophoresed on a 0.5% alkaline agarose gel in 30 mM NaOH, 1 mM EDTA at 30 V for 16 h, stained and visualized with ethidium bromide. The intensity of each high-molecular-weight band was determined using ImageQuant software (GE Biosciences). The fraction of lesion-free DNA fragments was quantified as a ratio of high-molecular-weight DNA in the T4 Endo V- or Fpg/Endo III-treated band (Enz_T) to the band without enzyme treatment (NoEnz_T) at each time point, T. To normalize for any nicks or AP sites present in the DNA before UV or H₂O₂ treatment, the ratio obtained at each time point was divided by the ratio calculated at the pretreatment time point as follows: (Enz_T/NoEnz_T)/R₀, where R₀ is (Enz_T/NoEnz_T) at the time immediately before UV or H₂O₂ addition.

DNA synthesis rate

For experiments using UV irradiation, overnight cultures were diluted 1:100 and grown at 37°C in DGcthy to an OD₆₀₀ of 0.25–0.35. Where indicated manganese was added to the medium as described above. At this time, half of the cells were mock irradiated, while the other half of the culture was irradiated with 50 J/m².

For experiments using H₂O₂, overnight cultures were diluted 1:100 and grown at 37°C to an OD₆₀₀ of 0.25–0.35 in DGcthy supplemented with 200 μM MnCl₂ where indicated. At this time, half of the cells were mock treated, while the remaining culture was exposed to 10 mM H₂O₂ for 5 min at 37°C. Following either mock or H₂O₂ treatment, cells were filtered on 0.45-μm membranes to remove excess H₂O₂ from the medium and resuspended in fresh DGcthy medium either supplemented with Mn or not based on initial growth conditions.

For both UV irradiation and H₂O₂ experiments, cultures were returned immediately to 37°C after treatment to allow recovery

and continued growth. At the times indicated, duplicate 0.5-ml aliquots of culture were pulse-labeled with 0.5 μCi/ml [³H]thymidine for 2 min at 37°C. Cells were then lysed, and the DNA was precipitated in cold 5% trichloroacetic acid and filtered onto Millipore glass fiber filters. The amounts of ³H on each filter were determined by scintillation counting.

Two-dimensional agarose gel electrophoresis

Cells containing the plasmid pBR322 were grown overnight in DGcthy medium supplemented with 100 μg/ml ampicillin (amp). A 0.2-ml aliquot of this culture was pelleted and resuspended in 20-ml DGcthy with 200 μM MnCl₂ where indicated and grown without amp selection to OD₆₀₀ of 0.5. At this time, cultures were either UV-irradiated with an incident dose of 50 J/m² and then returned to 37°C to allow recovery, or treated with 10 mM H₂O₂ for 5 min at 37°C, filtered on 0.45-μm membranes to remove excess H₂O₂ from the medium, then resuspended in fresh DGcthy medium and returned to 37°C for the duration of the time course. At the times indicated, a 0.75-ml aliquot was transferred to an equal volume of 2× NET. Total DNA (genomic and plasmid) was then purified from these cultures as described above.

DNA samples were dialyzed against 200-ml 1 mM Tris, pH 8.0, 1 mM EDTA, pH 8.0 for 30 min using 47-mm Millipore 0.025-μm pore disks and digested with PvuII restriction endonuclease (Thermo Fisher Scientific) overnight at 37°C. Samples were then extracted once with one volume of chloroform, loaded directly on a 0.4% agarose gel in 1× TBE (Tris-borate-EDTA) and DNA was separated initially at 1 V/cm for 15 h. For the second dimension, lanes were excised, rotated 90°, recast in 1% agarose in 1× TBE and electrophoresed at 6.5 V/cm for 7 h. DNA in the gel was transferred to Hybond N+ nylon membrane by standard Southern blotting techniques, and the plasmid DNA was detected using ³²P-labeled pBR322 plasmid prepared by random prime labeling (Agilent) using dCTP (3000 Ci/mmol; PerkinElmer) and visualized using a Storm 840 PhosphorImager (GE Biosciences) and its associated ImageQuant software.

Data availability

Strains used in these experiments are available upon request and can be found in Supplementary Table S1. Supplemental material is available at figshare: <https://doi.org/10.25387/g3.13426079>.

Results

DNA replication recovers in the absence of lesion removal following oxidative challenge

To examine how replication responds to H₂O₂-induced DNA damage *in vivo*, we compared the replication occurring in H₂O₂-treated wild-type cells to that occurring in *fpg nei nth* mutants, which lack the three predominant oxidative DNA glycosylases. Consistent with previous reports (Laschia and Wallace 1988; Asad et al. 1995; Saito et al. 1997; Schalow et al. 2011), the absence of Fpg, Endonuclease III (Endo III) and Endonuclease VIII (Endo VIII) did not render cells hypersensitive to H₂O₂ (Figure 1A). However, the absence of these glycosylases impaired or prevented the removal of the oxidative lesions recognized by these enzymes *in vivo* (Figure 1, B and C). To monitor repair, cultures were exposed to 10 mM H₂O₂ for 5 min, then filtered and resuspended in fresh medium to allow recovery in the absence of H₂O₂. Lesion removal was assayed at various times during the recovery by treating purified total genomic DNA with Fpg and Endo III (Schalow et al. 2011). Purified Endo VIII was not utilized because it lacks




Figure 1 DNA replication recovers even in the absence of lesion removal following H_2O_2 treatment, but not UV irradiation. (A) The survival of wild-type (open squares) and *fpg nei nth* (filled triangles) cells following 10 mM H_2O_2 treatment is plotted. Graphs represent the average of three independent experiments. Error bars represent one standard error of the mean. (B) Wild-type (top) and *fpg nei nth* (bottom) cultures were treated with 10 mM H_2O_2 for 5 min, allowed to recover and then genomic DNA was purified at the indicated times. DNA was either treated with Fpg and Endo III (+Fpg, Endo III), or no glycosylase (untreated) for 1 h at 37°C and then analyzed on alkali agarose gels. A representative gel is shown. Arrows indicate lesion-free DNA. (C) The percentage of lesion-free, high-molecular-weight DNA in Fpg/Endo III-treated samples is plotted for each time point relative to mock-treated samples. Graphs represent an average of at least two independent experiments. (D) [³H]thymidine was added to cultures grown with manganese for 2 min at the indicated times following H_2O_2 treatment at time zero. The rate of DNA synthesis (³H/2 min) relative to the amount incorporated immediately prior to exposure is plotted for wild-type and *fpg nei nth* cultures exposed to mock treatment (open symbols) or 10 mM H_2O_2 for 5 min (filled symbols). Graphs represent an average of at least two independent experiments. Error bars represent one standard error of the mean. (E) The survival of wild-type (open squares) and *uvrA* (filled triangles) cells following UV irradiation is plotted. Graphs represent the average of three independent experiments. Error bars represent one standard error of the mean. (F) Wild-type (top) and *uvrA* (bottom) cultures were UV irradiated at 50 J/m², then genomic DNA was purified at the indicated times and either treated with T4 endonuclease V (+T4 endo V) or no T4 endo V (untreated) for 1 h at 37°C and analyzed on alkali agarose gels. A representative gel is shown. Arrows indicate lesion-free, high-molecular-weight DNA. (G) The percentage of lesion-free, high-molecular-weight DNA in T4 Endo V-treated samples is plotted for each time point relative to mock-treated samples. Graphs represent an average of at least two independent experiments. (H) [³H]thymidine was added to cultures grown with manganese for 2 min at the indicated times following UV irradiation at time zero. The rate of DNA synthesis (³H/2 min) relative to the amount incorporated immediately prior to exposure is plotted for wild-type and *uvrA* cells exposed to mock treatment (open symbols) or 50 J/m² UV irradiation (filled symbols) treatment. Graphs represent an average of at least two independent experiments. Error bars represent one standard error of the mean.

enzymatic turnover *in vitro* and its substrate specificity overlaps with Fpg and Endo III (Melamede et al. 1994; Jiang et al. 1997). The glycosylase activity of Fpg and Endo III results in incisions at lesions recognized by the three oxidative glycosylases, while the AP lyase activity associated with these enzymes nicks the DNA backbone (Bailly and Verly 1987; Melamede et al. 1994; Bhagwat and Gerlt 1996; Jiang et al. 1997). Any non-incised AP sites created by base excision would also be cleaved under the alkali conditions used during electrophoresis. Thus, following alkali-agarose gel electrophoresis, the presence of oxidative lesions in the genomic DNA is observed as the loss of high-molecular-weight species. In wild-type cells, prior to treatment, DNA fragments averaged greater than 40 kb in length, which is the approximate limit of resolution in our agarose gels. Incubation of the genomic DNA with Fpg and Endo III glycosylases resulted in a loss of high-molecular-weight fragments at times immediately after H₂O₂ exposure, indicating the presence of lesions. Over time, the number of lesions in the DNA decreased and within 30 min, greater than 60% of the DNA had been restored as lesion-free high-molecular-weight fragments (Figure 1C). In contrast, the removal of lesions in H₂O₂-treated *fpg nei nth* cultures was severely impaired with only 18% of high-molecular-weight DNA restored by the end of the 60-min recovery time course (Figure 1C). The actual amount of repair is likely to be significantly less as much of this high-molecular-weight DNA is likely to represent newly synthesized DNA, rather than repair events (see Figure 1D and text below).

To examine how H₂O₂-induced damage affects DNA replication, cultures were treated with 10 mM H₂O₂ for 5 min, as before, and at the times indicated, aliquots of cultures were pulse-labeled with [³H]thymidine for 2 min before the DNA was precipitated and the amount of ³H incorporated was quantified. The rate of DNA replication (³H incorporation in the DNA/2 min) was determined for each time point and expressed relative to the rate immediately prior to treatment. In both wild-type and *fpg nei nth* cells, a rapid inhibition of DNA replication was observed immediately following H₂O₂ treatment. Surprisingly, replication rapidly resumed even in the absence of lesion removal and the *fpg nei nth* mutant restored DNA synthesis with kinetics that were modestly altered or reduced relative to wild-type cells (Figure 1D). The result indicates that the resumption of replication following H₂O₂ treatment does not require lesion repair by the glycosylases tested.

The recovery of replication in the absence of oxidative lesion repair stands in contrast to what occurs after UV irradiation. *uvrA* mutants are defective for nucleotide excision repair and are unable to remove UV-induced lesions (reviewed in (Van Houten 1990)). Immediately after treatment with 50 J/m² UV light, a similar inhibition of replication is observed in both wild-type and *uvrA* cultures. Using the pyrimidine dimer-specific glycosylase, T4 Endonuclease V (T4 Endo V), to monitor repair (Spivak and Hanawalt 1995) and measuring the rate of replication as before demonstrates that in wild-type cultures, replication resumes at a time that correlates with the removal of the lesions. However, in *uvrA* mutants, the lesions are not removed, DNA synthesis does not resume, and high levels of lethality are observed (Figure 1, E–H).

Thus, unlike UV-induced damage, the results demonstrate that lesions removed by Fpg, Endo III, or Endo VIII do not prevent replication from progressing *in vivo*, and imply that these lesions do not disrupt replication. However, an inhibition of replication is clearly observed immediately following exposure to H₂O₂. This latter observation would appear to suggest that replication is disrupted by these lesions. We considered two possibilities to

explain these apparently contradictory observations. It is possible that other forms of H₂O₂-induced DNA damage, not recognized by these glycosylases, disrupt replication and are responsible for the inhibition that is observed. Alternatively, the inhibition could result from an oxidative sensor or another aspect of the cellular oxidative stress response that is not directly associated with DNA damage.

The recovery of replication following H₂O₂ occurs independently of *recF* pathway and lacks intermediates associated with disrupted replication forks

To determine whether the observed replication inhibition results from disruption by other forms of damage that are not repaired by the three predominant oxidative DNA glycosylases, we examined *recF* mutants. In contrast to other DNA damage hypersensitive mutants, the defect in *recF* pathway mutants is specific to their ability to restore replication following disruption (Supplementary Figures S1 and S2). Other hypersensitive mutants, including *recBCD*, *recG*, and *ruvABC* process and restore replication normally following disruption, arguing that cell lethality in these mutants arises from other causes (Courcelle et al. 1997; Courcelle and Hanawalt 1999; Courcelle et al. 2003; Chow and Courcelle 2004; Donaldson et al. 2004; Donaldson et al. 2006; Wendel et al. 2014). RecF processes and maintains DNA at replication forks following disruption by DNA damage, and is required for DNA synthesis to resume (Hori and Clark 1973; Courcelle et al. 1997; Courcelle et al. 1999; Courcelle et al. 2003). As shown in Figure 2A, we found that although RecF contributes to survival following UV irradiation, it does not contribute to survival following H₂O₂ exposure. Indeed, the absence of RecF conferred a modest resistance in the presence of H₂O₂ compared to wild-type cells. As controls, we examined *uvrA* and *xthA* (encoding the dominant AP endonuclease, Exo III) mutants (Figure 2A), which have previously been shown to be hypersensitive to UV and H₂O₂, respectively (Hill 1958; Setlow et al. 1963; Demple et al. 1983).

We also examined whether the absence of RecF impaired the recovery of DNA synthesis following exposure to H₂O₂, as before. Although RecF was required for replication to resume after UV irradiation (Figure 2B and (Courcelle et al. 1997, 1999)), DNA synthesis resumed independently of RecF following H₂O₂ treatment. Taken together, these results are consistent with the idea that the inhibition of replication caused by H₂O₂ is not due to disruption of replication by DNA damage, and demonstrate that RecF-mediated processing is not required for survival or for replication to resume.

Previous studies have shown that replication forks disrupted by UV-induced DNA damage undergo processing events that displace the DNA polymerase and restore the region to its double-stranded form, allowing repair enzymes to access the lesion and effect repair (Courcelle et al. 2003; Chow and Courcelle 2004; Donaldson et al. 2006; Belle et al. 2007; Jeiranian et al. 2012). The processing forms unique structural intermediates that can be observed using 2D agarose gel electrophoresis. To determine if similar structural intermediates or processing occurs following the inhibition of replication observed after H₂O₂ stress, we compared the structural intermediates that arise on replicating plasmids after treatment with H₂O₂ to that occurring after UV irradiation using 2D agarose gel electrophoresis. To this end, wild-type cultures of *E. coli* containing the plasmid pBR322 were either exposed to H₂O₂ or irradiated with UV, as before. At various times following treatment, total genomic DNA was isolated, digested with the restriction enzyme, PvuII, to linearize the plasmid just




Figure 2 RecF is not required for survival or replication recovery following oxidative challenge. (A) The survival of wild-type (open squares), *recF* (filled diamonds), *uvrA* (filled squares) and *xthA* (filled circles) cells following UV irradiation or 10 mM H₂O₂ treatment is plotted. Wild-type plots for each treatment and *uvrA* plot for UV irradiation are reproduced from Figure 1 for comparison. Graphs represent the average of three independent experiments. Error bars represent one standard error of the mean. (B) Data were obtained and plotted as for Figure 1. The rate of DNA synthesis (³H/2 min) relative to the amount incorporated immediately prior to exposure is shown for wild-type and *recF* cells exposed to mock treatment (open symbols), UV irradiation, or 10 mM H₂O₂ exposure for 5 min (filled symbols) in Mn-supplemented medium. Graphs represent an average of at least two independent experiments. Error bars represent one standard error of the mean.

downstream of its origin of replication, then separated and analyzed by 2D agarose gel electrophoresis. Non-replicating plasmid molecules migrate as a linear 4.4-kb fragment and form the prominent spot observed on the gels. Replicating molecules form Y-shaped structures that migrate more slowly due to their larger size and nonlinear shape, and appear as an arc radiating out from the prominent linear spot (Figure 3, A and B). Following the disruption of replication by UV-induced damage, the displacement of the DNA polymerase and processing of the replication fork DNA form a set of molecules that contain four arms (X- or double Y-shaped structures) which migrate slower through the gel and accumulate in the cone region located above the Y-arc (Figure 3, A and B). Previous work from our lab has shown that these cone region intermediates persist until the time corresponding with when the lesions are repaired and DNA replication resumes (Courcelle et al. 2003). When we examined H₂O₂-treated wild-type cells using this technique, we did not observe the accumulation of any cone region intermediates, despite the inhibition of replication (Figure 3C). Instead, only normal Y-shaped intermediates were observed throughout the recovery time course. The absence of any arrested replication fork intermediates indicates that H₂O₂-induced lesions are processed differently from UV lesions and suggests that following H₂O₂-treatment the replication fork machinery is not disrupted and the replication forks do not require processing similar to that which occurs after UV.

DNA replication, but not DNA repair, contains an iron-sensitive component that is inactivated and restored by manganese after H₂O₂ treatment

In previous work, we found that limiting levels of manganese in the growth medium impaired the recovery of DNA replication (Hutfilz et al. 2019). Based on these results, we proposed that H₂O₂ treatment oxidizes and inactivates some essential iron-dependent enzyme(s) and that manganese was needed to re-metallate these or alternative enzymes before genomic DNA replication can resume. These observations would be consistent with the results presented here and suggest that an oxidized 'sensor' protein is responsible for the observed inhibition rather than H₂O₂-induced lesions. The sensor could be a replication protein, or a protein that associates with the replisome to prevent DNA synthesis from progressing during oxygen stress. Alternatively, the sensor could be a repair protein(s) that is inactivated to prevent repair, thereby preventing replication from resuming. To examine this latter possibility, we compared the ability of cells to repair H₂O₂-induced lesions under varying manganese growth conditions. To this end, cultures were grown in defined medium supplemented with 200 μM MnCl₂ or without additional metals, treated with H₂O₂, and monitored for the recovery of replication and repair of the H₂O₂-induced lesions as before. Although the absence of manganese reduced the ability of the cells to resume replication, it did not alter the rate or time that the H₂O₂-induced lesions were removed (Figure 4, A–C). Consistent with this, no replication fork processing intermediates were observed by 2D agarose gel analysis either in the presence or absence of manganese (Supplementary Figure S3). As a control, we also examined the rate of DNA synthesis and lesion removal following UV irradiation in the presence or absence of supplemented manganese. In contrast, to H₂O₂ treatment, the recovery of replication following UV irradiation occurred with similar kinetics in both the presence and absence of manganese supplementation. Repair of UV-induced lesions was similarly unaffected by the presence of




Figure 3 The inhibition, and subsequent recovery, of DNA replication after oxidative stress is not associated with DNA damage processing intermediates. (A) The migration pattern of PvuII-digested pBR322 plasmid DNA observed by 2D gel electrophoresis is diagrammed. Non-replicating plasmids run as a 4.4-kb linear fragment. Replicating plasmids form Y-shaped structures that migrate slower than non-replicating linear DNA and form an arc that extends from the linear region. Following UV irradiation, double-Y or X-shaped intermediates are observed that migrate in the cone region behind the arc of Y-shaped molecules. Two-dimensional agarose gels of total DNA purified from wild-type cultures containing plasmid exposed to (B) 50 J/m² UV irradiation or (C) 10 mM H₂O₂ for 5 min were probed with labeled pBR322 at the indicated times after treatment. Arrow indicates double-Y or X-shaped intermediates in the cone region.

manganese (Figure 4, D–F). The observation that manganese promotes the recovery of replication in H₂O₂-treated, but not UV-treated cultures, supports the idea that H₂O₂-induced replication inhibition results from an oxidized sensor that stalls replication, and is not due to disruption by DNA damage. Further, the results also demonstrate that lesion removal does not require manganese supplementation.

Taken together, the observations that replication recovers and continues in the presence of H₂O₂-induced DNA lesions, that RecF processing is not required for the H₂O₂-inhibited replication to resume, that processing intermediates at inhibited replication forks are not observed, and that the initial inhibition is not caused by a lack of repair, indicate that H₂O₂-induced DNA lesions do not disrupt replication *in vivo*. Instead, we infer that the initial inhibition of replication is likely to be caused by an iron-dependent replication protein or replication-associated protein that is oxidized and inactivates replication under these conditions.

Discussion

The recovery and continuation of DNA synthesis suggests that the predominant lesions generated by H₂O₂ do not block replication progression *in vivo*. Early studies showed that thymine glycols, which comprise up to 30% of the H₂O₂-induced base damage, represent strong blocks to *E. coli* DNA polymerase I and phage T4 DNA polymerase *in vitro* (Ide et al. 1985; Rouet and Essigmann 1985; Clark and Beardsley 1986; Hayes and Leclerc 1986; Blakely et al. 1990), and it was often inferred that this would extend generally to other DNA polymerases. However, since these initial studies, several other polymerases have been examined and found to efficiently bypass this lesion, including many in humans (Fischhaber et al. 2002; Takata et al. 2006; Belousova et al. 2010; Yoon et al. 2010; Hogg et al. 2011; Makarova et al. 2018). Further, auxiliary proteins associated with the replisome, such as single-strand binding protein and processivity factors, can increase bypass efficiency in some cases (Maga et al. 2007; McCulloch et al. 2009). Finally, many of the other prominent lesions, such as 8-oxo-guanine, dihydrothymine, and uracil glycols are bypassed with fairly high frequencies *in vitro* (Purmal et al. 1994; Shibutani and Grollman 1994; Purmal et al. 1998; McCulloch et al. 2009). Taken together, these observations suggest that the replicative polymerases of *E. coli*, in the context of the full replisome, may bypass these lesions during genomic replication.

It is also possible that bypass frequencies at these oxidative lesions increase following the upregulation of manganese import that occurs after oxidative challenge (Kehres et al. 2002; Anjem et al. 2009). Consistent with this idea, we have previously shown that the recovery promoted by Mn is associated with an elevated level of mutagenesis that depends on both replicative and translesion polymerases (Hutfilz et al. 2019). Typical intracellular manganese concentrations in minimal medium can range from 15 to 150 μM when Mn transporters are fully induced (Anjem et al. 2009; Martin et al. 2015). The Mn-dependent mutagenesis may arise by reducing polymerase fidelity. Several studies have shown that manganese alters polymerase fidelity *in vitro*, including *E. coli* DNA polymerase I, human polymerases iota, Dpo4, and Primpol, and several polymerases used for PCR-mediated mutagenesis (Kunkel and Loeb 1979; Goodman et al. 1983; Beckman et al. 1985; Vaisman et al. 2005; Frank and Woodgate 2007; Tokarsky et al. 2017). If this phenomenon also extended to the replicative polymerase III in *E. coli*, it could explain why replication is initially inhibited, but then recovers in the absence of repair. However, the need for lesion bypass also predicts that replication should be disrupted by thymine glycols and depend on RecF processing for resumption, which is not observed.

Here we show that the inhibition and recovery of replication following H₂O₂-treatment is distinct from that following UV in several aspects. Following UV, the inhibition of replication results from disruption of the replisome by UV-induced pyrimidine dimers. In addition to the core proteins of the replication machinery, restoring replication following disruption *in vivo* requires processing of the fork by RecF and the repair of the lesions. In contrast, the inhibition of replication following H₂O₂ treatment appears to result from the inactivation of the replisome by an iron-sensitive protein. Restoring replication does not require lesion repair but is promoted by the presence of manganese. We did not detect any difference in the quantity of lesions present in DNA purified from cells grown in the presence or absence of manganese (Figure 4), arguing against the idea that the difference in recovery is due to a protective effect of manganese preventing




Figure 4 The absence of Mn in growth medium inhibits replication recovery, but not DNA repair, following oxidative stress. (A) Data were obtained and plotted as for Figure 1. The rate of DNA synthesis ($^3H/2 min$) relative to the amount incorporated immediately prior to exposure is plotted for wild-type cells exposed to mock treatment (open symbols) or 10 mM H_2O_2 (filled symbols) treatment in the presence or absence of Mn. Mn-supplemented plot is reproduced from Figure 1C for the purpose of comparison and control. Graphs represent an average of at least two independent experiments. Error bars represent one standard error of the mean. (B) Wild-type cultures grown with (top) or without Mn (bottom) were treated with 10 mM H_2O_2 for 5 min, allowed to recover and then genomic DNA was purified at the indicated times. DNA was either treated with Fpg and Endo III (+Fpg, Endo III), or no glycosylase (untreated) for 1 h at 37°C and then analyzed on alkali agarose gels. A representative gel is shown for each treatment. Arrows indicate lesion-free DNA. (C) The percentage of lesion-free, high-molecular-weight DNA in Fpg/Endo III-treated samples is plotted for each time point relative to mock-treated samples. Graphs represent an average of at least two independent experiments. Error bars represent one standard error of the mean. (D) The rate of DNA synthesis ($^3H/2 min$) relative to the amount incorporated immediately prior to exposure is plotted for wild-type cells exposed to mock treatment (open symbols) or 50 J/m² UV irradiation (filled symbols) in the presence or absence of Mn. Mn-supplemented plot is reproduced from Figure 1. Graphs represent an average of at least two independent experiments. Error bars represent one standard error of the mean. (E) Wild-type cultures grown with (top) or without Mn (bottom) were UV irradiated at 50 J/m², then genomic DNA was purified at the indicated times and either treated with T4 endo V (+T4 Endo V) or no T4 endo V (untreated) for 1 h at 37°C and analyzed on alkali agarose gels. A representative gel is shown for each treatment. Arrows indicate lesion-free DNA. (F) The percentage of lesion-free, high-molecular-weight DNA in T4 Endo V-treated samples is plotted for each time point relative to mock-treated samples. Graphs represent an average of at least two independent experiments. Error bars represent one standard error of the mean.

lesion formation. However, we cannot exclude the possibility that different ratios or otherwise novel lesions are generated in these two conditions. Other differences that have been noted are that the increased mutation frequency after UV depends entirely on the UmuC polymerase, whereas after H₂O₂ the induced mutagenesis is more modest and involves both replicative and translesion DNA polymerases (Bagg et al. 1981; Imlay and Linn 1987; Napolitano et al. 2000; Hutfilz et al. 2019).

The results offer an explanation that may resolve a long standing paradox as to why oxidative glycosylase mutants are defective in repair, but not hypersensitive to H₂O₂ (Laschia and Wallace 1988; Asad et al. 1995; Saito et al. 1997; Spek et al. 2001; Schalow et al. 2011). However, several critical questions remain to be addressed as to the mechanism of cell death caused by oxygen free radicals. H₂O₂ induces a modest SOS response following H₂O₂ treatment (Imlay and Linn 1987; Goerlich et al. 1989; Gifford et al. 2000). Our current understanding suggests that the SOS response is typically induced in response to disrupted replication forks or double-strand breaks that subsequently form replication forks during repair (reviewed in (Walker 1984; Kogoma 1997; Hanawalt 2001; Crowley and Courcelle 2002)). The modest SOS induction observed could suggest that the replication fork still encounters some disruptive lesions either before inactivation occurs or due to incomplete inhibition of the forks. Early work has shown that cell lethality follows a curious bimodal pattern as the H₂O₂ concentration increases (Imlay and Linn 1986, 1987). 'Mode 1' killing occurs at concentrations below 2 mM H₂O₂ and appears to involve RecA and the SOS response. It could be these lower doses do not inactivate the replication machinery, allowing it to progress and be disrupted by the lesions, thereby inducing SOS. 'Mode 2' killing occurs at high concentrations, and may involve the inactivation of the replisome as we observed here. The H₂O₂ concentrations we used here would fall under the conditions of 'mode 2' killing. We did not explore the conditions associated with low dose or 'mode 1' killing, as the reduced number of lesions generated under these conditions would make their detection prohibitive with the assays employed here.

Alternatively, the modest SOS induction may suggest that oxidative damage induces double-strand breaks that are removed from the replication fork and arise elsewhere in the genome. This latter explanation would explain the lack of requirement for RecF which is not required for the repair of double-strand breaks and would also explain the H₂O₂ hypersensitivity of *recBC* mutants, which are defective in the repair of double-strand breaks (Willets and Clark 1969; Willets and Mount 1969; Horii and Clark 1973; Youngs and Bernstein 1973; Howard-Flanders 1975; Lloyd and Thomas 1983). Many iron-containing proteins bind DNA and James Imlay's group has proposed that these binding sites may be hotspots for DNA strand breaks due to Fenton chemistry generating oxygen free radicals at these loci (Imlay 2003; Djaman et al. 2004; Imlay 2014).

It is possible that the oxygen-sensitive pausing of replication represents an adaptive response to prevent genomic replication from occurring during periods of oxygen stress when potentially lethal or mutagenic damage is present. Pausing replication to allow more time for repair to occur has been demonstrated to be beneficial both for reducing mutations and increasing survival (Castellani et al. 1964; Ganessan and Smith 1969). This may also explain the increased resistance of *recF* mutants to H₂O₂ treatment, which are impaired in resuming replication after disruption (Figure 2A). In the absence of manganese, this inhibition can last for several hours, without compromising viability (Imlay and Linn 1986; Hutfilz et al. 2019).

There are other examples of oxygen sensors with the most notable example being OxyR, which following oxidation, transcriptionally upregulates genes associated with the oxidative stress response (Zheng et al. 1998). In the case of OxyR, activation/inactivation occurs through a reversible disulfide bond and does not involve iron or divalent metals. While this could also be the case with replication, the requirement for manganese to recover would argue that the sensor in this case is a metal-containing enzyme (Djaman et al. 2004; Anjem and Imlay 2012; Imlay 2014). Many mononuclear iron enzymes are inactivated by H₂O₂ treatment, several of which can be reactivated following re-metallation with manganese, which is insensitive to oxidation, to restore function (Anjem and Imlay 2012; Imlay 2014). The sensor inhibiting replication could be an essential component of the replisome itself or a secondary protein that upon oxidation, associates to inhibit replication. However, we use the term 'sensor' in this case in the broadest possible sense, since there are a large number of mechanistic possibilities one could envision that could regulate the inactivation and reactivation of the replisome following a strong oxidative challenge.

Funding

This study was supported by National Science Foundation grant MCB1916625.

Conflicts of interest: None declared.

Literature cited

- Ames BN. 1983. Dietary carcinogens and anticarcinogens. Oxygen radicals and degenerative diseases. *Science*. 221:1256–1264.
- Anjem A, Imlay JA. 2012. Mononuclear iron enzymes are primary targets of hydrogen peroxide stress. *J Biol Chem*. 287: 15544–15556.
- Anjem A, Varghese S, Imlay JA. 2009. Manganese import is a key element of the OxyR response to hydrogen peroxide in *Escherichia coli*. *Mol Microbiol*. 72:844–858.
- Asad NR, de Almeida CEB, Asad LMBO, Felzenszwalb I, Leitão AC. 1995. Fpg and UvrA proteins participate in the repair of DNA lesions induced by hydrogen peroxide in low iron level in *Escherichia coli*. *Biochimie*. 77:262–264.
- Bagg A, Kenyon CJ, Walker GC. 1981. Inducibility of a gene product required for UV and chemical mutagenesis in *Escherichia coli*. *Proc Natl Acad Sci U S A*. 78:5749–5753.
- Bailly V, Verly WG. 1987. *Escherichia coli* endonuclease III is not an endonuclease but a beta-elimination catalyst. *Biochem J*. 242: 565–572.
- Beckman RA, Mildvan AS, Loeb LA. 1985. On the fidelity of DNA replication: manganese mutagenesis *in vitro*. *Biochemistry*. 24: 5810–5817.
- Belle JJ, Casey A, Courcelle CT, Courcelle J. 2007. Inactivation of the DnaB helicase leads to the collapse and degradation of the replication fork: a comparison to UV-induced arrest. *JB*. 189: 5452–5462.
- Belousova EA, Maga G, Fan Y, Kubareva EA, Romanova EA, et al. 2010. DNA polymerases beta and lambda bypass thymine glycol in gapped DNA structures. *Biochemistry*. 49:4695–4704.
- Bhagwat M, Gerlt JA. 1996. 3'- and 5'-strand cleavage reactions catalyzed by the Fpg protein from *Escherichia coli* occur via successive beta- and delta-elimination mechanisms, respectively. *Biochemistry*. 35:659–665.

- Bichara M, Pinet I, Lambert IB, Fuchs RP. 2007. RecA-mediated excision repair: a novel mechanism for repairing DNA lesions at sites of arrested DNA synthesis. *Mol Microbiol.* 65:218–229.
- Blakely WF, Fuciarelli AF, Wegher BJ, Dizdaroglu M. 1990. Hydrogen peroxide-induced base damage in deoxyribonucleic acid. *Radiat Res.* 121:338–343.
- Caron PR, Kushner SR, Grossman L. 1985. Involvement of helicase II (*uvrD* gene product) and DNA polymerase I in excision mediated by the *uvrABC* protein complex. *Proc Natl Acad Sci U S A.* 82:4925–4929.
- Castellani A, Jagger J, Setlow RB. 1964. Overlap of photoreactivation and liquid holding recovery in *Escherichia coli* B. *Science.* 143:1170–1171.
- Chan GL, Doetsch PW, Haseltine WA. 1985. Cyclobutane pyrimidine dimers and (6-4) photoproducts block polymerization by DNA polymerase I. *Biochemistry.* 24:5723–5728.
- Chow KH, Courcelle J. 2004. RecO Acts with RecF and RecR to protect and maintain replication forks blocked by UV-induced DNA damage in *Escherichia coli*. *J Biol Chem.* 279:3492–3496.
- Clark JM, Beardsley GP. 1986. Thymine glycol lesions terminate chain elongation by DNA polymerase I *in vitro*. *Nucleic Acids Res.* 14:737–749.
- Cole JM, Acott JD, Courcelle CT, Courcelle J. 2018. Limited capacity or involvement of excision repair, double-strand breaks, or translesion synthesis for Psoralen cross-link repair in. *Genetics.* 210:99–112.
- Courcelle CT, Chow KH, Casey A, Courcelle J. 2006. Nascent DNA processing by RecJ favors lesion repair over translesion synthesis at arrested replication forks in *Escherichia coli*. *Proc Natl Acad Sci U S A.* 103:9154–9159.
- Courcelle J, Crowley DJ, Hanawalt PC. 1999. Recovery of DNA replication in UV-irradiated *Escherichia coli* requires both excision repair and *recF* protein function. *J Bacteriol.* 181:916–922.
- Courcelle J, Donaldson JR, Chow KH, Courcelle CT. 2003. DNA damage-induced replication fork regression and processing in *Escherichia coli*. *Science.* 299:1064–1067.
- Courcelle J, Ganesan AK, Hanawalt PC. 2001. Therefore, what are recombination proteins there for? *Bioessays.* 23:463–470.
- Courcelle J, Hanawalt PC. 1999. RecQ and RecJ process blocked replication forks prior to the resumption of replication in UV-irradiated *Escherichia coli*. *Mol Gen Genet.* 262:543–551.
- Courcelle J, Hanawalt PC. 2001. Participation of recombination proteins in rescue of arrested replication forks in UV-irradiated *Escherichia coli* need not involve recombination. *Proc Natl Acad Sci U S A.* 98:8196–8202.
- Courcelle J, Hanawalt PC. 2003. RecA-dependent recovery of arrested DNA replication forks. *Annu Rev Genet.* 37:611–646.
- Courcelle J, Carswell-Crumpton C, Hanawalt PC. 1997. *recF* and *recR* are required for the resumption of replication at DNA replication forks in *Escherichia coli*. *Proc Natl Acad Sci U S A.* 94:3714–3719.
- Cross CE, Halliwell B, Borish ET, Pryor WA, Ames BN, et al. 1987. Oxygen radicals and human disease. *Ann Intern Med.* 107:526–545.
- Crowley DJ, Courcelle J. 2002. Answering the call: coping with DNA damage at the most inopportune time. *J Biomed Biotechnol.* 2:66–74.
- Demple B, Halbrook J, Linn S. 1983. *Escherichia coli* xth mutants are hypersensitive to hydrogen peroxide. *J Bacteriol.* 153:1079–1082.
- Djaman O, Outten FW, Imlay JA. 2004. Repair of oxidized iron-sulfur clusters in *Escherichia coli*. *J Biol Chem.* 279:44590–44599.
- Donaldson JR, Courcelle CT, Courcelle J. 2004. RuvAB and RecG are not essential for the recovery of DNA synthesis following UV-induced DNA damage in *Escherichia coli*. *Genetics.* 166:1631–1640.
- Donaldson JR, Courcelle CT, Courcelle J. 2006. RuvABC is required to resolve holliday junctions that accumulate following replication on damaged templates in *Escherichia coli*. *J Biol Chem.* 281:28811–28821.
- Efrati E, Tocco G, Eritja R, Wilson SH, Goodman MF. 1999. “Action-at-a-distance” mutagenesis. 8-oxo-7, 8-dihydro-2'-deoxyguanosine causes base substitution errors at neighboring template sites when copied by DNA polymerase beta. *J Biol Chem.* 274:15920–15926.
- Evans J, Maccabee M, Hatahet Z, Courcelle J, Bockrath R, et al. 1993. Thymine ring saturation and fragmentation products: lesion bypass, misinsertion and implications for mutagenesis. *Mutat Res.* 299:147–156.
- Fischhaber PL, Gerlach VL, Feaver WJ, Hatahet Z, Wallace SS, et al. 2002. Human DNA polymerase kappa bypasses and extends beyond thymine glycols during translesion synthesis *in vitro*, preferentially incorporating correct nucleotides. *J Biol Chem.* 277:37604–37611.
- Frank EG, Woodgate R. 2007. Increased catalytic activity and altered fidelity of human DNA polymerase iota in the presence of manganese. *J Biol Chem.* 282:24689–24696.
- Fridovich I. 1978. The biology of oxygen radicals. *Science.* 201:875–880.
- Ganesan AK, Smith KC. 1969. Dark recovery processes in *Escherichia coli* irradiated with ultraviolet light. II. Effect of *uvr* genes on liquid holding recovery. *J Bacteriol.* 97:1129–1133.
- Gifford CM, Blaisdell JO, Wallace SS. 2000. Multiprobe RNase protection assay analysis of mRNA levels for the *Escherichia coli* oxidative DNA glycosylase genes under conditions of oxidative stress. *J Bacteriol.* 182:5416–5424.
- Goerlich O, Quillardet P, Hofnung M. 1989. Induction of the SOS response by hydrogen peroxide in various *Escherichia coli* mutants with altered protection against oxidative DNA damage. *J Bacteriol.* 171:6141–6147.
- Goodman MF, Keener S, Guidotti S, Branscomb EW. 1983. On the enzymatic basis for mutagenesis by manganese. *J Biol Chem.* 258:3469–3475.
- Hanawalt PC. 2001. Controlling the efficiency of excision repair. *Mutat Res.* 485:3–13.
- Hayes RC, Leclerc JE. 1986. Sequence dependence for bypass of thymine glycols in DNA by DNA polymerase I. *Nucleic Acids Res.* 14:1045–1061.
- Hill RF. 1958. A radiation-sensitive mutant of *Escherichia coli*. *Biochim Biophys Acta.* 30:636–637.
- Hogg M, Seki M, Wood RD, Doublé S, Wallace SS. 2011. Lesion bypass activity of DNA polymerase θ (POLQ) is an intrinsic property of the pol domain and depends on unique sequence inserts. *J Mol Biol.* 405:642–652.
- Horii Z, Clark AJ. 1973. Genetic analysis of the *recF* pathway to genetic recombination in *Escherichia coli* K12: isolation and characterization of mutants. *J Mol Biol.* 80:327–344.
- Howard-Flanders P, Theriot L. 1966. Mutants of *Escherichia coli* K-12 defective in DNA repair and in genetic recombination. *Genetics.* 53:1137–1150.
- Howard-Flanders P, Theriot L, Stedford JB. 1969. Some properties of excision-defective recombination-deficient mutants of *Escherichia coli* K-12. *J Bacteriol.* 97:1134–1141.
- Howard-Flanders P. 1975. Repair by genetic recombination in bacteria: overview. *Basic Life Sci.* 5A:265–274.
- Hutchinson F. 1985. Chemical changes induced in DNA by ionizing radiation. *Prog Nucl Acid Res Mol Biol.* 32:115–154.

- Hutfilz CR, Wang NE, Hoff CA, Lee JA, Hackert BJ, et al. 2019. Manganese is required for the rapid recovery of DNA synthesis following oxidative challenge in *Escherichia coli*. *J Bacteriol.* 201: 28121.
- Ide H, Kow YW, Wallace SS. 1985. Thymine glycols and urea residues in M13 DNA constitute replicative blocks *in vitro*. *Nucl Acids Res.* 13:8035–8052.
- Ide H, Petrullo LA, Hatahet Z, Wallace SS. 1991. Processing of DNA base damage by DNA polymerases. Dihydrothymine and beta-ureidoisobutyric acid as models for instructive and noninstructive lesions. *J Biol Chem.* 266:1469–1477.
- Imlay JA. 2003. Pathways of oxidative damage. *Annu Rev Microbiol.* 57:395–418.
- Imlay JA. 2014. The mismetallation of enzymes during oxidative stress. *J Biol Chem.* 289:28121–28128.
- Imlay JA, Linn S. 1986. Bimodal pattern of killing of DNA-repair-defective or anoxically grown *Escherichia coli* by hydrogen peroxide. *J Bacteriol.* 166:519–527.
- Imlay JA, Linn S. 1987. Mutagenesis and stress responses induced in *Escherichia coli* by hydrogen peroxide. *J Bacteriol.* 169:2967–2976.
- Jeiranian HA, Courcelle CT, Courcelle J. 2012. Inefficient replication reduces RecA-mediated repair of UV-damaged plasmids introduced into competent *Escherichia coli*. *Plasmid.* 68:113–124.
- Jeiranian HA, Schalow BJ, Courcelle CT, Courcelle J. 2013. Fate of the replisome following arrest by UV-induced DNA damage in *Escherichia coli*. *Proc Natl Acad Sci U S A.* 110:11421–11426.
- Jiang D, Hatahet Z, Melamede RJ, Kow YW, Wallace SS. 1997. Characterization of *Escherichia coli* endonuclease VIII. *J Biol Chem.* 272:32230–32239.
- Kehres DG, Janakiraman A, Schlauch JM, Maguire ME. 2002. Regulation of *Salmonella enterica* serovar Typhimurium *mntH* transcription by H₂O₂, Fe(2+), and Mn(2+). *JB.* 184:3151–3158.
- Kogoma T. 1997. Stable DNA replication: interplay between DNA replication, homologous recombination, and transcription. *Microbiol Mol Biol Rev.* 61:212–238.
- Kunkel TA, Loeb LA. 1979. On the fidelity of DNA replication. Effect of divalent metal ion activators and deoxyribose nucleoside triphosphate pools on *in vitro* mutagenesis. *J Biol Chem.* 254:5718–5725.
- Laschia MF, Wallace SS. 1988. Excision repair of thymine glycols, urea residues, and apurinic sites in *Escherichia coli*. *J Bacteriol.* 170: 3359–3366.
- Lloyd RG, Thomas A. 1983. On the nature of the RecBC and RecF pathways of conjugal recombination in *Escherichia coli*. *Mol Gen Genet.* 190:156–161.
- Maga G, Villani G, Crespan E, Wimmer U, Ferrari E, et al. 2007. 8-oxo-guanine bypass by human DNA polymerases in the presence of auxiliary proteins. *Nature.* 447:606–608.
- Makarova AV, Boldinova EO, Belousova EA, Lavrik OI. 2018. *In vitro* lesion bypass by human PrimPol. *DNA Repair.* 70:18–24.
- Martin JE, Waters LS, Storz G, Imlay JA. 2015. The *Escherichia coli* small protein MntS and exporter MntP optimize the intracellular concentration of manganese. *PLoS Genet.* 11: e1004977.
- Mcculloch SD, Kokoska RJ, Garg P, Burgers PM, Kunkel TA. 2009. The efficiency and fidelity of 8-oxo-guanine bypass by DNA polymerases delta and eta. *Nucleic Acids Res.* 37:2830–2840.
- Melamede RJ, Hatahet Z, Kow YW, Ide H, Wallace SS. 1994. Isolation and characterization of endonuclease VIII from *Escherichia coli*. *Biochemistry.* 33:1255–1264.
- Mellon I, Hanawalt PC. 1989. Induction of the *Escherichia coli* lactose operon selectively increases repair of its transcribed DNA strand. *Nature.* 342:95–98.
- Napolitano R, Janel-Bintz R, Wagner J, Fuchs RP. 2000. All three SOS-inducible DNA polymerases (Pol II, Pol IV and Pol V) are involved in induced mutagenesis. *EMBO J.* 19:6259–6265.
- Nathan C, Cunningham-Bussel A. 2013. Beyond oxidative stress: an immunologist's guide to reactive oxygen species. *Nat Rev Immunol.* 13:349–361.
- Pages V, Fuchs RP. 2003. Uncoupling of leading- and lagging-strand DNA replication during lesion bypass *in vivo*. *Science.* 300: 1300–1303.
- Purmal AA, Lampman GW, Bond JP, Hatahet Z, Wallace SS. 1998. Enzymatic processing of uracil glycol, a major oxidative product of DNA cytosine. *J Biol Chem.* 273:10026–10035.
- Purmal AA, Lampman GW, Kow YW, Wallace SS. 1994. The sequence context-dependent mispairing of 5-hydroxycytosine and 5-hydroxyuridine *in vitro*. *Ann NY Acad Sci.* 726:361–363.
- Rouet P, Essigmann JM. 1985. Possible role for thymine glycol in the selective inhibition of DNA synthesis on oxidized DNA templates. *Cancer Res.* 45:6113–6118.
- Rupp WD, Howard-Flanders P. 1968. Discontinuities in the DNA synthesized in an excision-defective strain of *Escherichia coli* following ultraviolet irradiation. *J Mol Biol.* 31:291–304.
- Rupp WD, Wilde CE, Reno DL, Howard-Flanders P. 1971. Exchanges between DNA strands in ultraviolet-irradiated *Escherichia coli*. *J Mol Biol.* 61:25–44.
- Saito Y, Uraki F, Nakajima S, Asaeda A, Ono K, et al. 1997. Characterization of endonuclease III (*nth*) and endonuclease VIII (*nei*) mutants of *Escherichia coli* K-12. *J Bacteriol.* 179:3783–3785.
- Schalow BJ, Courcelle CT, Courcelle J. 2011. *Escherichia coli* Fpg glycosylase is nonredundant and required for the rapid global repair of oxidized purine and pyrimidine damage *in vivo*. *J Mol Biol.* 410: 183–193.
- Setlow RB, Swenson PA, Carrier WL. 1963. Thymine dimers and inhibition of DNA synthesis by ultraviolet irradiation of cells. *Science.* 142:1464–1466.
- Shibutani S, Grollman AP. 1994. Miscoding during DNA synthesis on damaged DNA templates catalysed by mammalian cell extracts. *Cancer Lett.* 83:315–322.
- Singer B, Kusmirek JT. 1982. Chemical mutagenesis. *Annu Rev Biochem.* 51:655–693.
- Spek EJ, Wright TL, Stitt MS, Taghizadeh NR, Tannenbaum SR, et al. 2001. Recombinational repair is critical for survival of *Escherichia coli* exposed to nitric oxide. *J Bacteriol.* 183:131–138.
- Spivak G, Hanawalt PC. 1995. Determination of damage and repair in specific DNA sequences. *Methods.* 7:147–161.
- Takata K, Shimizu T, Iwai S, Wood RD. 2006. Human DNA polymerase N (POLN) is a low fidelity enzyme capable of error-free bypass of 5S-thymine glycol. *J Biol Chem.* 281:23445–23455.
- Tokarsky EJ, Wallenmeyer PC, Phi KK, Suo Z. 2017. Significant impact of divalent metal ions on the fidelity, sugar selectivity, and drug incorporation efficiency of human PrimPol. *DNA Repair.* 49: 51–59.
- Vaisman A, Ling H, Woodgate R, Yang W. 2005. Fidelity of Dpo4: effect of metal ions, nucleotide selection and pyrophosphorolysis. *EMBO J.* 24:2957–2967.
- Van Houten B. 1990. Nucleotide excision repair in *Escherichia coli*. *Microbiol Rev.* 54:18–51.
- Walker GC. 1984. Mutagenesis and inducible responses to deoxyribonucleic acid damage in *Escherichia coli*. *Microbiol Rev.* 48:60–93.
- Wallace SS. 1994. DNA damages processed by base excision repair: biological consequences. *Int J Radiat Biol.* 66:579–589.
- Wallace SS. 2002. Biological consequences of free radical-damaged DNA bases. *Free Radic Biol Med.* 33:1–14.

- Wendel BM, Courcelle CT, Courcelle J. 2014. Completion of DNA replication in *Escherichia coli*. *Proc Natl Acad Sci U S A*. 111:16454–16459.
- Willetts NS, Clark AJ. 1969. Characteristics of some multiply recombination-deficient strains of *Escherichia coli*. *J Bacteriol*. 100:231–239.
- Willetts NS, Mount DW. 1969. Genetic analysis of recombination-deficient mutants of *Escherichia coli* K-12 carrying *rec* mutations cotransducible with *thyA*. *J Bacteriol*. 100:923–934.
- Yoon JH, Bhatia G, Prakash S, Prakash L. 2010. Error-free replicative bypass of thymine glycol by the combined action of DNA polymerases kappa and zeta in human cells. *Proc Natl Acad Sci U S A*. 107:14116–14121.
- Youngs DA, Bernstein IA. 1973. Involvement of the *recB-recC* nuclease (exonuclease V) in the process of x-ray-induced deoxyribonucleic acid degradation in radiosensitive strains of *Escherichia coli* K-12. *J Bacteriol*. 113:901–906.
- Yu D, Ellis HM, Lee EC, Jenkins NA, Copeland NG, et al. 2000. An efficient recombination system for chromosome engineering in *Escherichia coli*. *Proc Natl Acad Sci U S A*. 97:5978–5983.
- Zheng M, Aslund F, Storz G. 1998. Activation of the OxyR transcription factor by reversible disulfide bond formation. *Science*. 279:1718–1721.

Communicating editor: N. Rhind