

Portland State University

PDXScholar

Psychology Faculty Publications and
Presentations

Psychology

8-2023

Describing Physical Activity Patterns of Truck Drivers Using Actigraphy

Bradley Wipfli

OHSU-PSU School of Public Health, bwipfli@pdx.edu

Sean P.M. Rice

OHSU-PSU School of Public Health

Ryan Olson

Portland State University

Kasey Ha

Harvard School of Dental Medicine and Boston Children's Hospital

Caitlyn F. Trullinger-Dwyer

Oregon Health & Science University

See next page for additional authors

Follow this and additional works at: https://pdxscholar.library.pdx.edu/psy_fac



Part of the [Medicine and Health Sciences Commons](#), and the [Psychology Commons](#)

Let us know how access to this document benefits you.

Citation Details

Wipfli, Brad, Sean PM Rice, Ryan Olson, Kasey Ha, Caitlyn Trullinger-Dwyer, and Todd Bodner. "Describing Physical Activity Patterns of Truck Drivers Using Actigraphy." *Safety and Health at Work* (2023).

This Article is brought to you for free and open access. It has been accepted for inclusion in Psychology Faculty Publications and Presentations by an authorized administrator of PDXScholar. Please contact us if we can make this document more accessible: pdxscholar@pdx.edu.

Authors

Bradley Wipfli, Sean P.M. Rice, Ryan Olson, Kasey Ha, Caitlyn F. Trullinger-Dwyer, and Todd Bodner



Contents lists available at ScienceDirect

Safety and Health at Work

journal homepage: www.e-shaw.net

Short Communication

Describing Physical Activity Patterns of Truck Drivers Using Actigraphy

Brad Wipfli^{1,2,*}, Sean P.M. Rice^{1,2}, Ryan Olson^{1,2,3}, Kasey Ha⁴, Caitlyn Trullinger-Dwyer², Todd Bodner³¹ School of Public Health, Oregon Health & Science University-Portland State University, Portland, OR, USA² Oregon Institute of Occupational Health Sciences, Oregon Health & Science University, Portland, OR, USA³ Department of Psychology, Portland State University, Portland, OR, USA⁴ Harvard School of Dental Medicine and Boston Children's Hospital, Boston, MA, USA

ARTICLE INFO

Article history:

Received 20 March 2023

Received in revised form

10 August 2023

Accepted 14 August 2023

Available online 19 August 2023

Keywords:

accelerometer
commercial driver
exercise
objective measurement

ABSTRACT

Background: Truck driving is a highly sedentary occupation that places workers at risk for chronic health conditions, such as obesity and high blood pressure. The primary purpose of this study was to objectively describe truck drivers' typical physical activity (PA) patterns.**Methods:** We used ~7–10-day baseline PA actigraphy data samples from drivers in the Safety & Health Involvement For Truckers (SHIFT) study ($n = 394$). Driver PA patterns (e.g., average number of ≥ 10 minute Freedson bouts per week, time in bouts, and common days/times for PA) were summarized with descriptive analyses. We also compared objective accelerometer data to self-reports.**Results:** Drivers' weekly PA averaged 14.4 minutes ($SD = 37.0$), and most PA occurred between 5–6 pm on Tuesdays and Wednesdays. Drivers overestimated self-reported weekly exercise by over 60 min/week compared to accelerometer data.**Conclusion:** Our results suggest that objective PA assessment may be warranted over self-report when possible, and timing may be key in future PA intervention work with truck drivers.© 2023 The Authors. Published by Elsevier B.V. on behalf of Occupational Safety and Health Research Institute, Korea Occupational Safety and Health Agency. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Obesity rates among truck drivers are over 30% higher than that of the general working population [1]. Numerous structural and systemic factors contribute to truck driver obesity, including long and sedentary driving shifts, food swamps at truck stops, variable schedules, and limited off-duty time and safe environments for physical activity (PA) [2]. Interventions to improve drivers' working conditions, schedules, and health are important, including those that aim to increase PA. However, little is known about truck drivers' actual PA levels and patterns, and opportunities to create or protect time for drivers to maintain or increase PA. Studies of drivers' PA are scarce [1,3,4] and tend to rely on self-report data that often result in overestimates of PA (correlations between objective and subjective measurements range between -0.71 and + 0.96) [5]. There is a need for studies that objectively measure truck drivers' PA to guide and evaluate future health interventions [6].

Actigraphy is a common and reliable method for objectively measuring PA [7], but it is rarely used in research with commercial

drivers because vehicular movement and vibrations in trucks can artificially inflate actigraphy estimates of energy expenditure [8–10]. However, focusing on bouts of PA ≥ 10 minutes in duration—instead of the sum of energy expenditure estimates—decreases the likelihood of artificial inflation due to vehicular movement and vibration. Prior research has shown that this approach does not produce false bouts during workers' driving periods and thus opens the door for conservative and objective measurement of PA among truck drivers [10]. By analyzing patterns of PA bouts in truck drivers and the context in which PA bouts occur, we can better understand how and when to intervene to increase PA and inform employers' approaches to managing drivers' route assignments and schedules. To our knowledge, only five previous studies have used actigraphy to measure PA among truck drivers [11–15], and these studies failed to account for vehicular movement and vibrations and had small sample sizes. As such, the purpose of this study was to apply a bout-based analysis approach

Brad Wipfli: <https://orcid.org/0000-0001-7261-0551>; Sean P.M. Rice: <https://orcid.org/0000-0001-7712-4667>; Ryan Olson: <https://orcid.org/0000-0002-2751-4402>; Kasey Ha: <https://orcid.org/0009-0009-4153-6921>; Caitlyn Trullinger-Dwyer: <https://orcid.org/0009-0004-9153-6966>; Todd Bodner: <https://orcid.org/0000-0002-7001-274X>

* Corresponding author. OHSU-PSU School of Public Health, 1810 SW 5th Ave, Suite 510, Portland, OR 97201, USA.

E-mail address: bwipfli@pdx.edu (B. Wipfli).

with PA actigraphy data collected from a large sample of US truck drivers in order to describe their typical patterns of PA during a work week. We aimed to describe the days and times drivers usually engaged in PA, the number and length of activity bouts per week, and contextual characteristics associated with PA. We also compared objective measures of PA from actigraphy data to drivers' PA self-report data.

1. Method

1.1. Participants

This study focuses on baseline data collected from truck drivers ($N = 452$) who participated in the Safety and Health Involvement for Truck Drivers (SHIFT) randomized controlled trial conducted in the US [3]. All participants provided informed consent prior to participation, and all methods were approved by the human subjects institutional review board at Oregon Health & Science University. Participants included truck drivers at five companies in the Western USA. Descriptions of participant recruitment, demographics, and work and health characteristics of the full sample have been previously published [3]. Table 1 describes characteristics of the participants included in this study. Participants were mostly male (87.0%) with an average age of 48.4 (SD = 11.0). Average tenure as a truck driver was 11.9 years (SD = 10.3), and current company tenure was 5.0 years (SD = 5.7).

1.2. Devices and implementation procedure

PA was measured with a tri-axial accelerometer and inclinometer (ActiGraph GT3X+, ActiGraph LLC, Pensacola, FL). Using Actilife software, GT3X+ devices were initialized with participant demographics, set with a sampling rate of 30 Hz, and set to begin recording 2 hours after each participant had completed an in-person data collection visit (survey and objective health measures). Participants were provided with an elastic belt to wear the device on their non-dominant hip [16]. Researchers guided participants as they put the belt and ActiGraph on, and provided verbal and written instructions for wearing the device during the sampling period. Participants were asked to only remove device when changing clothes or during activities where the device could get

wet. They were also asked to wear the devices for 7–10 consecutive days, including work and non-work days, though some participants wore them for different lengths. Participants returned devices to researchers using a prepaid mailing envelope. All devices were serviced and maintained according to the manufacturer's recommendations during the study period.

1.3. Actigraphy measures and analyses

Wear time was validated with a modified algorithm [17] based on a study of PA in a highly sedentary population [18], changing the requirement from 90 consecutive minutes of zero activity counts to 120 consecutive minutes of zero activity counts to be classified as non-wear time. Using this method, a sample day was considered valid if wear time compliance was equal to or exceeded 8 hours per day. Participants were required to have three or more valid days to be included in the analysis. Due to missing data or insufficient wear time, 58 drivers were excluded from analyses, leaving a total sample of $n = 394$. Activity bouts were calculated using the Freedson Adult (2011) algorithm [16], with an activity threshold of 1952 counts/minute. Only bouts at least 10 minutes long, inclusive of a 2-minute drop period, were analyzed.

Electronic logbook data were obtained for a subsample of drivers ($n = 46$), which indicated driving and non-driving periods. We compared these data to records of activity bouts to confirm that bouts of PA observed in actigraphy data were not falsely generated by truck vibrations during driving periods. In this sub-sample, 14 of the 46 drivers generated a total of 79 activity bouts, and all bouts occurred during non-driving times.

1.4. Additional measures and analyses

In the SHIFT trial, participants completed a range of survey measures, including demographics and self-reported PA. Directly measured health factors included blood pressure (Omron HEM-907XL, Kyoto, Japan) and body composition (Tanita TBF-310GS scale, Tanita Corporation, Tokyo, Japan). Self-reported PA was measured using the Healthy Physical Activity Questionnaire [19], and days per week with at least 30 minutes of moderate and vigorous PA in the past month were summed. Average PA minutes per week was conservatively extrapolated by multiplying this sum

Table 1
Participant characteristics

Variables	No freedson bouts ($N = 266$)	≥ 1 freedson bouts ($N = 128$)	Total sample ($N = 394$)	p
	Mean (SD) or n (%)	Mean (SD) or n (%)	Mean (SD) or n (%)	
Age (years)	48.8 (11.4)	47.5 (10.2)	48.4 (11.0)	0.30
Sex (Female)	37 (14.0%)	14 (10.9%)	51 (13.0%)	0.40
Weight (lb)	258.0 (64.2)	237.2 (46.0)	251.2 (59.6)	<0.01
Tenure (years)				
As truck driver	12.9 (10.9)	9.8 (8.5)	11.9 (10.3)	<0.01
At current company	4.7 (5.5)	5.4 (5.9)	5.0 (5.7)	0.29
Employers in past 3 years	1.3 (1.3)	1.0 (1.0)	1.2 (1.2)	0.03
Driving hours per week*	59.4 (11.8)	59.6 (10.7)	59.5 (11.5)	0.79
Days away from home	3.8 (1.5)	3.8 (1.5)	3.8 (1.5)	0.69
Company				
1	32 (12.0%)	13 (10.2%)	45 (11.4%)	
2	48 (18.0%)	17 (13.3%)	65 (16.5%)	
3	99 (37.2%)	40 (31.3%)	139 (35.3%)	
4	75 (28.2%)	57 (44.5%)	132 (33.5%)	
5	12 (4.5%)	1 (0.8%)	13 (3.3%)	
Self-reported physical activity				
Moderate physical activity (0–7 days)	1.8 (1.8)	1.9 (1.8)	1.8 (1.8)	0.58
Vigorous physical activity (0–7 days)	0.9 (1.5)	1.1 (1.6)	0.9 (1.5)	0.28
Extrapolated physical activity time per week (minutes)	79.0 (86.6)	87.5 (84.5)	81.8 (85.9)	0.36

* Variable compared using Mann–Whitney U test because of substantial skew and kurtosis. All other continuous variables compared using independent samples t -test.

by 30 (i.e., the minimum amount of PA participants could have engaged in according to the prompt). Descriptive statistics (e.g., means and standard deviations) of participants' demographic and work characteristics were computed for the whole sample and compared between those with and without activity bouts, using independent samples t-test for continuous variables and logistic regression for categorical variables (for non-normal continuous variables Mann–Whitney U test was used).

Within the group of drivers who had activity bouts, associations between participant characteristics and bouts, average time in bouts, and average PA time per week were assessed via Spearman's rho (ρ), and patterns of PA (e.g., day-of-week and time-of-day) were descriptively analyzed and visualized. Finally, PA self-reports were compared to actigraphy data. Time-of-day of PA was determined by the hour. For example, if some had a bout that lasted between 13:25 and 13:35, then that bout and 10 minutes of activity would be classified as occurring at 1 pm. If, however, a bout occurred between multiple-hour cutoffs (e.g., a bout that begins at 8:50 and ends at 9:05), then that bout was classified as the hour in which the majority of PA occurred (e.g., 8 am). If the bout time is equally spaced between hour cutoffs, (e.g., 8:55–9:05), then we coded that bout as the later hour. The minutes of activity, however, were only classified as occurring in each respective hour (e.g., 5 minutes at 8 am and 5 minutes at 9 am for a bout between 8:55–9:05). Self-reported PA was compared to objective actigraphy data via Spearman's rho and Wilcoxon signed rank test. An alpha level of 0.05 was used to indicate significance from statistical tests; however, interpretations of such results are limited to describing potentially meaningful associations rather than making causal inferences.

2. Results

The sample ($n = 394$) produced a total 389 Freedson bouts over the monitoring period (number of valid days ranged from 3 to 24 days; mean = 10.1, SD = 2.8). The majority of drivers ($n = 266$) did not complete any PA bouts lasting 10 minutes or more. The drivers ($n = 128$) who produced at least one bout of PA completed an average of 3.0 bouts (SD = 3.2) per person, and the mean bout duration was 19.2 minutes (SD = 11.5). A descriptive summary of driver characteristics for those with and without bouts is provided in Table 1. Some identified differences were that drivers who completed at least one bout tended to weigh less (mean difference [MD] = -20.8 lb, $p < 0.01$), have shorter career tenure (MD = -3.1 years, $p < 0.01$), report fewer employers in the past three years (MD = -0.3, $p = 0.03$), and have a lower computed likelihood of cardiovascular disease [20,21] (MD = -2.3 total points, $p = 0.01$). Overall, drivers averaged 14.4 minutes of PA per week (SD = 37.0).

Within the activity bout group, this average increases to 44.4 minutes (SD = 53.8).

Within the activity bout group, we computed correlations between activity variables and driver characteristics (see Table 2). Weight was negatively associated with Freedson bouts per week, minutes per bout, and minutes per week of PA. Tenure was positively associated with minutes per bout, but not bouts or minutes per week. Similarly, days away from home were negatively associated with minutes per bout, but had no relationship to bouts or minutes per week. Company tenure was positively associated with minutes in bout and minutes per week, but not bouts per week. Finally, though driving hours per week was only significantly negatively correlated with minutes per week, the correlation size ($r = -0.19$) was similar to that of bouts per week ($r = -0.14$) and minutes per bout ($r = -0.17$).

See Figs. 1 and 2 for visualizations of PA by time-of-day and day-of-week, respectively. Most PA bouts occurred in the 5 pm (total bouts = 40; 10.3%) and 6 pm (total bouts = 38; 9.8%) periods. Further, the majority of bouts occurred mid-week on Tuesdays (total bouts = 70; 18.0%), Wednesdays (total bouts = 61; 15.7%), and Thursdays (total bouts = 64; 16.5%). Patterns in total PA minutes mirrored data on the number of bouts, with a total of 694 (9.6%), 716 (9.9%), and 665 (9.2%) minutes occurring in the 5 pm, 6 pm, and 7 pm hour periods, respectively. Similarly, 1,319 (17.7%); 1,215 (16.3%); and 1,149 (15.4%) total minutes of PA were recorded on Tuesday, Wednesday, and Thursday, respectively.

According to self-report for the past month, drivers engaged in at least 30 minutes of moderate PA on about 1.8 days per week (SD = 1.8) and vigorous activity on about 0.9 days per week (SD = 1.5). After an extrapolation to standardize these self-reports relative to actigraphy-measured bouts, participants averaged 81.8 minutes (SD = 85.6) of self-reported PA per week. Thus, drivers overestimated their PA by 67.3 minutes per week (SD = 90.5, $p < 0.01$) compared to their more objective actigraphy data. Additionally, self-report and objective measures of total time in PA were not correlated ($\rho = 0.08$, $p = 0.11$). When excluding drivers with 0 bouts, drivers still overestimated their PA by about 43.2 minutes per week (SD = 93.9), $p < 0.01$. In this subsample with bouts, the rho correlation between self-reported and objective PA minutes per week was non-significant ($\rho = 0.13$, $p = 0.16$).

3. Discussion

To our knowledge, this study represents the first actigraphy measurement of truck driver PA patterns using a method that excludes potential false activity generated by truck vibrations. Results from this sample describe truck drivers as a highly sedentary


Table 2
Correlations between driver characteristics and objective physical activity

Variable	1	2	3	4	5	6	7	8	9	10
1. Bouts per week										
2. Min per bout	0.38*									
3. Min per week	0.91*	0.70*								
4. Age	0.11	0.17	0.14							
5. Sex	0.08	-0.11	0.02	-0.11						
6. Weight	-0.29*	-0.18*	-0.28*	-0.06	-0.09					
7. Tenure	0.09	0.20*	0.15	0.48*	0.02	0.03				
8. C tenure	0.14	0.30*	0.23*	0.41*	0.01	-0.01	0.72*			
9. # Employers	0.12	-0.13	0.05	-0.35*	0.11	0.01	-0.07	-0.38*		
10. Driving hours per week	-0.14	-0.17	-0.19*	0.05	-0.05	-0.04	-0.06	-0.19*	0.01	
11. Days away from home	-0.01	-0.18*	-0.09	0.01	0.02	-0.01	-0.11	-0.18*	-0.05	0.50*

Note. All values reflect Spearman's rho. Correlations were computed with the sample that had at least one Freedson bout. Min = minutes. C Tenure = company tenure. *Significant at $p < 0.05$.

A)

Activity Bouts at Time-of-Day



B)

Activity Minutes at Time-of-Day




Fig. 1. Physical activity and time-of-day.

population in need of improvements to working conditions, scheduling support from employers, and access to additional health promotion programs. In this study, two-thirds of the sample did not engage in any PA bouts of 10 minutes or greater during a work week, and the average total minutes of PA per week was less than 30 minutes. When drivers engaged in PA, they tended to do so in the early evening (around 5–6 pm) on mid-weekdays (Tuesday and Wednesday). Results also indicated that drivers substantially overestimated their level of PA in a self-report measure.


Future intervention work may benefit from these findings in a few ways. First, given that PA on driving days is more likely to occur in the evening after a driving period, employers and supervisors (dispatchers) should work to manage driver schedules in ways that protect early mid-week evening off-duty time, as well as access to truck or rest stops that provide safe places to walk or exercise. On this theme, it is possible that with progressive days on duty, or when work hours extend to late in the evening, driver fatigue increases and motivation for engaging PA reduces. While we did not collect indicators of consecutive days on duty in this study, it is possible that the predominance of early mid-week PA and limited presence of later week and weekend PA could indicate progressive fatigue, recovery on weekends, and then higher motivation to exercise after off-duty recovery time at the beginning of a work week. In trucking a typical work week many not start Monday and end

Friday, but this observed pattern and possible relationship between consecutive work days and PA may warrant future investigation.

Observed patterns in PA may also inform health programs or systems, which could utilize ecologically timed reminders (or other motivational tools) to prime drivers during their usual peak PA times (i.e., Tuesdays and Wednesdays between 5–6 pm). Increasing motivational prompts in temporal proximity to normal activity bout opportunities may make it more likely for drivers to engage in PA and may make them more motivated to exercise for longer. The more tailored an intervention is to one's motivational readiness, the more exercise they are likely to engage in [22]. Interventions involving just-in-time adaptive prompts to engage in PA were reviewed in 2019 [23]. Most of the studies reviewed ($k = 19$) were self-described as intervention pilots, and thus likely underpowered. Outcomes across the studies were mixed, yet promising. To illustrate, one of two workplace interventions reviewed monitored participants' step counts and locations using wearable technology and then provided "context-aware" coaching via smartphone to encourage PA at opportune moments [24]. In the active intervention phase, participants in the context-aware coaching group showed a larger increase in daily step counts (about 600 steps greater) compared to control.

Further, day-of-week patterns in PA for drivers also suggest there is a need for addressing and encouraging PA during weekend

A)



B)




Fig. 2. Physical activity and day-of-week.

or restart periods, perhaps by involving friends and family members in health program educational and motivational components. In total, the patterns observed in truck drivers' PA may serve to guide employers' efforts to organize and schedule work to support driver well-being, as well as to improve the tailoring and efficacy of future PA-promoting educational and motivational programs.

3.1. Strengths and limitations

Strengths of the study include the objective measurement of PA, utilization of a method that minimized potential for false PA from vehicular movement and vibration, a large sample of workers from a priority occupation for health interventions, and the novelty of the PA pattern assessment with this occupational group. There are often large discrepancies between self-report and objective measures of PA in general population samples [5], and our own findings support this divergence in a sample of truck drivers as well. Typically, and as was found in our results, individuals will overestimate their usual PA. Using actigraphy to measure PA objectively presents a more accurate picture of how much, how often, and when individuals are active. Truck drivers are at particularly high risk for health issues as a result of the characteristics of their work, and the results of this study provide an important incremental step forward for guiding future interventions at trucking companies that benefit driver health and wellbeing.

To place findings in context, the current study provides measurements based on a substantially larger sample size and contributes additional information about the timing of PA compared to other relatively recent accelerometer-based studies of commercial

drivers. In a small pilot study of team truck drivers ($n = 8$) using the same actigraph and bout-based analysis as the current study, Olson et al. found drivers at baseline averaged 4.40 ($SD = 7.36$) minutes per week in moderate-intensity PA bouts [14]. Gilson et al. used the same actigraph devices as the current study, but worn on the wrist, to cross-sectionally measure driver PA [13]. Drivers in this sample ($n = 21$) averaged less than one 10+ minute bout of PA per day, with fewer bouts on non-workdays compared to workdays. Verela Mato et al. [11] reported moderate to vigorous stepping behavior of 0.1 ($SD = 0.1$) hours per day among 43 truck drivers using the activePAL3 monitor placed on the thigh. Gilson et al. used a wrist-worn GENEActiv accelerometer and reported that drivers ($n = 44$) spent 5% to 9% of their on-duty and off-duty workday time, respectively, being physically active (walking or running) [12]. Additionally, Clemes et al. reported a significant post-intervention difference in daily step count as measured by wrist-worn Fitbit devices, resulting from a decrease in step count in the control group and maintenance of step count in the intervention group [15]. However, these studies did not address how their particular devices may have been affected by large truck vibrations or driving tasks (e.g., operating steering wheel and shifting gears).

The dissimilarity of measurement anchors in our comparison of objective and subjective PA is one limitation of this study. The nature of the survey scale required that we make an extrapolation from two self-report items in order to transform answers into minutes per week of PA. Those two items asked participants to report days per week of at least 30 minutes of moderate and vigorous PA in the last month. It may be more valid for future research to use a self-report measure of PA that is directly anchored in minutes of

activity per week (e.g., the International Physical Activity Questionnaire [IPAQ] [25]) for comparisons to objective data. Additionally, we asked participants about their general activity in the past month and compared it to a single week of objective data (the self-report period also came before the actigraphy data collection period). Activity levels may vary week-to-week, so future research may benefit from using the same time period for subjective-objective comparisons (e.g., IPAQ uses “the past 7 days” as the time anchor). Similarly, another limitation was its focus on a specific context and population which is long-haul truck drivers in the USA. The patterns identified may not be generalizable for different kinds of truck drivers (e.g., short-haul) or drivers from different countries. For example, federal driving regulations in the USA allow up to 11 hours on-duty per day (for property-carrying drivers), requiring 30 minutes of rest every 8 hours of driving [26]. As such, evenings as a time where many drivers will engage in most of their PA makes sense (i.e., after a full day of driving). In the European Union, in contrast, there is a limit of 9 hours per day (and 56 hours per week), with requirements of 45 rest minutes every 4.5 hours of driving [27]. European drivers may therefore have more opportunities to engage in PA throughout the work day. In general, there are many factors that may impact the generalizability of the present study’s results and conclusions (e.g., geography, climate, local and federal laws). Future research should investigate these potential impacts in more depth. A final limitation of the study is our inability to note off-duty and on-duty days, as well as consecutive days on duty during the sampling periods, for the full sample of drivers. Future research may benefit from collecting electronic logbook data for every participant, or from utilizing daily diary survey methods during the sampling period, to investigate deeper questions about PA as it relates to daily work hours, consecutive work days, off-duty PA opportunities, and work and non-work days.

3.2. Conclusions

Overall, our results suggest that truck drivers tend to engage in PA infrequently but do have certain patterns to their PA (e.g., during evenings) that may be leveraged to guide future intervention work. Results also strongly encourage the use of actigraphy instead of, or as a supplement to, driver self-reports of PA. Driver self-reports greatly overestimated their PA levels. In the absence of analysis algorithms that strongly account for or exclude non-human vibrations, future research on PA with truck drivers should employ activity bouts as an outcome variable to prevent attributing truck-generated activity to drivers. Patterns in driver PA, as well as exploratory analyses related to driver sleep quality, encourage future research on the relationship between driver work exposures, fatigue, sleep quality, and PA. Such work could be facilitated by actigraphy studies employing both hip and wrist-worn devices to measure both PA and sleep in the same study. The priority for future interventions should be placed upon improving drivers’ working conditions and creating or protecting time and space for them to be physically active. Such work can then be supplemented with tailored educational and motivational programs to support increases in PA and improved overall driver health.

Data statement

Data will be provided upon request from qualified researchers.

Conflicts of interest

The authors declare they have no conflicts of interest.

Acknowledgments

The research was supported with a grant from the National Heart, Lung, and Blood Institute (R01HL1054950). This study is registered as a clinical trial (NCT02105571).

<https://clinicaltrials.gov/ct2/show/NCT02105571?term=SHIFT+truck&rank=1>. The authors would like to thank Kevin Bransford, Sydney Running, Melanie Koren, Katrina Bettencourt, Jessica Sanata, Louis Moore, Kristy Luther, Noal Clemons, Emma Robson, Annie Buckmaster, Annie Cannon, Rossmary Vasquez, Diane Elliot, Nancy Perrin, and Kent Anger for their contributions to the study.

References

- [1] Sieber WK, Robinson CF, Birdsey J, Chen G-X, Hitchcock EM, Lincoln JE, Nakata A, Sweeney MH. Obesity and other risk factors: the national survey of U.S. long-haul truck driver health and injury. *Am J Ind Med* 2014;57:615–26.
- [2] Lincoln JE, Birdsey J, Sieber WK, Chen G-X, Hitchcock EM, Nakata A, Robinson CF. A pilot study of healthy living options at 16 truck stops across the United States. *Am J Health Promot* 2018;32:546–53.
- [3] Olson R, Wipfli B, Thompson SV, Elliot DL, Anger WK, Bodner T, Hammer LB, Perrin NA. Weight control intervention for truck drivers the SHIFT randomized controlled trial, United States. *Am J Public Health* 2016;106:1698–706.
- [4] Birdsey J, Sieber WK, Chen G-X, Hitchcock EM, Lincoln JE, Nakata A, Robinson CF, Sweeney MH. National survey of US long-haul truck driver health and injury: health behaviors. *J Occup Environ Med* 2015;57:210–6.
- [5] Prince SA, Adamo KB, Hamel ME, Hardt J, Gorber SC, Tremblay M. A comparison of direct versus self-report measures for assessing physical activity in adults: a systematic review. *Int J Behav Nutr Phys Act* 5;56:1-24.
- [6] Pritchard EK, Kim HC, Nguyen N, van Vreden C, Xia T, Iles R. The effect of weight loss interventions in truck drivers: systematic review. *PLoS One* 2022;17:e0262893.
- [7] Freedson PS, Melanson E, Sirard J. Calibration of the computer science and applications, Inc. accelerometer. *Med Sci Sports Exerc* 1998;30:777–81.
- [8] Cohen MD, Cutaia M, Brehm R, Brutus V, Pike VC, Lewendowski D. Detecting motor vehicle travel in accelerometer data. *COPD: J Chronic Obstr Pulm Dis* 2012;9:102–10.
- [9] Steele BG, Holt L, Belza B, Ferris S, Lakshminaryan S, Buchner DM. Quantitating physical activity in COPD using a triaxial accelerometer. *Chest* 2000;117:1359–67.
- [10] Wipfli B, Olson R, Uba U, Moore L, Clemons N, Thompson S, Bransford K. Physical activity actigraphy among truck drivers: accounting for seat vibrations. *Med Sci Sports Exerc* 2014;46:792–3.
- [11] Varela Mato V, Caddick N, King JA, Johnson V, Edwardson C, Yates T, Stensel DJ, Daly H, Nimmo MA, Cledes SA. The impact of a novel structured health intervention for truckers (SHIFT) on physical activity and cardiometabolic risk factors. *J Occup Environ Med* 2018;60:368–76.
- [12] Gilson ND, Pavey TG, Wright ORL, Vandelanotte C, Duncan MJ, Gomersall S, Trost SG, Brown WJ. The impact of an m-health financial incentives program on the physical activity and diet of Australian truck drivers. *BMC Public Health* 2017;17:1–11.
- [13] Gilson ND, Mielke GI, Coombes JS, Feter N, Smith E, Duncan MJ, Wallis G, Brown WJ. VO2peak and 24-hour sleep, sedentary behavior, and physical activity in Australian truck drivers. *Scand J Med Sci Sports* 2021;31:1574–8.
- [14] Olson R, Johnson P, Shea SA, Marino M, Rimby J, Womak K, Wang F, Springer R, Donovan C, Rice SPM. Advancing the safety, health, and well-being of commercial driving teams who sleep in moving semi-trucks: the techArest pilot study. *J Occup Environ Med* 2020;62:1082–96.
- [15] Clemes SA, Varela-Mato V, Bodicoat DH, Brookes CL, Chen Y-L, Cox E, Edwardson CL, Gray LJ, Guest A, Johnson V, Munir F, Paine NJ, Richardson G, Ruettger K, Sayyah M, Sherry A, Di Paola AS, Troughton J, Walker S, Yates T, King J. A multicomponent structured health behaviour intervention to improve physical activity in long-distance HGV drivers: the SHIFT cluster rct. *Public Health Res* 2022;10:1–210.
- [16] Sasaki JE, John D, Freedson PS. Validation and comparison of ActiGraph activity monitors. *J Sci Med Sport* 2011;14:411–6.
- [17] Choi L, Liu Z, Matthews CE, Buchowski MS. Validation of accelerometer wear and nonwear time classification algorithm. *Med Sci Sports Exerc* 2011;43:357–64.
- [18] Hutto B, Howard VJ, Blair SN, Colabianchi N, Vena JE, Rhodes D, Hooker SP. Identifying accelerometer nonwear and wear time in older adults. *Int J Behav Nutr Phys Act* 2013;10:1–8.
- [19] Elliot DL, Goldberg L, Kuehl KS, Moe EL, Breger RKR, Pickering MA. The PHLAME (promoting healthy lifestyles: alternative models’ effects) firefighter study: outcomes of two models of behavior change. *J Occup Environ Med* 2007;49:204–13.
- [20] Goff DC, Lloyd-Jones DM, Bennett G, Coady S, D’Agostino RB, Gibbons R, Greenland P, Lackland DT, Levy D, O’Donnell CJ, Robinson JG, Schwartz JS,

- Shero ST, Smith SC, Sorlie P, Stone NJ, Wilson PWF, Jordan HS, Nevo L, Wnek J, Anderson JL, Halperin JL, Albert NM, Bozkurt B, Brindis RG, Curtis LH, DeMets D, Hochman JS, Kovacs RJ, Ohman EM, Pressler SJ, Sellke FW, Shen W-K, Smith SC, Tomaselli GF. ACC/AHA guideline on the assessment of cardiovascular risk. 2013. *Circulation* 2014;129:S49–73.
- [21] Framingham Heart Study. Hard coronary artery disease (10-year risk); 2001 [cited 2023 Jan 23]. Available from: <https://www.framinghamheartstudy.org/fhs-risk-functions/hard-coronary-heart-disease-10-year-risk/>.
- [22] Marcus BH, Emmons KM, Simkin-Silverman LR, Linnan LA, Taylor ER, Bock BC, Roberts MB, Rossi JS, Abrams DB. Evaluation of motivationally tailored vs. standard self-help physical activity interventions at the workplace. *Am J Health Promot* 1998;12:246–53.
- [23] Hardeman W, Houghton J, Lane K, Jones A, Naughton F. A systematic review of just-in-time adaptive interventions (JITAI) to promote physical activity. *Int J Behav Nutr Phys Act* 2019;16:31.
- [24] Van Dantzig S, Bulut M, Krans M, Van der Lans A, De Ruyter B. Enhancing physical activity through context-aware coaching. In: Proceedings of the 12th EAI international conference on pervasive computing technologies for healthcare. New York (NY): Association for Computing Machinery; 2018. p. 187–90. 2018 May 21–24; New York, NY.
- [25] Lee PH, Macfarlane DJ, Lam TH, Stewart SM. Validity of the international physical activity Questionnaire short form (IPAQ-SF): a systematic review. *Int J Behav Nutr Phys Act* 2011;8:1–11.
- [26] Federal Motor Carrier Safety Administration. Summary of hours of service regulations [Internet]. United States Department of Transportation. [updated 2022 March 28; cited 2023 Aug 9]. Available from: <https://www.fmcsa.dot.gov/regulations/hours-service/summary-hours-service-regulations>.
- [27] Your Europe. EU rules for working in road transport. [updated 2023 May 1; cited 2023 Aug 9]. Available from: https://europa.eu/youreurope/citizens/work/work-abroad/rules-working-road-transport/index_en.htm.