

Portland State University

PDXScholar

---

Chemistry Faculty Publications and  
Presentations

Chemistry

---

4-24-2024

# Chemical and Physiological Interactions Between E-Liquid Constituents: Cause for Concern?

Robert M. Strongin

*Portland State University, strongin@pdx.edu*

Eva Sharma

*Westat Inc.*

Hanno C. Erythropel

*Yale University*

Nada O F Kassem

*San Diego State University Research Foundation*

Alexandra Noël

*Louisiana State University*

*See next page for additional authors*

Follow this and additional works at: [https://pdxscholar.library.pdx.edu/chem\\_fac](https://pdxscholar.library.pdx.edu/chem_fac)

 Part of the [Chemistry Commons](#)

Let us know how access to this document benefits you.

---

## Citation Details

Strongin, R. M., Sharma, E., Erythropel, H. C., Kassem, N. O. F., Noël, A., Peyton, D. H., & Rahman, I. (2024). Chemical and physiological interactions between e-liquid constituents: cause for concern? *Tobacco Control*, tc-2023-058546.

This Article is brought to you for free and open access. It has been accepted for inclusion in Chemistry Faculty Publications and Presentations by an authorized administrator of PDXScholar. Please contact us if we can make this document more accessible: [pdxscholar@pdx.edu](mailto:pdxscholar@pdx.edu).

---

**Authors**

Robert M. Strongin, Eva Sharma, Hanno C. Erythropel, Nada O F Kassem, Alexandra Noël, D H. Peyton, and Irfan Rahman



OPEN ACCESS

# Chemical and physiological interactions between e-liquid constituents: cause for concern?

Robert M Strongin ,<sup>1</sup> Eva Sharma ,<sup>2</sup> Hanno C Erythropel ,<sup>3</sup>  
Nada O F Kassem ,<sup>4</sup> Alexandra Noël,<sup>5</sup> D H Peyton ,<sup>1</sup> Irfan Rahman <sup>6</sup>

<sup>1</sup>Chemistry, Portland State University, Portland, Oregon, USA

<sup>2</sup>Westat Inc, Rockville, Maryland, USA

<sup>3</sup>Chemical and Environmental Engineering, Yale University, New Haven, Connecticut, USA

<sup>4</sup>CBEACH, San Diego State University Research Foundation, San Diego, California, USA

<sup>5</sup>Department of Comparative Biomedical Sciences, Louisiana State University, Baton Rouge, Louisiana, USA

<sup>6</sup>Department of Environmental Medicine, University of Rochester, Rochester, New York, USA

## Correspondence to

Dr Robert M Strongin, Chemistry, Portland State University, Portland, Oregon, USA; rmstrongin@gmail.com

Received 8 December 2023

Accepted 11 April 2024

## ABSTRACT

Studies of Electronic Nicotine Delivery Systems (ENDS) toxicity have largely focused on individual components such as flavour additives, base e-liquid ingredients (propylene glycol, glycerol), device characteristics (eg, model, components, wattage), use behaviour, etc. However, vaping involves inhalation of chemical mixtures and interactions between compounds can occur that can lead to different toxicities than toxicity of the individual components. Methods based on the additive toxicity of individual chemical components to estimate the health risks of complex mixtures can result in the overestimation or underestimation of exposure risks, since interactions between components are under-investigated. In the case of ENDS, the potential of elevated toxicity resulting from chemical reactions and interactions is enhanced due to high operating temperatures and the metallic surface of the heating element. With the recent availability of a wide range of e-liquid constituents and popularity of do-it-yourself creation of e-liquid mixtures, the need to understand chemical and physiological impacts of chemical combinations in ENDS e-liquids and aerosols is immediate. There is a significant current knowledge gap concerning how specific combinations of ENDS chemical ingredients result in synergistic or antagonistic interactions. This commentary aims to review the current understanding of chemical reactions between e-liquid components, interactions between additives, chemical reactions that occur during vaping and aerosol properties and biomolecular interactions, all of which may impact physiological health.

## INTRODUCTION

The impact of Electronic Nicotine Delivery Systems (ENDS) toxicity on physiological health is an ongoing concern and a better understanding is needed to guide research themes and regulations. The toxicity of ENDS depends on factors including vaping patterns of an individual, device design and operation, power and resulting heat output, as well as the chemical makeup of the refill liquid ('e-liquid').<sup>1</sup> Vaping involves aerosolising mixtures of nicotine, solvents, flavorants and various other additives. As of 2017, there were approximately 15 500 available e-liquid formulations<sup>2</sup> comprised of >200 flavorant chemicals.<sup>3</sup> The number of e-liquid formulations appears to have trended upwards to the present time since a 2017 study, which is the most thorough recent report to date. For example, a study published in 2021 reported close to 20 000 different commercial e-liquids with 250 flavour descriptors.<sup>4</sup> Furthermore, do-it-yourself (DIY) e-liquid preparation, wherein

## WHAT THIS PAPER ADDS

- ⇒ Electronic Nicotine Delivery Systems (ENDS) usage results in exposure to complex chemical mixtures.
- ⇒ ENDS risk assessments for regulatory purposes mainly rely on assessments of individual chemicals or of composite commercial mixtures of e-liquids and/or their corresponding aerosols. There is a significant knowledge gap concerning the health risks from the combined exposure to multiple chemicals from ENDS usage.
- ⇒ This brief report highlights recent studies showing that specific combinations of compounds associated with ENDS can lead to altered chemical emission profiles or non-additive biological effects.

people who use ENDS create their own e-liquid mixtures, has become increasingly popular, in part as a reaction to an enforcement policy of flavoured cartridge-based ENDS.<sup>5 6</sup> The popularity of mixing e-liquid flavours in the USA was reported in Wave 2 (2014–2015) of the Population Assessment of Tobacco and Health Study, a nationally representative study of youth (12–17 years) and adults (18+ years), where 38.7% youth and 21.8% adults who used ENDS in the past 30 days reported using more than one flavour in their e-liquid.<sup>7</sup>

There has been extensive effort in the ENDS field towards the analysis of commercial e-liquid and aerosol chemical components. Gas chromatography and liquid chromatography, incorporating a variety of detection systems, particularly mass spectrometry, are the major analytical methods used to date.<sup>1</sup> However, a wide variety of specific sampling and analytical techniques have been developed for ENDS research<sup>1</sup> to address the challenging complexity and dynamics of the ENDS product landscape.<sup>8</sup>

ENDS risk assessments for regulatory purposes largely rely on assessments of individual chemicals or of composite commercial mixtures of e-liquids and/or their corresponding aerosols. There is a significant knowledge gap concerning the health risks from the combined exposure to multiple chemicals from ENDS usage. For example, chemical studies have shown that ENDS solvents and other molecular components can undergo incomplete combustion (thermal oxidation) reactions to a varying degree, depending on the device power, e-liquid makeup and puffing behaviour.<sup>1</sup> Indeed, e-cigarettes have also been described as 'chemical reactors' in literature.<sup>9 10</sup> Other reactions besides



© Author(s) (or their employer(s)) 2024. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

**To cite:** Strongin RM, Sharma E, Erythropel HC, et al. *Tob Control* Epub ahead of print: [please include Day Month Year]. doi:10.1136/tc-2023-058546

incomplete combustion may also occur between specific e-liquid ingredients, such as flavorant acetal formation during storage, thereby further altering molecular structures and properties, which contributes to the complex chemical mixtures of ENDS aerosols.<sup>11</sup> Moreover, studies of the *in vitro* or *in vivo* biological effects of vaping have focused on either a specific e-liquid chemical ingredient, such as a single additive, or on exposure to commercial, composite e-liquids or aerosols thereof.<sup>12</sup> However, it is well-known in the pharmaceutical field that different drug molecules may produce non-additive synergistic or antagonistic physiological effects when used in combination.<sup>13</sup>

Herein, this paper aims to describe the current understanding of chemical and biological properties of specific chemical combinations found in ENDS e-liquids and aerosols. This issue is relatively under-investigated,<sup>14,15</sup> but is timely and significant considering the extensive availability of e-liquid formulations and the popularity of DIY vaping. The examples described below provide evidence that interactions between certain e-liquid ingredients can lead to unique, non-additive toxicological effects.

### Chemical reactions between e-liquid components

A major difference between e-cigarettes and cigarettes is the fact that e-cigarette chemical formulations are liquids: it is well-known that organic molecules are typically more reactive in solution since dissolution enhances intermolecular interactions. For example, a striking illustration of changes to e-liquid compositions on storage was discovered by Erythropel *et al.* Their findings suggested that common aldehyde flavorants such as benzaldehyde, cinnamaldehyde, citral, ethylvanillin and vanillin readily react with the e-liquid solvents propylene glycol (PG) and glycerol (GL) to form substantial amounts of new compounds (acetals, conversions of 50%–95% depending on the aldehyde), within a timeframe of days to weeks, depending on the aldehyde in question. As expected, the resulting acetals exhibited differential physiological properties from either the flavorant or solvent components that form the respective acetal under the exclusion of a water molecule. PG-derived acetals diminished cellular energy metabolic functions, including basal respiration, ATP production and spare respiratory capacity.<sup>16</sup> PG/GL acetals induced cultured bronchial epithelial cell death at lower concentrations than the parent aldehydes.<sup>17</sup>

Interestingly, a study from the Netherlands (NL) did not detect benzaldehyde in a representative sample of 320 NL-marketed e-liquids.<sup>18</sup> However, benzaldehyde was reported by industry in the EU-CEG dataset to be present in 12.4% of >16k NL-marketed e-liquids.<sup>19</sup> It is quite possible that this inconsistency between reported data and analytical measurements is a result of benzaldehyde reactivity, such as with PG or GL to form acetals, or that the sample of commercial e-liquids did not contain benzaldehyde. Kerber *et al* later demonstrated that acetal formation could be inhibited by nicotine or water, but was promoted by the presence of benzoic acid.<sup>20</sup> Gschwend *et al* identified additional flavorant adducts, including acetals derived from ketones. They also found acetals in 32% (n=142) of the sampled commercial e-liquids.<sup>21</sup> Overall, it is concerning that several frequently used aldehyde flavorants are transformed to different species through reactions with the e-liquid solvents under storage conditions. This leads to new compounds that also exhibit unique toxicological properties, and occurs in potentially one-third, or more,<sup>20</sup> of all commercial products.

### Toxicological interactions between flavorant/additive molecules

Muthumalage *et al* were the first to determine that mixtures of e-liquid components can exhibit synergistic interactions.<sup>22</sup> They discovered that flavorants trigger reactive oxygen species (ROS) production and an associated inflammatory response in monocytes. Higher production of ROS (and cytotoxicity) was found in a 10 flavorant e-liquid mixture than was anticipated from the sum of the contributions of each specific flavorant.<sup>22</sup> The authors concluded that mixing multiple e-liquid flavorants caused the greatest cytotoxicity, implying a bigger health risk when mixtures of flavorants are present vs a single flavorant.

Marescotti *et al* found that, of 28 flavorants studied, 2-acetylthiazole, allyl hexanoate,  $\alpha$ -pinene, citronellol, guaiacol, linalool, methyl anthranilate, 3-methyl-2,4-nonanedione, 3-(methylthiol) propionaldehyde and phenethyl alcohol each resulted in human bronchial epithelial cell cytotoxicity.<sup>23,24</sup> When mixtures were investigated, cytotoxic properties deviated from the cytotoxicity observed using one specific flavorant. Citronellol had the greatest impact on mixture toxicity, whereas other chemicals resulted in synergistic effects.<sup>23</sup>

A study by Baldovinos *et al* analysed the cytotoxic effects of individual compounds and binary mixtures of a representative terpene (*R*-limonene) and an additive (triethyl citrate) on human lung cell models.<sup>25</sup> Data were analysed to determine the effects of 97:3 and 80:20% v/v (triethyl citrate/limonene) binary mixtures on BEAS-2B and A549 cells. LC<sub>50</sub> values and isobolograms were used to assess toxicity and chemical interactions. The data showed that limonene were more cytotoxic than triethyl citrate. Isobolographic analyses confirmed that the mixtures resulted in an antagonistic chemical interaction (ie, reduced toxicity). Further testing of different ratios of binary mixtures is needed for chemical interaction screening to inform safety assessments.

Larcombe *et al* investigated potential synergistic toxicity between nicotine and e-liquid solvents.<sup>26</sup> Using a mouse model, they showed that inhalation of 0/100% PG/GL aerosols, with or without 12 mg/mL of nicotine, impaired lung function (airway resistance, tissue damping and elastance) in a more pronounced manner than similar ENDS aerosols composed of 100/0% PG/GL. In this study, the presence of nicotine did not significantly impact the altered lung physiological responses, indicating that the inhalation of ENDS aerosols can impair lung function independently of nicotine content. Although an interaction between nicotine and e-liquid solvents leading to impaired lung function was not observed, this study revealed that ENDS solvents alone can induce adverse pulmonary effects *in vivo*.<sup>26</sup> However, molecular interactions between e-liquid solvents and nicotine impact nicotine exposure. For example, e-liquids composed of >70% PG have been found to aerosolise faster and lead to greater nicotine aerosol concentrations than GL-rich e-liquids.<sup>27</sup> This was further confirmed in people who used ENDS wherein PG-rich e-liquids was associated with increased nicotine delivery.<sup>28,29</sup>

Taken together, these *in vitro* and *in vivo* studies show toxicological interactions following exposures to e-liquid constituents, since non-additive toxicological effects of individual e-liquid components were noted. Despite these observations, no mechanisms or chemical reactions/interactions underlying these effects were evaluated. Thus, the chemical interactions between the e-liquid constituents leading to synergistic toxicological effects are still mainly unknown. A recent study by Pappas *et al*,<sup>30</sup> however, showed that interaction between acids in nicotine salt with coil metals resulted in enhanced transfer of metal oxide, highlighting a mechanism by which e-liquid constituents and

ENDS design characteristics interact, and by this mean, may lead to increased toxicity.

### Chemical reactions that occur during vaping

Molecular changes can be promoted during vaping due to catalytic reactions on the metal surface of ENDS heating filaments. Saliba *et al*<sup>31</sup> reported that the metal wires (eg, the iron/chromium/aluminium alloy Kanthal, stainless steel and the nickel/chromium alloy black nichrome) promoted the breakdown of PG to toxic carbonyls such as formaldehyde, methyl glyoxal and acetaldehyde, at temperatures <250°C, and as low as <80°C.<sup>31</sup> Importantly, this is further evident that dry coils and overheating are not the sole reasons<sup>32</sup> for elevated carbonyl toxicant emissions in vaped aerosols.

Formaldehyde derived from partial combustion of PG and GL during vaping has been shown to react with PG and GL to form new formaldehyde derivatives (hemiacetals).<sup>33</sup> Unlike gaseous formaldehyde, which is water soluble and deposits largely in the upper respiratory tract, the hemiacetals are less water soluble and preferentially partition into the aerosol particulate phase. They can thus be delivered more deeply into the airways/lungs than gaseous formaldehyde, which is concerning.<sup>34</sup>

### CONCLUSION

The examples described above embody strong evidence that interactions between certain e-liquid ingredients can lead to chemical reactions as well as cause unique, non-additive toxicological effects. Awareness and understanding of the health-related impact of specific combinations of ingredients is thus a necessary and important step to better understand the complex mixtures that e-cigarette liquids and importantly, the aerosols that users are actually exposed to, are. In turn, regulators and researchers should build on the existing although limited knowledge of the toxicity of mixtures as one important tool of many to better understand the toxicity of e-cigarette aerosol mixtures if the goal is to offer safer, truly lower risk alternatives to cigarettes, and to effectively protect public health in light of thousands of existing flavour combinations.

X Hanno C Erythropel @hanno\_ery

**Contributors** The authors contributed equally to writing and editing this manuscript. RMS, HCE, ES, NOFK, AN, DHP and IR contributed to planning, research, writing and proofreading.

**Funding** This study is a cross-institution collaborative project from the Toxicity Special Interest Group (SIG) supported, in part, by the Center for Coordination of Analytics, Science, Enhancement and Logistics (CASEL) in Tobacco Regulatory Science U54DA046060 (National Institute of Drug Abuse (NIDA) and the Food and Drug Administration's Center for Tobacco Products (FDA CTP). Support for authors was also provided by NIDA and FDA CTP awards R01ES025257 (RMS and DHP), U54DA036151 (HCE), U54DA036105, R03ES029441-02S1 (AN), U54 CA228110 (IR), T30IR0894 (NOFK) and U54DA046060-01 (ES). The content of this paper is solely the responsibility of the authors and does not represent the official views of the co-authors' institutions, the NIH, or the FDA.

**Competing interests** There are no competing interests.

**Patient consent for publication** Not applicable.

**Ethics approval** Not applicable.

**Provenance and peer review** Not commissioned; externally peer reviewed.

**Open access** This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

### ORCID iDs

Robert M Strongin <http://orcid.org/0000-0003-3777-8492>

Eva Sharma <http://orcid.org/0000-0002-1328-508X>  
 Hanno C Erythropel <http://orcid.org/0000-0003-3443-9794>  
 Nada O F Kassem <http://orcid.org/0000-0002-5403-7234>  
 D H Peyton <http://orcid.org/0000-0001-5828-055X>  
 Irfan Rahman <http://orcid.org/0000-0003-2274-2454>

### REFERENCES

- Strongin RM. E-cigarette chemistry and analytical detection. *Annu Rev Anal Chem (Palo Alto Calif)* 2019;12:23–39.
- Hsu G, Sun JY, Zhu SH. Evolution of electronic cigarette brands from 2013–2014 to 2016–2017: analysis of brand websites. *J Med Internet Res* 2018;20:e80.
- Aszyk J, Kubica P, Kot-Wasik A, *et al*. Comprehensive determination of flavouring additives and nicotine in e-cigarette refill solutions. part i: liquid chromatography-tandem mass spectrometry analysis. *J Chromatogr A* 2017;1519:45–54.
- Havermans A, Krüsemann EJZ, Pennings J, *et al*. Nearly 20 000 e-liquids and 250 unique flavour descriptions: an overview of the Dutch market based on information from manufacturers. *Tob Control* 2021;30:57–62.
- El-Hellani A, Soule EK, Daoud M, *et al*. Assessing toxicant emissions from e-liquids with DIY additives used in response to a potential flavour ban in e-cigarettes. *Tob Control* 2022;31:s245–8.
- Record RA, Groznik M, Sussman MA. Using the theory of planned behavior to anticipate DIY e-juice mixing among young adult International e-cigarette users. *Addict Health* 2023;15:23–30.
- Schneller LM, Bansal-Travers M, Goniewicz ML, *et al*. Use of flavored electronic cigarette refill liquids among adults and youth in the US—results from wave 2 of the population assessment of tobacco and health study (2014–2015). *PLoS One* 2018;13:e0202744.
- Strongin RM, Sharma E, Erythropel HC, *et al*. Emerging ENDS products and challenges in tobacco control toxicity research. *Tob Control* 2023;33:110–5.
- Jensen RP, Strongin RM, Peyton DH. Solvent chemistry in the electronic cigarette reaction vessel. *Sci Rep* 2017;7:42549.
- El-Hellani A, El-Hage R, Salman R, *et al*. Electronic cigarettes are chemical reactors: implication to toxicity. *Chem Res Toxicol* 2020;33:2489–90.
- Erythropel HC, Jabba SV, DeWinter TM, *et al*. Formation of flavorant–propylene glycol adducts with novel toxicological properties in chemically unstable e-cigarette liquids. *Nicotine Tob Res* 2019;21:1248–58.
- Stefaniak AB, LeBouf RF, Ranpara AC, *et al*. Toxicology of flavoring-and cannabis-containing e-liquids used in electronic delivery systems. *Pharmacol Ther* 2021;224:107838.
- Cascorbi I. Drug interactions—principles, examples and clinical consequences. *Dtsch Arztebl Int* 2012;109:546–55.
- Bopp SK, Barouki R, Brack W, *et al*. Current EU research activities on combined exposure to multiple chemicals. *Environ Int* 2018;120:544–62.
- Committee ES, More SJ, Bampidis V, *et al*. Guidance on harmonised methodologies for human health, animal health and ecological risk assessment of combined exposure to multiple chemicals. *Efsa Journal* 2019;17:e05634.
- Jabba SV, Diaz AN, Erythropel HC, *et al*. Chemical adducts of reactive flavor aldehydes formed in e-cigarette liquids are cytotoxic and inhibit mitochondrial function in respiratory epithelial cells. *Nicotine Tob Res* 2020;22:525–34.
- Jordt S-E, Caceres AI, Erythropel H, *et al*. Flavor-solvent reaction products in electronic cigarette liquids activate respiratory irritant receptors and elicit cytotoxic metabolic responses in airway epithelial cell. ERS International Congress 2020 abstracts; September 7, 2020
- Krüsemann EJZ, Pennings JLA, Cremers J, *et al*. GC-MS analysis of e-cigarette refill solutions: a comparison of flavoring composition between flavor categories. *J Pharm Biomed Anal* 2020;188:113364.
- Krüsemann EJZ, Havermans A, Pennings JLA, *et al*. Comprehensive overview of common e-liquid ingredients and how they can be used to predict an e-liquid's flavour category. *Tob Control* 2021;30:185–91.
- Kerber PJ, Peyton DH. Kinetics of aldehyde flavorant-acetal formation in e-liquids with different e-cigarette solvents and common additives studied by 1H NMR spectroscopy. *Chem Res Toxicol* 2022;35:1410–7.
- Gschwend G, Jenkins C, Jones A, *et al*. A wide range of flavoring-carrier fluid adducts form in e-cigarette liquids. *Chem Res Toxicol* 2023;36:14–22.
- Muthumalage T, Prinz M, Ansah KO, *et al*. Inflammatory and oxidative responses induced by exposure to commonly used e-cigarette flavoring chemicals and flavored e-liquids without nicotine. *Front Physiol* 2017;8:1130.
- Marescotti D, Mathis C, Belcastro V, *et al*. Systems toxicology assessment of a representative e-liquid formulation using human primary bronchial epithelial cells. *Toxicol Rep* 2020;7:67–80.
- Marescotti D, Mathis C, May A, *et al*. 2021. Toxicological assessment of flavors used in E-vapor products. Toxicological Evaluation of Electronic Nicotine Delivery Products: Elsevier. 367–83.
- Baldovinos Y, Archer A, Salamanca J, *et al*. Chemical interactions and cytotoxicity of terpene and diluent vaping ingredients. *Chem Res Toxicol* 2023;36:589–97.
- Larcombe AN, Janka MA, Mullins BJ, *et al*. The effects of electronic cigarette aerosol exposure on inflammation and lung function in mice. *Am J Physiol Lung Cell Mol Physiol* 2017;313:L67–79.

- 27 Talih S, Balhas Z, Salman R, *et al.* Transport phenomena governing nicotine emissions from electronic cigarettes: model formulation and experimental investigation. *Aerosol Sci Technol* 2017;51:1–11.
- 28 Spindle TR, Talih S, Hiler MM, *et al.* Effects of electronic cigarette liquid solvents propylene glycol and vegetable glycerin on user nicotine delivery, heart rate, subjective effects, and puff topography. *Drug Alcohol Depend* 2018;188:193–9.
- 29 Yan XS, D’Ruiz C. Effects of using electronic cigarettes on nicotine delivery and cardiovascular function in comparison with regular cigarettes. *Regul Toxicol Pharmacol* 2015;71:24–34.
- 30 Pappas RS, Gray N, Halstead M, *et al.* Lactic acid salts of nicotine potentiate the transfer of toxic metals into electronic cigarette aerosols. *Toxics* 2024;12:65.
- 31 Saliba NA, El Hellani A, Honein E, *et al.* Surface chemistry of electronic cigarette electrical heating coils: effects of metal type on propylene glycol thermal decomposition. *J Anal Appl Pyrolysis* 2018;134:520–5.
- 32 Jaegers NR, Hu W, Weber TJ, *et al.* Low-temperature (&lt;200 °C) degradation of electronic nicotine delivery system liquids generates toxic aldehydes. *Sci Rep* 2021;11:7800.
- 33 Jensen RP, Luo W, Pankow JF, *et al.* Hidden formaldehyde in e-cigarette aerosols. *N Engl J Med* 2015;372:392–4.
- 34 Pankow JF. Calculating compound dependent gas-droplet distributions in aerosols of propylene glycol and glycerol from electronic cigarettes. *J Aerosol Sci* 2017;107:9–13.