Portland State University

PDXScholar

University Honors Theses

University Honors College

Spring 2021

NHANES III Mortality Follow Up: Social Determinants Creating Disparity within the Health Outcomes of Persons with Diabetes

Claire B. Adams

Portland State University

Follow this and additional works at: <https://pdxscholar.library.pdx.edu/honorstheses>

Part of the [Clinical Epidemiology Commons](#), [Community Health and Preventive Medicine Commons](#), and the [Endocrinology, Diabetes, and Metabolism Commons](#)

Let us know how access to this document benefits you.

Recommended Citation

Adams, Claire B., "NHANES III Mortality Follow Up: Social Determinants Creating Disparity within the Health Outcomes of Persons with Diabetes" (2021). *University Honors Theses*. Paper 1131.

<https://doi.org/10.15760/honors.1162>

This Thesis is brought to you for free and open access. It has been accepted for inclusion in University Honors Theses by an authorized administrator of PDXScholar. Please contact us if we can make this document more accessible: pdxscholar@pdx.edu.

NHANES III Mortality Follow Up: Social Determinants Creating Disparity within the Health
Outcomes of Persons with Diabetes

Claire B. Adams ¹ and Carlos J. Crespo, DrPH ²

¹Portland State University Honors College, ²OHSU-PSU School of Public Health

Abstract

Given the recent research into the social determinants affecting health outcomes, the goal of this thesis is to examine whether a non-urban setting, among other social determinants of health, increases the risk of mortality among the United States diabetic population. We examined the relationship of urban and non-urban diabetes rates among those aged 40-74 years using a national sample of diabetic and nondiabetic Americans. The data was pulled from the Third National Health and Nutrition Examination Survey (NHANES III) which was conducted from 1988 to 1994. After excluding pregnant women, those who did not complete a fasting AM glucose, and those with missing information on variables for our analysis, a total of 3,219 American subjects (1,635 women and 1,584 men) were studied. Our results show no significant association between urban living and dying with diabetes. However, we found a significant interaction between urban living and insurance status in predicting diabetes mortality among persons with diabetes. Further research should investigate the influence of having health insurance in urban and rural settings. Therefore, expanding research on the social determinants of health currently impacting diabetes mortality rates has the potential to influence the work being done by healthcare professionals to address access within rural communities.

Keywords: National Health and Nutrition Examination Survey (NHANES), NHANES III, social determinants of health, health disparities, Centers for Disease Control and Prevention (CDC)

Introduction

Effective management and care of persons with diabetes in America has become an increasing challenge within the Public Health community over the past several decades¹. As one of the most rapidly growing chronic diseases, diabetes has placed a growing economic burden on the United States healthcare system and the American taxpayers, with an approximately 26% rise in diabetic healthcare costs from 2015-2017². In 2019, a study found the mean annual direct cost per person was between \$220 and \$7,600, with medications and hospitalizations contributing to the majority of the costs³. Given the high cost incurred by those with diabetes, access to consistent health insurance coverage, whether private or public, is associated with improved diabetes care management and health outcomes⁴. Diabetes is associated with a multitude of risk factors, therefore proper access to care is pivotal in preventing negative health outcomes and higher rates of mortality.

The common risk factors for diabetes studied and discussed by the majority of those in the discourse community revolve around the individuals smoking status, obesity, and physical inactivity, amongst others. However, what if those are just the foundation for a larger issue at hand? One must look at the social determinants of this disease. In reality, the social determinants of health are just as likely, if not more, to have an impact on the risks and health outcomes of those with diabetes⁵. This goes beyond the associated comorbidities of the disease instead focusing on the conditions in which people were born in, raised, and currently live. Although all the social determinants of health play essential roles in health effects, “geography and health are intrinsically linked. Where we are born, live, study and work directly influences our health experiences: the air we breathe, the food we eat, the viruses we are exposed to and the health

services we can access”⁶. Traditional clinical interventions for health continue to create gaps of care and access for those whose social determinants place them at a disadvantage to others. Clinical outcomes for persons with diabetes continue to be highly influenced by social determinants of health, many of which still lack proper research and policy-based solutions⁵.

As the number of persons living with diabetes continues to increase, providing proper care necessary to reduce associated health outcomes and mortality becomes an increasingly important issue for the Public Health community to tackle. Based on the most recent estimates by the Center for Disease Control (CDC), there are an estimated 34.2 million Americans, or 10.5% of the population, living with diabetes². The CDC also estimates that approximately 88 million people adults have prediabetes². With proper preventative medicine and constant healthcare access, the progression to type 2 diabetes mellitus for these 88 million people is preventable. Diabetes is associated by many long-term comorbidities which currently lead to more adult cases of blindness, renal failure, and amputations than any other disease in America⁷. Prevention of this disease is necessary to keep healthcare costs down, as well as communities in good health. Yet, for rural people, these geographic barriers to healthcare access continue to exacerbate the prevention of proper management of the disease⁵. These barriers are not limited to zip code alone but encompass other obstacles in health such as proximity to public parks, safe places to walk or exercise, food security, and a primary care physician or a needed specialist nearby.

Methods

The NHANES III study is a nationwide probability sample of 39,695 people aged 2 months and older which was conducted from 1988 to 1994 in two phases⁸. This Diabetic Mortality follow-up study was performed through linking the death certificate data available in the National Death Index (NDI) to the NHANES III data. To perform our analysis, we excluded

women who were pregnant at the time of the study, those who did not complete a morning fasting glucose blood test, those outside of the 40-74 year age range, and those with missing variables necessary for our analysis. Our analytic sample includes the remaining 3,219 persons with complete data on diabetes status.

All analysis was conducted using the statistical software SAS 9.4 program. We used Cox Proportional Hazards Models to investigate the association between the mortality of persons with diabetes while controlling for exposures in insurance status and geographic areas. We used the geographic areas *urban* and *non-urban* as defined and published by NHANES III study. The age range of 40-74 was selected given a relative stability with education and income levels within this age group and that this age group completed an Oral Glucose Tolerance Test.

Results

After analysis using SAS 9.4, we were left with the subjects shown in Table 1, which were then further evaluated by exposure to urban and non-urban residence, as seen in Table 2.

Table 1. Description of Subjects

	Men (95% CI)	Women (95% CI)
Age, years	53.96 (53.19-54.72)	54.57 (53.68-55.46)
Race		
Non-Hispanic Whites, %	49.00 (47.00-51.00)	51.00 (49.00-53.00)
Non-Hispanic Blacks, %	43.47 (40.78-46.17)	56.53 (53.83-59.22)
Mexican Americans, %	49.50 (45.96-53.05)	50.50 (46.95-54.04)
Other race/ethnicity, %	49.01 (42.54-55.47)	51.00 (44.53-57.46)
Education Level		
Less than High School, %	27.56 (23.55-31.56)	25.00 (22.17-27.83)
High School, %	25.49 (21.62-29.35)	37.60 (34.70-40.48)
More than High School, %	46.75 (42.73-50.76)	37.24 (33.29-41.18)
Insurance status		
Yes, %	92.56 (89.42-95.66)	90.63 (88.16-93.09)
Urban Residence		
Yes, %	47.77 (37.21-58.33)	44.82 (35.76-53.88)

Table 2. Demographic Distribution According to Exposure Variable (Urban vs. Non-Urban)

	Urban Residence	Non-Urban Residence
	% (95% CI)	% (95% CI)
Sex		
Men, %	50.12 (47.01-53.24)	47.16 (45.06-49.25)
Women, %	49.88 (46.76-52.99)	52.84 (50.75-54.94)
Age, years	54.18 (52.13-55.24)	54.35 (53.34-55.36)
Race		
Non-Hispanic Whites, %	70.41 (63.70-77.11)	87.19 (83.73-90.65)
Non-Hispanic Blacks, %	11.25 (8.72-13.78)	6.82 (4.83-8.80)
Mexican Americans, %	4.50 (3.04-5.95)	2.91 (2.02-3.80)
Other race/ethnicity, %	13.85 (7.83-19.88)	3.08 (1.13-5.04)
Education Level		
Less than High School, %	22.03 (17.83-26.22)	29.87 (24.60-35.13)
High School, %	28.38 (24.67-32.10)	34.58 (31.43-37.74)
More than High School, %	49.24 (43.90-54.59)	35.49 (30.22-40.76)
Insurance status		
Yes, %	90.13 (86.17-94.09)	92.81 (90.14-95.47)

Then based on the demographic distribution, the subjects were categorized based on the exposure to the outcome variables (diabetes mellitus status or non-diabetes status) which appears in Table 3.

Table 3. Demographic Distribution According to Outcome Variable (Diabetes Mellitus)

	Diabetes Mellitus (95% CI)	No Diabetes (95% CI)
Sex		
Men, %	57.70 (50.49-64.90)	39.57-36.21-42.93
Women, %	42.30 (35.10-49.51)	60.43 (57.07-63.79)
Age, years		
	59.03 (57.17-60.90)	51.58 (50.82-52.33)
Race		
Non-Hispanic Whites, %	12.52 (10.64-14.41)	44.92 (41.13-48.71)
Non-Hispanic Blacks, %	17.26 (14.28-20.25)	31.35 (27.51-35.20)
Mexican Americans, %	22.57 (19.37-25.77)	28.33 (24.81-31.82)
Other race/ethnicity, %	10.35 (5.51-15.20)	33.01 (23.35-42.66)
Education Level		
Less than High School, %	35.36 (26.77-43.95)	17.02 (14.23-19.81)
High School, %	37.86 (28.49-47.23)	30.25 (26.60-33.90)
More than High School, %	26.78 (17.09-36.48)	52.63 (48.37-56.88)
Insurance status		
Yes, %	90.24 (84.39-96.08)	92.61 (90.09-95.13)
Urban Residence		
Yes, %	41.80 (30.71-52.89)	46.80 (36.53-57.07)

Lastly to examine the role of insurance coverage among those in urban and non-urban residence, Table 4 shows the prevalence of diabetes according to these variables. As seen here, when exposed to non-urban residence and lack of health insurance coverage, there is a 17.51 prevalence of diabetes compared to the appropriate exposures (8.61 prevalence).

Table 4. Prevalence of Diabetes According to Health Insurance Status in Urban and Non-Urban Areas

	Diabetes Mellitus (95% CI)
Live in urban area	
Covered by Health Insurance	12.99 (10.55-15.44)
Not Covered by Health Insurance	8.61 (3.90-13.31)
Live in non-urban area	
Covered by Health Insurance	13.17 (10.69-15.64)
Not Covered by Health Insurance	17.51 (7.88-27.14)

Given the exposures, to examine the Hazard Risk for diabetes mortality and the interaction with urban and non-urban residence, as well age and insurance coverage status the Cox Proportional Hazard Model was applied to the data (Table 5).

Table 5. Cox Proportional Hazard Models

	Living in Non-Urban Area	
	HR (95% CI)	N
Model 1. Diabetes mortality	1.45 (0.85-1.55)	57
Model 2. Diabetes mortality adjusted for age	1.117 (0.83-1.50)	55
Model 3. Diabetes mortality adjusted for age and health insurance status	1.079 (0.804-1.449)	55

Discussion

The most meaningful findings from this mortality study concerns the mortality of those with diabetes who live in a non-urban setting without health insurance coverage. Living in an urban setting was not a predictor for dying with diabetes. However, there was a predictive significance with a $P=0.0006$ ($HR=2.191$ ($1.43-3.356$)) between people without health insurance and non-urban counties for a model ran with the Cox Proportional Hazard. But due to the minute sample size in the model, we were unable to further stratify the study population. Among the small sample size being a limitation to our study, there were many people with missing variables that were unable to be included, missing variables with the specific category of insurance coverage, and no differentiation between type 1 and type 2 diabetes mellitus. These limitations, along with a lack of health insurance and non-urban residency, exhibit the need for future research within the Public Health and endocrinology community. A longitudinal study which

focuses on people with diabetes in rural and urban communities and their access to consistent healthcare is needed.

In highlighting the role of insurance coverage combined with geographic location it exposes the potential barriers present to access proper healthcare for those with diabetes.

Figure 1. Free Clinic Locations by State, as of May 2015

Source: The Henry J. Kaiser Family Foundation¹⁰

According to a recent study, adequate access to primary care is associated with potential healthcare savings related to preventable hospitalizations for those living in rural areas that are uninsured⁹. This calls for further attention to the association present with geographic areas and health insurance coverage. However, insurance coverage only explains a portion of this issue. Access to healthcare for those in rural areas may be just as, if not more, important. Does having health insurance provide immediate access to a healthcare provider or is this access only determined by one's geographic location?

Given that only 22% of free clinics within the United States are located in rural areas compared to 78% in urban areas, those without health insurance in rural areas are left at a notable disadvantage¹⁰. Not only are many rural counties at a disadvantage, but there are also seventeen states without a single free clinic within a rural area¹⁰. As shown in Figure 1, those uninsured who do not live in or near rural communities are unable to access medical care within a reasonable distance. It can be assumed improved diabetes control could easily be attained through better access to healthcare^{11, 12}.

The social determinant of geographic location and environment plays an essential role in health outcomes impacting the social services provided which allow greater access to certain groups. Future research within this could provide insight into optimal future locations for free medical clinics and educational outreach programs for diabetes, as well as increase overall access to healthcare for disease prevention and management. This would allow for greater access to those who may be several hundred miles from their nearest specialist. Adequate access to consistency in healthcare could be the key to lowering negative health outcomes, mortality rates, and providing economic savings for those living with diabetes¹³.

References

1. Centers for Disease Control and Prevention. National Diabetes Statistics Report, 2020. Atlanta, GA: Centers for Disease Control and Prevention, U.S. Dept of Health and Human Services; 2020.
2. American Diabetes Association. Economic costs of diabetes in the U.S. in 2017. *Diabetes Care*. 2018;41(5):917-928. doi:10.2337/dci18-0007
3. Ramzan S, Timmins P, Hasan SS, Babar Z-U-D. Cost analysis of type 2 diabetes mellitus treatment in economically developed countries. *Expert Review of Pharmacoeconomics & Outcomes Research*. 2018;19(1):5-14. doi:10.1080/14737167.2018.1513790
4. Doucette ED, Salas J, Scherrer JF. Insurance coverage and diabetes quality indicators among patients in NHANES. *Am J Manag Care*. 2016;22(7):484-490.
5. Walker RJ, Strom Williams J, Egede LE. Influence of Race, Ethnicity and Social Determinants of Health on Diabetes Outcomes. *Am J Med Sci*. 2016;351(4):366-373. doi:10.1016/j.amjms.2016.01.008
6. Dummer TJB. Health geography: Supporting public health policy and planning. *Canadian Medical Association Journal*. 2008;178(9):1177. doi:10.1503/cmaj.071783
7. Crandall JP, Knowler WC, Kahn SE, et al. The prevention of type 2 diabetes. *Nat Clin Pract Endocrinol Metab*. 2008;4(7):382-393. doi:10.1038/ncpendmet0843
8. NHANES III (1988-1994). Centers for Disease Control and Prevention. <https://wwwn.cdc.gov/nchs/nhanes/nhanes3/default.aspx>
9. Zhang W, Mueller KJ, Chen LW. Uninsured hospitalizations: rural and urban differences. *J Rural Health*. 2008;24(2):194-202. doi:10.1111/j.1748-0361.2008.00158.x

10. Martin A, Bhavsar G, Workman L, Probst JC. Free Clinics in the Rural Safety Net, 2014. *South Carolina Rural Health Research Center*. September 2015:1-11
11. Gertz AM, Frank S, Blixen CE. A survey of patients and providers at free clinics across the United States. *Journal of Community Health*. 2010;36(1):83-93. doi:10.1007/s10900-010-9286-x
12. McLendon SF, Wood FG, Stanley N. Enhancing diabetes care through care coordination, telemedicine, and education: Evaluation of a rural pilot program. *Public Health Nurs*. 2019;36(3):310-320. doi:10.1111/phn.12601
13. Mallow JA, Theeke LA, Whetsel T, Barnes ER. Diabetes group medical visits and outcomes of care in low-income, rural, uninsured persons. *Open J Nurs*. 2013;3(3):314-322. doi:10.4236/ojn.2013.33043