Portland State University

PDXScholar

Dissertations and Theses

Dissertations and Theses

1984

Inquiry into the causes and significance of cytoplasmic vacuolation of neutrophils in the peripheral circulation

Gary Xavier Haight
Portland State University

Follow this and additional works at: https://pdxscholar.library.pdx.edu/open_access_etds

Part of the [Biology Commons](#), and the [Cell Biology Commons](#)

Let us know how access to this document benefits you.

Recommended Citation

Haight, Gary Xavier, "Inquiry into the causes and significance of cytoplasmic vacuolation of neutrophils in the peripheral circulation" (1984). *Dissertations and Theses*. Paper 3400.

<https://doi.org/10.15760/etd.5283>

This Thesis is brought to you for free and open access. It has been accepted for inclusion in Dissertations and Theses by an authorized administrator of PDXScholar. Please contact us if we can make this document more accessible: pdxscholar@pdx.edu.

AN ABSTRACT OF THE THESIS OF Gary Xavier Haight for the
Master of Science in Biology presented May 4, 1984.

Title: Inquiry into the Causes and Significance of
Cytoplasmic Vacuolation of Neutrophils in the
Peripheral Circulation.

APPROVED BY MEMBERS OF THE THESIS COMMITTEE:

[REDACTED]

Lester J. Newman, Co-Chairman

[REDACTED]

Deborah A. Duffield, Co-Chairman

[REDACTED]

Malcolm S. Lea

[REDACTED]

Quentin D. Clarkson

One-half million cases of septicemia are reported annually with a mortality of around 35%. Diagnosis depends in part on blood cultures which require one to two weeks. It would be advantageous if an early sign of septicemia were available.

Cytoplasmic vacuolation of polymorphonuclear neutrophils is occasionally seen in the peripheral blood smears of

patients who have infections. The object of this work was to determine the cause and significance of cytoplasmic vacuolation with the goal of using the occurrence of vacuolation as an early indicator of infection.

It was necessary first to establish that the phenomenon was not simply an artifact of slide preparation technique. Having established that, I examined the records of 59 patients displaying vacuolation. The major finding was that presence of vacuolation is weakly correlated with a number of variables. Sixty-three variables were examined for each of 59 patients using multiple regression analysis. Fifteen variables made a significant contribution to the regression equation and accounted for 62% of the variability seen in percent vacuolation. Among the factors contributing were patient age, sex, presence of infection and history of alcohol abuse.

An animal model was developed using mice; a number of interesting but conflicting results were obtained. Although female in patients were represented in the vacuolation group twice as often as male inpatients, in mice the males were affected at an equal frequency but had twice as high a percent vacuolation. In addition I observed that the control animals ultimately demonstrated vacuolation to a degree equivalent to that of the experimental animals. Animals fed ethanol and animals who received bacterial injection developed vacuolation but those who got both the injection and

the ethanol had a lower incidence of vacuolation.

Finally in vitro induction of vacuolation was attempted using plasma from affected individuals and cells from healthy volunteers. Vacuolation appeared first in the experimental tubes (patient plasma and volunteer cells) but was eventually seen in the control tubes (volunteer plasma and cells).

The usefulness of cytoplasmic vacuolation as a diagnostic tool is limited by the fact that its presence is transient and is correlated with a number of factors besides infection. However, since 72% of the patients having vacuolated neutrophils are ultimately diagnosed as having an infection, vacuolation, when it is present, should serve as a warning of the possibility of infection.

INQUIRY INTO THE CAUSES AND SIGNIFICANCE OF
CYTOPLASMIC VACUOLATION OF NEUTROPHILS
IN THE PERIPHERAL CIRCULATION

by

GARY XAVIER HAIGHT

A Thesis submitted in partial fulfillment of the
requirements for the degree of

MASTER OF SCIENCE
in
BIOLOGY

Portland State University

1984

TO THE OFFICE OF GRADUATE STUDIES AND RESEARCH:

The members of the Committee approve the thesis of
Gary Xavier Haight presented May 4, 1984.

Lester J. Newman, Co-Chairman

Deborah A. Duffield, Co-Chairman

Malcolm S. Lea

Quentin D. Clarkson

APPROVED:

W. Herman Taylor, Head, Department of Biology

Stanley E. Rauch, Dean of Graduate Studies and Research

ACKNOWLEDGEMENTS

It is unlikely that without the enthusiastic teaching, encouragement and friendship of Dr. Debbie Duffield this Master's project would ever have begun.

The friendship and guidance of Dr. Lester Newman and Dr. Malcolm Lea and the statistical expertise of Dr. Deane Clarkson have been invaluable and much appreciated.

My thanks go to Carol Wilson who initiated me into the mysteries of multiple regression and the Honeywell 66.

I am grateful to Paul Funatake for his professional production of the graphic figures which went into this thesis.

I wish to thank Karen Dodge, Carol Titsworth, and the hematology staffs of Emanuel and Providence Hospitals for staining the hundreds of slides generated in this project and for their dedication in supplying me with blood smears from affected patients.

Very special thanks go to my wife, Rose, for her unwavering support and encouragement, and for never allowing me to give up when things appeared impossible. I couldn't have done it without you.

TABLE OF CONTENTS

	PAGE
ACKNOWLEDGEMENTS	iii
LIST OF TABLES	vi
LIST OF FIGURES	vii
INTRODUCTION	1
 CHAPTER I EVALUATION OF ANTICOAGULANTS	
INTRODUCTION	7
MATERIALS AND METHODS	7
RESULTS	8
DISCUSSION	10
 CHAPTER II VACUOLATION IN PATIENTS	
INTRODUCTION	12
MATERIALS AND METHODS	12
RESULTS	13
DISCUSSION	18
 CHAPTER III VACUOLATION IN ANIMALS	
INTRODUCTION	26
MATERIALS AND METHODS	27
RESULTS	29
DISCUSSION	34

CHAPTER IV NITROBLUE TETRAZOLIUM REDUCTION

INTRODUCTION	39
MATERIALS AND METHODS	40
RESULTS	41
DISCUSSION	43

CHAPTER V IN VITRO VACUOLATION

INTRODUCTION	47
MATERIALS AND METHODS	47
RESULTS	48
DISCUSSION	49
FINAL DISCUSSION	55
REFERENCES	58
APPENDIX	62

LIST OF TABLES

TABLE		PAGE
I	Occurrence of Vacuolation in Anticoagulated Blood	10
II	Distribution by Diagnosis, Age, and Sex of Inpatients Displaying Vacuolation	17
III	Sex Effect of Neutrophil Vacuolation in Inpatients.	20
IV	Correlation Coefficients for Independent and Combined Variables.	22
V	Variables in the Regression Equation.	23
VI	Presence of Vacuolation and Positive Nitroblue Tetrazolium Tests in Four Groups of Mice Run Number One	42
VII	Presence of Vacuolation and Positive Nitroblue Tetrazolium Tests in Four Groups of Mice Run Number Two	43
VIII	Percent Vacuolation of Neutrophils Exposed to Patient Plasma Versus Time of Exposure. .	49
IX	Hourly Comparison of Mean Percent Vacuolation of Experimentals Versus Controls	52

LIST OF FIGURES

FIGURE	PAGE
1. Mature polymorphonuclear neutrophil. 2100x. . .	2
2. Mature neutrophil showing cytoplasmic vacuolation. 2800x.	6
3. Neutrophil showing toxic granulation at 4 hours. EDTA used as anticoagulant. 2300x.	9
4. Neutrophil showing slight vacuolation at 1 hour. Heparin used as anticoagulant. 1900x. . .	9
5. Frequency of occurrence versus percent vacuolation for hospital inpatients.	15
6. Mature neutrophil with large vacuole. 2300x..	19
7. Mature neutrophils with many small cytoplasmic vacuoles. 2300x.	19
8. Normal mature neutrophil from a mouse. 2300x. .	30
9. Mean percent vacuolation for mice by sex and experimental group.	31
10. Blood smear with distorted background. 2300x. .	33
11. Occurrence of disrupted background by sex and experimental group.	36
12. Mouse neutrophil grossly distorted by cytoplasmic vacuoles. 2300x.	38
13. Positive NBT test stimulated by phorbol 12-myristate 13-acetate. 2300x.	44

14. Positive NBT test stimulated experimentally. . .	44
15. Percent vacuolation versus time for neutrophils <u>in vitro</u>	50

INTRODUCTION

Phagocytic leukocytes are divided into two primary groups: the monocytic and the granulocytic series. The granulocytes are further divided into basophils, eosinophils, and neutrophils based on the staining characteristics of their cytoplasmic granules. In a normal peripheral blood smear the predominant leukocyte is the neutrophil. The most commonly seen neutrophil is the mature segmented form, easily identified by its multilobed nucleus and characteristic staining (Figure 1). The only other developmental stage normally seen in a peripheral blood smear is the form with an unsegmented or "band" nucleus.

Many morphologic changes in the neutrophil have been described in disease states (Stiene-Martin 1982a, Stiene-Martin 1982b). Sutro (1933) described the phenomenon of vacuolation of neutrophil cytoplasm as occasionally being seen in patients having bacterial infections. Meranze (1935) described neutrophils having a "motheaten" appearance from cases of both infectious and non-infectious disease. One investigator (Gordin 1952) attempted to correlate extent of vacuolation with severity of infection, while others (Ponder et al 1942) used cytoplasmic vacuolation of neutrophils as a measure of the severity of disease. Zieve et al (1966) found vacuolated cytoplasm in the neutrophils of 3%

Figure 1. Mature polymorphonuclear neutrophil.
2100X.

of 3500 blood smears from a general hospital population. Of this 3%, 98% were from patients whose primary diagnosis was bacterial infection. Cytoplasmic vacuolation has been observed in the neutrophils of persons treated for ethanol abuse (Lindenbaum and Lieber 1969; Davidson and McPhie 1980) and in the neutrophils of dogs being fed ethanol (Beard and Knott 1966).

Vacuolation of neutrophil cytoplasm was reported as being transient, lasting only 12 to 24 hours (Zieve et al 1966; Emerson 1970). The transient nature of cytoplasmic vacuolation in patients who respond to treatment corresponds well with the turnover rate of neutrophils in the peripheral circulation. The implication is that it is the circulating neutrophils which are affected, and that when stimulation ceases, they are rapidly cleared from the peripheral circulation. The half-life for neutrophils in the peripheral circulation has been variously reported (Cline 1975; Greenwalt 1977; Wintrobe et al 1981) as 6 to 7.5 hours. The persistence of vacuolated cells was seen as a very negative prognostic sign (Wintrobe et al 1981; Grange 1980).

Spontaneous formation of cytoplasmic vacuoles in neutrophils in wet mounts was reported by Gordin (1952) and by Bessis (1972). Gordin (1952) noted that vacuole size increased at a constant rate, once formation had begun, leading him to the conclusion that vacuole formation was due to a physical process not related to cellular metabolism.

He further concluded that since vacuoles form in wet films and are more often seen in older, more mature neutrophil forms, the origin of vacuoles is not a maturational disturbance, but is due to some peripheral insult. McCall et al (1969, 1979) claimed that cytoplasmic vacuoles were autophagosomes caused by lysosomal membrane instability and that this instability was precipitated by exposure to bacterial chemotoxins. Bessis (1972) believed the vacuoles formed from a swelling of the mitochondria, rough endoplasmic reticulum and the golgi apparatus.

Solberg and Hellum (1972) examined neutrophils from bacterially infected patients and found that they were able to ingest but were less able to kill bacteria in vivo than were neutrophils taken from afebrile patients. Vacuolated neutrophils were present in 27% of the infected patients and in only 8% of the control patients. Cytoplasmic vacuolation is generally held to be a sign of degeneration and impending cell death (Bessis 1972, Wintrobe et al 1981). The phenomenon has been noted in senescent human fibroblasts (Finch and Hayflick 1977) and in the senescent hyphae of the mold Podospora anserina (Belcour and Begel 1978, Esser and Keller 1976).

As many as one-half million cases of septicemia are reported annually with an overall mortality around 35% (Rubin 1981). Survival depends upon rapid diagnosis and treatment. A complication is that blood cultures require

one to two weeks. It would be advantageous to have an early and reliable sign of septicemia.

I have noted from personal observations that cytoplasmic vacuolation of polymorphonuclear neutrophils is occasionally seen in the peripheral blood smears of patients having septicemia. Since an admission blood smear is examined for most hospital patients and since the vacuolation is readily apparent (Figure 2) I decided to examine more closely the relationship between infection and cytoplasmic vacuolation.

It was necessary first to assess the influence of the anticoagulants used in specimen collection on vacuole formation (Chapter I). Information was then gathered on patients displaying vacuolation to find the variables which best correlated with appearance of vacuoles (Chapter II). An animal model for cytoplasmic vacuolation was investigated (Chapter III), and an attempt was made, using the nitroblue tetrazolium reduction test, to determine the applicability of the animal model (Chapter IV). Finally, Chapter V deals with the in vitro induction of vacuolation in neutrophils.

Figure 2. Mature neutrophil showing cytoplasmic vacuolation. 2800X.

CHAPTER I

INTRODUCTION

There is a common belief both among hematology lab technologists and in the literature (Davidson and Henry 1969; Shafer 1982) that exposure of blood to anticoagulants induces artifacts, including vacuolation, which render blood smears useless for diagnostic purposes. However, several hospitals in this metropolitan area make their diagnostic smears from anticoagulated blood, usually using EDTA, as anticoagulant.

Since, in some phases of my work it was easiest to obtain anticoagulated blood and since it seemed strange that technologists who warned of anticoagulant artifact induction were, in fact, using anticoagulated blood for their smears, this study was undertaken to ascertain the vacuole-inducing capability of various anticoagulants given prolonged in vitro exposure.

MATERIALS AND METHODS

Ten 10 ml tubes of blood were drawn from a normal healthy subject. Pairs of tubes contained the following anticoagulants: sodium heparin, sodium oxalate, potassium oxalate/ sodium fluoride, EDTA, and sodium citrate. As a control, needle-tip smears were made using no anticoagulant.

One tube from each of the five pairs was held at room temperature (24°C) and the other tube of each pair was held at 4°C, the temperatures at which blood is commonly kept in a clinical laboratory. Blood smears from the anticoagulated blood were made at 0.5, 1, 2, 3, 4, and 14 hours. The smears were air dried, fixed for one minute in absolute methanol, and stained with Wright's Stain on an Ames Hematek slide stainer. The examination of these and all subsequent smears was performed by bright-field microscopy at 1000x.

The smears were examined primarily for vacuole formation but color, cell morphology, and presence of other artifacts was noted.

RESULTS

Smears made from sodium citrate anticoagulated blood showed normal white cell morphology through 14 hours. Slight crenation of erythrocytes was seen at 14 hours both at 4°C and at 24°C. The EDTA anticoagulated smears showed normal morphology through 4 hours. After 4 hours, toxic granulation, a coarse dark granulation of the cytoplasm commonly seen in toxic conditions, occurred in blood stored at 24°C but not in blood stored at 4°C (Figure 3). Again, red cell crenation was seen at 14 hours at both temperatures. Heparinized smears had 1% vacuolated neutrophils at 1, 3, and 14 hours at 4°C and at 3 hours at 22°C (Figure 4). Potassium oxalate/ sodium fluoride smears displayed

Figure 3. Neutrophil showing toxic granulation at 4 hours. EDTA used as anticoagulant. 2300X.

Figure 4. Neutrophil showing slight vacuolation at 1 hour. Heparin used as anticoagulant. 1900X.

erythrocyte crenation at 1 hour at both temperatures. Heparinized blood yielded smears with slight toxic granulation of the neutrophils at 4 hours at room temperature and at 14 hours at 4°C. Sodium oxalate smears had toxic granulation at 3 hours at 24°C and at 4°C. Twenty percent of the neutrophils on the 14 hour 24°C slide had hypersegmented nuclei and one neutrophil on the 14 hour 4°C slide had vacuolated cytoplasm. Some of these results are summarized in Table I.

TABLE I

OCCURRENCE OF VACUOLATION IN ANTICOAGULATED BLOOD

ANTICOAGULANT -----	TIME AT 4°C						TIME AT 22°C						
	0.5	1	2	3	4	14	0.5	1	2	3	4	1	
EDTA	-	-	-	-	-	-	-	-	-	-	-	-	
HEPARIN	-	+	-	+	-	+	-	-	-	+	-	-	
SODIUM CITRATE	-	-	-	-	-	-	-	-	-	-	-	-	
POTASSIUM OXALATE/ SODIUM FLUORIDE	-	-	-	-	-	-	-	-	-	-	-	-	
SODIUM OXALATE	-	-	-	+	-	+	-	-	-	-	-	-	

+ vacuolation of neutrophil cytoplasm observed

- vacuolation of neutrophil cytoplasm not observed

DISCUSSION

In no instance did gross morphological alterations appear prior to one hour. It is important that with the most commonly used anticoagulant, EDTA, no serious changes

in morphology were noted prior to four hours. Vacuolation was seen only in the presence of heparin or sodium oxalate.

Providing that the blood is maintained at 4°C, that the anticoagulant used is EDTA or sodium citrate, and that the smears are made within four hours of the time the blood is drawn, smears made from anticoagulated blood are perfectly suitable for laboratory examination. One can be certain that vacuolation, if present, is not an artifact due to anticoagulant.

CHAPTER II

INTRODUCTION

This second study was directed towards determining the existence of any feature common to individuals having vacuolated neutrophils. In addition, I felt that it should be possible to determine something about the development of vacuolation through examination of blood smears. Specifically, one would expect to see at least equal numbers of vacuolated bands and vacuolated segmented forms if vacuolation is a maturational disturbance. If it is not a maturational disturbance, the predominant vacuolated form most likely would be the older segmented neutrophil.

MATERIALS AND METHODS

Local hospitals cooperated by watching for patients with cytoplasmic vacuolation and by providing background information and Wright's stained blood smears. I drew blood daily for ten days from four healthy volunteers. Despite the fact that some vacuolation could be attributed to heparin treatment, smears were made from both heparinized and non-anticoagulated blood. The use of heparin was necessary because blood was obtained by finger stick and only capillary tubes containing heparin were available. For comparison smears were also made by touching the finger directly to the

slides. In addition twelve healthy volunteers were sampled by finger-stick and smears were made directly from the finger tip. All smears, heparinized and non-heparinized, were made immediately to prevent the possibility of vacuole formation caused by technique. Each peripheral smear was examined at 1000x and 100 neutrophils were counted. They were scored for maturity and presence of vacuoles.

Patient information included age, sex, diagnosis, causative agent, differential white cell count, total white cell count, erythrocyte count, hemoglobin concentration, hematocrit, mean corpuscular volume, mean corpuscular hemoglobin, and mean corpuscular hemoglobin concentration. Not all information was available in all cases.

The strength of the relationship between percent vacuolation (cells exhibiting vacuolated cytoplasm per 100 cells examined) and the other variables was measured using an SPSS multiple regression package on Portland State University's Honeywell computer.

RESULTS

One of the participating hospitals averaged 1412 monthly admissions and supplied me with an average of 2 patients per month displaying vacuolation. This amounts to only 0.14% of the entering hospital population. This number should be considered the maximum occurrence of vacuolation in the hospital population since blood smears are scanned

not only from newly admitted patients but also from resident patients and from patients who pass through the emergency room but are not admitted.

No difference was noted between smears made from heparinized or non-heparinized blood. Three of the four healthy individuals showed no neutrophil vacuolation on any of the ten days. The fourth individual had 1% vacuolation on four of the ten days. None of the twelve one-time volunteers showed vacuolation on a peripheral smear.

The fifty-nine hospital inpatients who were selected for presence of vacuolation had a mean percent vacuolation of 17.6% (segmented neutrophils 8.1%, band neutrophils 10.1%). As can be seen from the data in Figure 5, the most common percent vacuolation was in the 0-5% range.

Final diagnosis was available for forty-seven of the fifty-nine inpatients. Fifteen subjects (32%) had septicemia, nineteen (40%) had local infections, and thirteen (28%) had no demonstrable infection (Table II). Nine (19%) of the patients with vacuolation (infected and non-infected) were known ethanol abusers. Included in this group were the only two cases of septicemia due to enteric bacteria.

Figure 5. Frequency of occurrence versus percent vacuolation for hospital inpatients.

VACUOLATED NEUTROPHILS PER
100 NEUTROPHILS COUNTED

TABLE II
DISTRIBUTION BY DIAGNOSIS, AGE, AND SEX
OF INPATIENTS DISPLAYING VACUOLATION

DIAGNOSIS	MALES	MEAN AGE	FEMALES	MEAN AGE
SEPTICEMIA 15 cases: (3 ethanol abusers)	4	22	11	48
Cause:				
<u>H. influenzae</u>	3		2	
<u>S. pneumoniae</u>	1		3	
<u>E. coli</u>			4	
<u>N. meningitidis</u>			2	
<u>S. aureus</u>			3	
Unknown			1	
LOCAL INFECTION 19 cases: (3 ethanol abusers)	3	66	16	52
Cause:				
<u>H. influenzae</u>	1		2	
<u>S. pneumoniae</u>	2		2	
<u>E. coli</u>	2		8	
<u>S. aureus</u>			3	
fungi			1	
Unknown			3	
NO INFECTION 13 cases: (1 ethanol abuser)	6	40	7	63
Ailment:				
Lung	1		2	
Heart	1		1	
Stroke	3		1	
Cancer	1		2	
Nephritis			1	
UNKNOWN 12 cases:	4	61	8	62
TOTAL 59 cases:	19	46	40	52

ETHANOL ABUSERS 7 cases:	3	57	4	50

DISCUSSION

As noted earlier not all information was available on all patients. Access to patient records was understandably limited but this limitation did make the study more difficult and the conclusions drawn less certain. It would have been very helpful if such information as time of onset of the illness, antibiotics taken, date therapy was begun, complications, and final resolution of the illness had been available. Certainly all of this information could have been made available without compromising patient privacy. For patients currently showing cytoplasmic vacuolation smears should be made every two or three hours to track the course of the vacuolation.

Although no attempt was made to quantify the degree of vacuolation within an individual cell it was noted that vacuolation of the neutrophil cytoplasm was usually limited to a few small (1.0 micron) vacuoles. Figure 1, from the admission blood smear of an affected individual, is typical. On occasion, vacuoles were of much greater size (Figure 6) or present in greater numbers (Figure 7).

Systemic and local infections taken together accounted for 72% of the patients for whom diagnosis was known. This was anticipated since activation of neutrophils by bacterial products preceeding phagocytosis is known to be a cause of cytoplasmic vacuolation (Bigley, personal communication). It is not clear what the cause of vacuolation was in the 13

Figure 6. Mature neutrophil with large vacuole. 2300X.

Figure 7. Mature neutrophils with many small cytoplasmic vacuoles. 2300X.

cases who had non-infectious diseases. There was no observable qualitative difference in the vacuolation nor was there any observable difference in persistence of vacuolation. Table III displays the sex differences noted. Twice as many female as male patients were found to have vacuolation although the percent vacuolation was very similar between sexes.

Although the hospital population does have more women (61%) than men (39%) most of this disparity is due to a large number of obstetrics patients, none of whom are a part of our vacuolation group. The numbers of men and women admitted for treatment of an illness is approximately equal.

TABLE III

SEX EFFECT OF NEUTROPHIL VACUOLATION IN INPATIENTS

SEX	N	% VACUOLATION (vacuolated cells per 100 cells examined)
---	---	-----
MALE	20	8.4
FEMALE	40	7.8

A large amount of information had been collected on each of the fifty-nine patients. To find out if there was any relationship between these variables and the observed percent vacuolation a multiple regression program which is part of the SPSS statistical package was run. Correlation

coefficients of the independent variables with percent vacuolation are found in Table IV. A more complete list is in the Appendix. Since percent vacuolation is not a normally distributed variable (Figure 5) the skewness of its distribution makes the observed values inaccurate estimators of the population mean (Snedecor and Cochran 1968). To correct for the skewed distribution it was necessary to perform a transformation, using the square root of the arcsine of percent vacuolation as the dependent variable in the regression (Clarkson, personal communication).

Some of the results of the regression are shown in Table V. The letter r stands for the regression coefficient. It is a measure of the correlation between percent vacuolation and the variable listed. The notation r^2 represents the proportion of the variance of percent vacuolation which is accounted for by the listed variable. For example, 9% of the variance in percent vacuolation may be attributed to the combined variable patient age and presence of local infection. It is important to note that the r and r^2 columns display cumulative values. Therefore, for the fifteen variables listed in the table, the cumulative contribution is to account for 62% of the variance in percent vacuolation. Although thirty of the sixty-three original variables and combinations were significant at the 0.05 level, I listed only fifteen in the

TABLE IV

CORRELATION COEFFICIENTS FOR INDEPENDENT AND
CALCULATED VARIABLES

VARIABLE	CORRELATION COEFFICIENT
ETHANOL ABUSE	0.22
<u>Neisseria meningitidis</u>	0.16
LOCAL INFECTION	0.16
SEPTICEMIA	0.11
MENINGITIS	0.11
SYSTEMIC INFECTION	0.10
SEX	0.08
AGE	0.02
<u>Hemophilus influenzae</u>	0.01
WHITE BLOOD COUNT	-0.12
AGE X LOCAL INFECTION	0.30
SYSTEMIC X LOCAL INFECTION	-0.19
ETHANOL ABUSE X SYSTEMIC INFECTION	0.17
AGE X WHITE BLOOD COUNT	-0.07
LOCAL INFECTION X <u>H. influenzae</u>	0.01
AGE X SYSTEMIC INFECTION	-0.16
WHITE BLOOD COUNT X LOCAL INFECTION	0.08
WHITE COUNT X ETHANOL ABUSE	0.18
<u>Neisseria meningitidis</u> X SEPTICEMIA	0.23
AGE X SEX	0.08
SEX X ETHANOL ABUSE	0.10
AGE X <u>N. meningitidis</u>	0.23

TABLE V

VARIABLES IN THE REGRESSION EQUATION

VARIABLE	CUM. r	CUM. r ²	F	REJECTION REGION FOR F (.05)
-----	-----	-----	-----	-----
PATIENT AGE X LOCAL INFECTION	.30	.09	5.5	≤4.00
SYSTEMIC X LOCAL INFECTION	.41	.17	5.5	≤3.15
ETHANOL X SYSTEMIC INFECTION	.50	.25	6.1	≤2.76
AGE X WBC	.55	.30	5.8	≤2.53
LOCAL INFECTION	.58	.33	5.3	≤2.37
LOCAL INFECTION X <u>H. influenzae</u>	.62	.38	5.4	≤2.25
AGE X SYSTEMIC INFECTION	.65	.42	5.4	≤2.17
WBC X LOCAL INFECTION	.67	.45	5.2	≤2.10
WBC X ETHANOL ABUSE	.69	.48	5.1	≤2.04
ETHANOL ABUSE	.72	.52	5.1	≤1.99
<u>N. meningitidis</u> X SEPTICEMIA	.74	.55	5.2	
AGE X SEX	.76	.57	5.2	≤1.92
SEX X ETHANOL ABUSE	.77	.59	5.0	
AGE	.78	.60	4.8	
AGE X <u>N. meningitidis</u>	.78	.62	4.6	≤1.84

table. The reason for this is that beyond the fifteenth step in the regression the changes seen in r^2 were very small, all less than 0.01. Fifteen variables taken together accounted for no more than 62% of the variance in percent vacuolation. In fact, if all thirty variables were included still only 70% of the variance would be explained. The fact that so many variables are involved in percent vacuolation shows that the phenomenon very likely has a number of contributing causes. Specifically, vacuolation appears to be related primarily to increasing patient age, excessive ethanol use, and presence of bacterial infection. Of course the finding of an association does not prove cause. From the data it is clear that an increased probability of occurrence of vacuolation is correlated with increasing age of the patient, with increasing white blood count (specifically neutrophils), with presence of infection, and with history of ethanol abuse. Seventy-two percent of the patients for whom diagnosis was known were known to have a concurrent infection (Table II). One could predict then, in an older patient showing vacuolation on an admission blood smear, a high probability of infection. This is especially true if there is a history of ethanol abuse.

It is worth noting that the healthy volunteer who had 1% vacuolation on several days is both female and is over fifty years of age while the other three volunteers were all under forty. This is consistent with the age and sex fac-

tors we have noted above. Taken together there were sixteen healthy volunteers only one of whom showed any degree of vacuolation.

The percent vacuolation on an admission blood smear does not correlate strongly with either local infection (0.16) or systemic infection (0.10). However, in light of presence of vacuolation on the smear, my finding that 72% of these individuals are eventually confirmed as having an infection (32% septicemia, 40% local infection), and the fact that only 0.14% of the more than 33,000 patients entering this hospital in a two year period were found to have vacuolation, the presence of vacuolation, regardless of percentage, should be regarded as early presumptive evidence of a possibly dangerous infection.

CHAPTER III

INTRODUCTION

Due in part to the difficulty of obtaining daily specimens from hospital patients and in part to the lack of control over such variables as age, sex, type of disease, antibiotics administered, and other treatments, I felt that an animal model would be helpful. Mice were chosen for ease in obtaining, bleeding, and maintaining.

There are some notable hematologic differences between mice and humans. Differential counts of human peripheral leukocytes normally show 50 to 70% segmented neutrophils (Davidson and Henry 1969). Segmented neutrophils usually represent only 8 to 15% of the circulating leukocytes in mice (Hardy 1967). According to Hardy (1967), segmented neutrophil percentages in Swiss mice have been reported to have a mean of 58%.

In this study an attempt was made to induce cytoplasmic vacuolation in mice. Examination of patient histories reveal a high incidence of bacterial infection, ethanol abuse or a combination of the two in humans having cytoplasmic vacuolation. A positive relationship between the history of infection and ethanol abuse and the degree of vacuolation can be shown (Chapter II). Therefore attempts were made to induce vacuolation in mice with bacteria and

ethanol.

MATERIALS AND METHODS

Forty Swiss outbred mice (twenty females and twenty males) were obtained from Simonsen Laboratories. The mice were divided into four groups, each having five females and five males. To minimize fighting, males and females of each group were kept in separate cages. Within each cage the mice were marked with a felt pen for individual identification. Blood was drawn from all mice immediately before beginning the experiment using the rapid orbital bleeding technique of Riley (1960).

All forty mice were maintained on water ad libitum and Purina Rodent Chow except for the following experimental deviations: a control group of ten received water and Rodent Chow; a second group of ten had as their only source of liquid 10% ethanol in water; a third group was injected intraperitoneally with 3×10^8 heat killed Escherichia coli; and a fourth group had 10% ethanol as their only liquid and were injected with E. coli.

The bacterial suspension was prepared as follows: a loopful of E. coli was introduced into 20 ml of Luria broth and incubated at 37°C in a shaking incubator until a Klett reading of 70 was obtained (3×10^8 bacteria per ml). To ensure that the mice would survive during the experiment, the bacteria were killed by boiling the suspension for ten

minutes. McCall (1969 and 1979) and Bigley (personal communication) claim that the vacuolation was due to the presence of bacterial products and not the bacteria themselves so I felt it was not necessary to expose the mice to live bacteria at this time. Aliquots of the boiled suspension were plated on nutrient agar to assure that the bacteria were killed. The remaining suspension was spun at 7000 RPM for ten minutes to sediment the cells, the broth was poured off and the cells were resuspended in 0.85% saline. The cells were washed twice more, suspended in sterile 0.85% saline and 1.0 ml aliquots were injected into the mice in the bacteria and the bacteria/ ethanol groups.

All mice were bled daily for 14 days. The blood was collected in VWR micro-hematocrit capillary tubes which contained, as anticoagulant, 2.0 USP units of ammonium heparin. Although heparin has been shown to be correlated with vacuolation (Chapter I) it has only had that effect on white cells after a minimum one hour exposure at 4°C. Therefore care was taken that the blood was immediately drawn, smeared and air dried. There was neither a prolonged exposure to the heparin solution nor to reduced temperature. Blood smears were made on microscope slides, fixed for 60 seconds in absolute methanol, and subsequently stained with Wright's stain in an Ames Hematek slide stainer. Slides were examined in brightfield at 1000x for gross morphology. Fifty neutrophils were counted per slide and were scored for

maturity and presence of vacuoles.

RESULTS

The initial blood smears from the mice, showed vacuolation in only three mice, two males and one female. Figure 8 shows a normal blood smear with no evidence of vacuolation. The level of vacuolation was low; 2 to 4 percent.

The reactions of the bacteria, ethanol, and control group mice were very similar with mean percent vacuulations of 9.9%, 9.4%, and 8.1% respectively. The mice exposed to ethanol and bacteria however, had a mean percent vacuolation of only 4.4%. Although the variability within groups is very large it is roughly the same between groups. Males had consistently higher percent vacuolation in every experimental group than did females (Figure 9). The difference was one of degree not of occurrence since all but one of the forty mice developed vacuolation at some time during the experiment. The mean percent vacuolation for males was approximately twice that of the females. In humans sex was a factor in occurrence of vacuolation but not in percent affected cells.

Concurrent with vacuolation in 42% of the occurrences was a peculiarity of the plasma which resulted in smears having a bubbly background appearance. This feature may be seen in Figure 10. The presence of this bubbly, distorted background on the slide was far more common for the male

Figure 8. Normal mature neutrophil from a mouse.
2300X.

Figure 9. Mean percent vacuolation for mice by sex and experimental group.

Figure 10. Blood smear with distorted background.
2300X.

CRANES (S) CREST

than for the female mice (Figure 11). Comparison of Figures 9 and 11 reveal a high correlation between the disrupted background, vacuolation, and sex in mice. This phenomenon was never seen in the peripheral blood smears of humans. Those mice which had greater than 50% vacuolation frequently had neutrophils whose cytoplasm had a frothy appearance, showing such a mass of vacuoles that the nuclei were distorted (Figure 12). This feature also was never observed in blood smears from human subjects.

DISCUSSION

Since the control animals had a vacuolation response equivalent to that of the experimental groups the bacterial injections and/or ethanol feedings were evidently not the sole cause of the vacuolation. It is more likely that tissue necrosis due to daily orbital bleeding was the causative agent.

One curiosity is the lower levels of vacuolation noted in the ethanol/bacteria group. It appears as if ethanol consumption has in some way interfered with or modified the process of vacuole formation. Whether this is due to a reduction in the stress sensed by the animal, or due to a metabolic effect is not known. It is also not known why there is such a disparity between males and females in all experimental groups. Despite far higher percentages of vacuolation in males there was no difference in frequency of

occurrence of vacuolation between sexes. Human subjects, whether male or female, had nearly the same percent vacuolation but there were differences in occurrence between the sexes, with female subjects being affected twice as frequently.

Figure 11. Occurrence of disrupted background by sex and experimental group.

Figure 12. Mouse neutrophil grossly distorted by cytoplasmic vacuoles. 2300X.

CHAPTER IV

INTRODUCTION

Given the occurrence of vacuolated neutrophils in experimental animals it would be valuable to know whether the vacuolation is induced by bacterial injection, ethanol ingestion, or by some other stimulus.

During phagocytosis neutrophils experience a burst of aerobic respiration (Karnovsky 1962) coupled with an increase in hydrogen peroxide production (Karnovsky 1962). At this time nitroblue tetrazolium (NBT) can be reduced within the phagocytic vacuole to blue formazan by the action of NADH oxidase (Miller and Kaplan 1970). Nitroblue tetrazolium enters the neutrophil by precipitating with heparin or fibrinogen and being phagocytosed (Segal and Jones 1980). A low percentage of neutrophils from healthy individuals (6.6%) (Jongeling 1975), and those from patients with non-bacterial illnesses (5.8 to 9.5%) (Park et al 1968) give a positive reaction with NBT. Cells which have been "activated" by exposure to bacteria or their products give a positive reaction with NBT although no evidence of phagocytosis may be seen (Bigley personal communication). The percentage of positives is 29% to 47% in cases of bacterial infection (Park et al 1968) and might therefore be useful in distinguishing between cells vacuolating due to exposure to

bacterial products and cells reacting to tissue necrosis (Bigley personal communication).

MATERIALS AND METHODS

Two runs were performed. In each run twelve Swiss outbred mice were divided into four groups of three animals each. They were maintained on the same experimental regimen as the groups of mice in Chapter III, except that the E. coli used in run number one were viable, not heat killed. The reason for using viable bacteria was to see if there was a difference in the response of the mice depending on the condition of the bacteria. The bacteria used in run number two were heat killed. Peripheral blood smears were made daily for each animal and stained with giemsa (Davidson and Henry 1962). These smears were examined for presence of vacuolation. The nitroblue tetrazolium reduction test was also performed daily as outlined in Hennig (1979) with the following modifications: 1) the blood was drawn using heparinized microhematocrit capillary tubes, and 2) phorbol 12-myristate 13-acetate was omitted from the procedure and an equal volume of 0.85% NaCl was substituted. Slides were prepared and the cells were scored as to positive or negative NBT reaction according to the criteria of Jongeling (1975). Phorbol 12-myristate 13-acetate causes activation of the neutrophils and was used prior to the beginning of the experiment with blood from each mouse to establish the

maximal NBT response for the mice. The inclusion of phorbol 12-myristate 13-acetate in the original procedure was for the purpose of finding those individuals who have a deficit in bacterial killing capacity (Hennig 1979).

RESULTS

The results for runs number one and two are shown in Tables VI and VII respectively. The data listed under day one for both tables were acquired the day before the experimental protocol was instituted. On day one of the first run three animals showed vacuolation and two had a positive NBT test. On day one in run number two, all animals were free from cytoplasmic vacuolation and all were negative for NBT reduction. However, the ability of mouse neutrophils to react to NBT was demonstrated by the strong positive reactions obtained when the cells were stimulated with phorbol 12-myristate 13-acetate (Figure 13). All mice tested with phorbol 12-myristate 13-acetate showed a positive NBT reaction.

Animals whose cells became positive for NBT reduction during the course of the experiment showed a much weaker reaction (Figure 14). Ten of twelve animals in run number one developed a positive NBT reaction at some time during the experiment. All of the animals in run number 2 developed a positive NBT and for ten of them the reaction remained positive for the balance of the experiment.

Experimental animals developed a positive NBT reaction on the average one day sooner than control animals. Vacuolation was more prevalent early in controls and later in non-controls.

The correlation coefficient for the vacuolation/NBT relationship was only 0.23.

TABLE VI

PRESENCE OF VACUOLATION AND POSITIVE NITROBLUE TETRAZOLIUM TESTS IN FOUR GROUPS OF MICE
RUN NUMBER ONE

DAY	CONTROLS			ETHANOL			LIVE BACTERIA			ETHANOL / LIVE BACT.		
	1	2	3	4	5	6	7	8	9	10	11	12
1 VAC	-	-	+	-	-	-	+	+	-	-	-	-
1 NBT	-	-	+	-	-	-	+	-	-	-	-	-
2 VAC	-	-	+	-	-	-	+	+	+	-	-	-
2 NBT	-	-	+	-	-	-	+	+	-	-	-	-
3 VAC	-	-	-	-	-	+	D	+	D	+	-	-
3 NBT	-	-	-	-	-	-		+		-	-	-
4 VAC	-	+	-	+	-	+		D		-	+	-
4 NBT	-	-	-	-	-	-				-	-	-
5 VAC	-	+	+	-	-	+				+	-	-
5 NBT	-	-	-	-	-	-				+	-	-
6 VAC	-	+	-	+	-	-				-	-	-
6 NBT	-	+	-	-	-	-				+	+	+
7 VAC	-	-	-	-	-	-				-	-	-
7 NBT	-	-	+	+	-	+				+	+	+

+ = positive NBT test or presence of vacuolation

- = negative NBT test or absence of vacuolation

D = deceased

TABLE VII

PRESENCE OF VACUOLATION AND POSITIVE NITROBLUE TETRAZOLIUM
TESTS IN FOUR GROUPS OF MICE
RUN NUMBER TWO

DAY	CONTROLS			ETHANOL			KILLED BACTERIA			ETHANOL/ KILLED BACT.		
	1	2	3	4	5	6	7	8	9	10	11	12
1 VAC	-	-	-	-	-	-	-	-	-	-	-	-
NBT	-	-	-	-	-	-	-	-	-	-	-	-
2 VAC	-	+	+	-	-	+	-	-	-	+	-	-
NBT	-	-	-	-	+	-	-	+	+	-	+	-
3 VAC	-	-	-	-	-	-	-	-	-	+	+	-
NBT	-	-	+	+	+	+	+	+	+	+	+	+
4 VAC	+	-	-	-	-	+	+	-	-	-	-	-
NBT	+	+	+	+	+	+	+	+	+	-	+	+
5 VAC	-	+	-	-	-	-	+	-	-	+	+	-
NBT	+	+	+	+	-	+	+	+	+	+	+	+
6 VAC	-	-	-	+	-	-	-	-	-	+	-	+
NBT	+	+	+	+	+	+	+	+	+	+	+	+

+ = positive NBT test or presence of vacuolation

- = negative NBT test or absence of vacuolation

DISCUSSION

Due to the appearance of vacuoles and positive NBT reactions in control animals it is clear that the ETOH and bacteria were not the only stimuli to which these animals were reacting. The expected results were positive NBT reactions and the presence of vacuolation in experimental animals and negative NBT reactions and the absence of vacuolation in control animals. However, not only did control animals have positive NBT reactions and vacuolation

Figure 13. Positive NBT test stimulated by phorbol 12-myristate 13-acetate. 2300X.

Figure 14. Positive NBT test stimulated experimentally. 2300X.

but there was little correlation between the vacuolation and NBT responses.

Animals exposed to viable bacteria only succumbed to their infections within three days. For this reason this discussion will be primarily concerned with the animals in run #2.

The non-control animals did respond with a positive NBT reaction on the average one day earlier and it is probable that this enhanced response is the differential amount which may be attributed to the action of the ethanol and/or bacteria. The fact that all animals ultimately responded can be attributed to the tissue irritation and destruction which is unavoidable given daily bleeding which causes tissue necrosis, encourages eye rubbing by the animals and results in local infection. There was very little difference between groups (run #2) in number of days in which a positive NBT test result was seen (control= 10 days, ETOH= 12 days, Bacteria= 14 days, ETOH/Bacteria= 12 days).

The vacuolation response was less clear cut. The appearance of vacuolation averaged one day earlier in controls than in experimentals. Among the animals given viable bacteria only (run #1), vacuolation appeared within the first two days.

The highest incidence of vacuolation was among the members of the ETOH/Bacteria group (run #2). It is in this respect that the results of the two runs are in conflict.

The members of the ETOH/Bacteria group in run #1 had the lowest incidence of vacuolation. On the one hand ETOH seems to confer a resistance to the effects of bacterial injection, on the other hand it seems to have a synergistic effect. I believe the difference is dependent on whether or not the bacteria used are viable. The resistance effect was seen only in the case of live bacteria. This gives the impression that the growth of bacteria was inhibited in the animals receiving ETOH. It is especially interesting that the animals which received both viable bacteria and ETOH survived the entire course of the experiment with no apparent ill effects while those receiving only the bacteria died early in the experiment. In the case of animals exposed to heat killed bacteria and ETOH there was no possibility for inhibition of the bacteria and the injurious effects of the bacteria and the ETOH were summed.

The low correlation between presence of vacuolation and a positive NBT test suggests that the vacuoles seen are not phagocytic vacuoles but instead are the result of tissue necrosis due to the blood drawing technique.

CHAPTER V

INTRODUCTION

McCall et al (1969, 1979) believed the stimuli precipitating cytoplasmic vacuolation were circulating bacterial chemotoxins. It seems reasonable that whatever the factor is that stimulates neutrophil vacuolation it may be present in the plasma of the affected individuals, although in light of the results of Chapter IV the nature of this factor is in question. As a simple test of this hypothesis I induced cytoplasmic vacuolation in normal neutrophils, in vitro, by exposure to plasma from affected individuals.

MATERIALS AND METHODS

Blood was collected in tubes containing sodium heparin from four patients displaying vacuolation. The tubes were centrifuged to sediment erythrocytes and the plasma was drawn off and frozen. Plasma was maintained at -20°C until tested. Each sample was typed against types A₁ and B reagent blood cells to determine ABO blood group. Since in each of the four cases the individuals were type O, cells were obtained from a healthy volunteer (#1) who was also type O. Blood was collected in heparinized Vacutainer tubes as for the patients above. The type O cells were sedimented by centrifugation, the plasma was removed, the red cells and

buffy coat were mixed, and the cells were stored at 4°C until used. In addition, plasma from this volunteer was frozen to use in control #1. Control #2 used plasma taken from a healthy volunteer having type A blood. Plasma was chosen over serum to ensure that no factors which might be necessary to the vacuolation process were excluded.

Vacuolation of neutrophils held in wet mounts has been reported (Gordin 1950, Bessis 1972). To minimize any effect due to evaporation or surface contact the use of thin blood films on microscope slides was avoided.

Five 10 ml culture tubes were used in the experiment. Each experimental tube received 0.5 ml of cells from volunteer #1 and 0.5 ml of plasma from one of the patients. Control tube #1 contained cells and plasma from volunteer #1 (type O). Control tube #2 contained cells from volunteer #1 (type O) and serum from volunteer #2 (type A). Smears were made immediately from each mixture, the remainder was incubated at 37°C in a CO₂ atmosphere. Smears were made from the mixtures at intervals of 0.5, 1, 2, 3, 4, 8, and 16 hours, air dried, fixed in absolute methanol for one minute, stained with giemsa for twenty minutes, and finally examined at 1000x for presence of vacuoles in the cytoplasm of the neutrophils.

RESULTS

Vacuolation was in evidence in the experimental groups

early in the experiment, prior to hour 4. After that time the degree of vacuolation in the control tubes rapidly increased and ultimately equaled the vacuolation seen in the cells from most other tubes. These results are shown in Table VIII and in Figure 15.

TABLE VIII

PERCENT VACUOLATION OF NEUTROPHILS EXPOSED TO PATIENT PLASMA VERSUS TIME OF EXPOSURE

PATIENT	TIME (HOURS)						
	0.5	1	2	3	4	8	16
-----	---	--	--	--	--	--	--
#1	8	4	12	4	2	2	12
#2	18	16	14	14	8	10	20
#3	4	14	24	26	30	12	20
#4	6	2	10	4	14	0	50
CONTROL #1	0	6	4	8	12	0	44
CONTROL #2	2	4	4	14	18	16	50

DISCUSSION

It appears that the cells exposed to plasma from patients known to have vacuolation of neutrophils vacuolate sooner while those exposed to normal plasma develop vacuolation at a later time. The ultimate level of vacuolation was just as high in the control tubes as in the experimental tubes, only the time frame was altered.

The differences between the experimental and control

Figure 15. Percent vacuolation versus time for neutrophils
in vitro.

groups are given seen in Table VIII. Not until the fourth hour did the vacuolation of the controls exceed that of the experimentals. The effects of drying and surface contact must be considered when looking at the prolonged incubation times. Gordin (1952) observed vacuole development in neutrophils left in wet mounts for 6-8 hours. This causes some doubt as to the applicability of results compiled after this time. The process of vacuolation seems to be limited in the time over which it can be sustained. Those tubes which showed early vacuolation, for the most part, did not maintain that level of vacuolation and were overtaken by the tubes which showed a low initial level of vacuolation.

TABLE IX

HOURLY COMPARISON OF MEAN PERCENT VACUOLATION
OF EXPERIMENTALS VERSUS CONTROLS

	TIME (HOURS)						
	0.5	1	2	3	4	8	16
	---	--	--	--	--	--	--
EXPERIMENTAL	9	9	15	12	14	6	26
CONTROL	1	5	4	11	15	8	47

Some differences were apparent between experimentals. For example, one of the patients with the highest initial vacuolation was diagnosed as having a severely infected wound while one of the patients with the lowest initial vacuolation had a septicemia. It is possible that the neutrophils from the septicemic patient had had a relatively long and thorough exposure to bacterial products and were

unable to respond to the same degree as the cells from a patient having a local infection and possibly a lesser degree of exposure of neutrophils to bacterial products. However, the cells from the septicemic patient were the only ones to have as high a percent vacuolation at 16 hours as the controls.

The decrease in vacuolation seen in tubes which displayed vacuolation early could be the result of two factors: "curing" of the affected cells or death and disintegration of the affected cells. There was no apparent decrease in numbers of neutrophils present on smears with increasing time into the experiment, nor was there any observable increase in cellular debris which would lead one to think that cells were being destroyed. As a result of these observations, I believe that the transient nature of cytoplasmic vacuolation in vivo is due not only to rapid migration of the cells out of the peripheral circulation but also due to these cells surviving the vacuolation and regaining a normal appearance.

Once again there is no clear difference between the experimentals and controls. The early vacuolation of cells in the patient tubes may be taken as an indication that there is some factor present inducing vacuolation. The delayed vacuolation of control tube #1 cannot be due only to incubation of cells and plasma from the same individual since delayed vacuolation was seen also in control tube #2

which had cells and plasma from different individuals and different blood groups.

FINAL DISCUSSION

The appearance of cytoplasmic vacuolation in neutrophils in the peripheral circulation is a phenomenon which has received relatively little study. In Chapter I, I was able to show that the appearance of the vacuoles, within certain limits, is not an artifact. Only on prolonged standing of the blood in the presence of anticoagulant is it possible to attribute some of the vacuolation to artifact formation.

The appearance of vacuoles on the peripheral blood smears of hospital inpatients was shown to be most strongly related to the presence of infection, particularly in female or aged patients. The value of a report of neutrophil vacuolation as a predictor of the presence of infection is probably quite good but the overall usefulness of vacuolation as a diagnostic tool is limited by the facts that it is highly transient and that it is detected in only a small fraction of patients having infection. Whether the lack of detection of vacuolation in patients having infection is due only to the short time within which it can be observed or is due to the fact that it does not invariably occur is not known.

The attempt to use mice as an animal model raised far more questions than it answered. It was possible to demonstrate the development of vacuolation in mice but the

vacuolation had a very different appearance and probably a very different cause. Specifically, the vacuolation seen is likely to be due to tissue necrosis of the tissues around the eye caused by daily ocular bleeding. This was borne out by the NBT test which showed that there was essentially no correlation between a positive NBT test (stimulation of vacuolation by presence of bacterial products) and the occurrence of vacuoles in the experimental animals. The observation that male mice had almost twice the percentage of vacuolated cells seen in females contrasted with the observation in humans that females, while having approximately the same percent vacuolated cells were twice as frequently affected. In addition, while in humans ethanol abuse was weakly correlated with vacuolation, in mice fed ethanol the occurrence of vacuolation seemed to be suppressed.

I was able to show that incubation of normal cells with the plasma from patients demonstrating vacuolation did induce vacuolation but the cause was ambiguous. Although the cells exposed to patient plasma became vacuolated first, ultimately all samples, both experimental and control, showed vacuolation.

The occurrence of vacuolation in experimental and control animals, of vacuolation in patients not known to have infection, the absence of vacuolation in cases of known infection, and the occurrence of vacuolation in cells exposed to normal plasma as well as plasma from affected

patients demonstrates that the phenomenon of vacuolation of neutrophil cytoplasm does not have one simple cause.

It is possible that further study could define the biochemical nature of the factors which cause the development of vacuolation but I am not enthusiastic about the benefits that would derive from such a study.

REFERENCES

- Beard, J. and D. Knott. 1966. "Hematopoietic Response to Experimental Chronic Alcoholism". The American Journal of the Medical Sciences, 252: 518-525.
- Belcour, L. and O. Begel. 1978. "Lethal Mitochondrial Genotypes in Podospora anserina: A Model for Senescence". Molecular and General Genetics, 163: 113-123.
- Bessis, M. 1972. Living Blood Cells and Their Ultrastructure. Berlin: Springer-Verlag.
- Boggs, D. and A. Winkelstein. 1981. White Cell Manual. Philadelphia: F.A. Davis Company.
- Cline, M. 1975. The White Cell. Cambridge: Harvard University Press.
- Davidson, I. and J. Henry. 1969. Todd-Sanford Clinical Diagnosis By Laboratory Methods 14th ed. Philadelphia: W.B. Saunders Co.
- Davidson, R. and J. McPhie. 1980. "Cytoplasmic Vacuolation of Peripheral Blood Cells in Acute Alcoholism". Journal of Clinical Pathology, 33: 1193-1196.
- Emerson, W., P. Zieve, and J. Krevans. 1970. "Hematologic Changes in Septicemia". The Johns Hopkins Medical Journal, 126: 69-76.
- Esser, K. and W. Keller. 1976. "Genes Inhibiting Senescence in the Ascomycete Podospora anserina". Molecular and General Genetics, 144: 107-110.
- Finch, C. and L. Hayflick. 1977. Handbook of The Biology of Aging. New York: Van Nostrand Reinhold Company.
- Gordin, R. 1952. "Toxic Granulation In Leukocytes Development and Relation to Cloudy Swelling". Acta Med Scand (Suppl 270), 143: 1-50.
- Grange, M., F. Brivet, P. Boumier and G. Tchernia. 1980. "Polynucleaires Neutrophiles Vacuoles Valeurs Diagnostique et Pronostique". La Nouvelle Presse Medicale, 9: 2553-2554.

- Greenwalt, T., ed. 1977. The Granulocyte: Function and Clinical Utilization. Alan R. Liss Inc.
- Hardy, J. 1967. "Hematology of Rats and Mice". Pathology of Laboratory Rats and Mice. Ed. E. Cotchin and F. Roe. Blackwell Scientific Publications.
- Hennig, C. 1979. "Nitroblue Tetrazolium Reduction Test, Hematology Division Procedure No.0550". Portland: University of Oregon Health Sciences Center, Department of Clinical Pathology.
- Jacques, Y. and D. Bainton. 1978. "Changes in pH within the Phagocytic Vacuoles of Human Neutrophils and Monocytes". Laboratory Investigation, 39: 179-185.
- Jongeling, G. 1975. "Standardization of the Qualitative Nitroblue Tetrazolium Dye Reduction Test and Its Clinical Usefulness". Thesis. University of Oregon Medical School.
- Karnovsky, M. 1962. "Metabolic Basis of Phagocytic Activity". Physiological Reviews, 42: 143-168.
- Lindenbaum, J. and C. Lieber. 1969. "Hematologic Effects of Alcohol in Man in the Absence of Nutritional Deficiency". The New England Journal of Medicine, 281: 333-338.
- Malcolm, I., K. Flegel, M. Katz. 1979. "Vacuolization of the Neutrophil in Bacteremia". Archives of Internal Medicine, 139: 675-676.
- McCall, C., D. Bass, L. DeChatelet, A. Link and M. Mann. 1979. "In Vitro Responses of Human Neutrophils to N-Formyl-Methionyl-Leucyl-Phenylalanine: Correlation with Effects of Acute Bacterial Infection". Journal of Infectious Diseases, 140: 277-286.
- McCall, C., I. Katayama, R. Cotran and M. Finland. 1969. "Lysosomal and Ultrastructural Changes in Human 'Toxic' Neutrophils During Bacterial Infection". Journal of Experimental Medicine, 129: 267-281.
- Meranze, D., T. Mendel and T. Meranze. 1935. American Journal of Medical Science, 189: 639.
- Miller, D. and H. Kaplan. 1970. "Decreased Nitroblue Tetrazolium Dye Reduction in the Phagocytes of Patients Receiving Prednisone". Pediatrics, 45: 861-865.

- Moxon, E., A. Smith, D. Averill and D. Smith. 1974. "Hemophilus influenzae Meningitis in Infant Rats after Intranasal Inoculation". Journal of Infectious Diseases, 129: 154-162.
- Nathan, D., R. Baehner and D. Weaver. 1969. "Failure of Nitroblue Tetrazolium Reduction in the Phagocytic vacuoles of Leukocytes in Chronic Granulomatous Disease". The Journal of Clinical Investigation, 48: 1895-1904.
- Park, B., S. Gikrig and E. Smithwick. 1968. "Infection and Nitroblue-Tetrazolium Reduction by Neutrophils". The Lancet, 7: 532-534.
- Parmley, R., W. Crist, H. Abdelsalam, L. Boxer, A. Malluh, V. Lui and C. Darby. 1980. "Congenital Dysgranulopoietic Neutropenia: Clinical, Serologic, Ultrastructural, and In Vitro Proliferative Characteristics". Blood, 56: 465-475.
- Ponder, E. and R. Ponder. 1942. "The Cytology of The Polymorphonuclear Leucocyte in Toxic Conditions". The Journal of Laboratory and Clinical Medicine, 28: 316-322.
- Riley, V. 1960. "Adaptation of Orbital Bleeding Technic to Rapid Serial Blood Studies". Proceedings of the Society for Experimental Biology and Medicine, 104: 751-754.
- Rubin, S. 1981. "Survival From Septicemia Depends On Rapid Detection and Therapy". Lab World, Apr: 33-35.
- Shafer, J. 1982. "Artifactual Alterations in Phagocytes in the Blood Smear". American Journal of Medical Technology, 48: 507-518.
- Segal, A. and O. Jones. 1980. "Rapid Incorporation of the Human Neutrophil Plasma Membrane Cytochrome B into Phagocytic Vacuoles". Biochem. Biophys Res. Commun., 92: 710-715.
- Snedecor, G. and W. Cochran eds. 1968. Statistical Methods, 6th ed., Ames, Iowa: The Iowa State University Press.
- Solberg, C. and K. Hellum. 1972. "Neutrophil Granulocyte Function in Bacterial Infections". The Lancet, Oct: 727-729.

- Stiene-Martin, E. 1982a. "Reactive Morphology in Blood Phagocytes: Introduction". American Journal of Medical Technology, 48: 485-490.
- Stiene-Martin, E. 1982b. "An Historical Review of Toxic and Reactive Morphologic Alterations in Granulocytes and Monocytes". American Journal of Medical Technology, 48: 491-505.
- Sutro, C. 1933. Archives of Internal Medicine, 51: 747. Referenced in: Gordin, R. 1952. "Toxic Granulation in Leukocytes, Development and Relation to Cloudy Swelling". Acta Med Scand (Suppl 270), 143: 1-50.
- Ward, P. 1980. "The Myeloid Leukocytoses". Postgraduate Medicine, 67(1): 219-229.
- Wintrobe, M., G. Lee, D. Boggs, J. Bithell, J. Foerster, J. Athens and J. Lukens eds. 1981. Clinical Hematology, 8thed., Philadelphia: Lea and Febiger.
- Zieve, P., M. Haghshenass, M. Blanks and J. Krevans. 1966. "Vacuolation of the Neutrophil". Archives of Internal Medicine, 118: 356-357.

APPENDIX

CORRELATION COEFFICIENTS

CORRELATION COEFFICIENTS

VARIABLES	(1)	(2)	(3)	(4)	(5)	(6)	(7)
VACUOLATED NEUTROPHILS (1)	1.0	.22	.10	.16	.01	.16	.11
ETHANOL ABUSE (2)	.22	1.0	.11	.08	-.12	.11	.15
SYSTEMIC INFECTION (3)	.10	.11	1.0	-.04	.41	.42	.92
LOCAL INFECTION (4)	.16	.08	-.04	1.0	.13	-.19	.01
<u>H. influenzae</u> (5)	.01	-.12	.41	.13	1.0	-.09	.45
<u>N. meningitidis</u> (6)	.16	.11	.42	-.19	-.09	1.0	.15
SEPTICEMIA (7)	.11	.15	.92	.01	.45	.15	1.0
MENINGITIS (8)	.11	.08	.48	-.21	.10	.64	.24
AGE (9)	.02	-.02	-.32	.04	-.34	-.25	-.20
SEX (10)	-.14	.08	-.04	-.16	.25	-.07	.05
WHITE BLOOD COUNT (11)	-.12	.17	-.11	.08	.06	-.15	-.08
AGE X LOCAL INF. (12)	.30	.03	-.09	.83	.01	-.15	-.05
SYSTEMIC X LOCAL INF. (13)	-.19	.08	.48	.44	.30	-.08	.52
ETHANOL X SYSTEMIC (14)	.17	.63	.36	.01	-.08	.24	.40
AGE X WHITE COUNT (15)	-.07	.06	-.22	.07	-.19	-.20	-.15
LOCAL INF. X <u>H. influenzae</u> (16)	.01	-.08	.19	.34	.69	-.06	.22
AGE X SYSTEMIC INF. (17)	-.16	.14	.73	.03	.16	.16	.79
WHITE COUNT X LOCAL (18)	.08	.05	-.11	.80	.16	-.15	-.07
WHITE COUNT X ETHANOL (19)	.18	.84	.01	.01	-.10	-.01	.03
<u>N. meningitidis</u> X SEPTICEMIA (20)	.23	.22	.29	-.13	-.06	.69	.32
AGE X SEX (21)	.08	.15	-.20	-.08	-.12	-.14	-.16
SEX X ETHANOL (22)	.10	.63	-.15	.17	-.08	-.06	-.14
AGE X <u>N. meningitidis</u> (23)	.23	.27	.29	-.13	-.06	.69	.31

CORRELATION COEFFICIENTS (CONT.)

VARIABLES		(8)	(9)	(10)	(11)	(12)	(13)	(14)
VACUOLATED NEUTROPHILS	(1)	.11	.02	.08	-.12	.30	-.19	.17
ETHANOL ABUSE	(2)	.08	-.02	.08	.17	.03	.08	.63
SYSTEMIC INFECTION	(3)	.48	-.32	-.04	-.11	-.09	.48	.36
LOCAL INFECTION	(4)	-.21	.04	-.16	.08	.83	.44	.01
<u>H. influenzae</u>	(5)	.10	-.34	.25	.06	.01	.30	-.06
<u>N. meningitidis</u>	(6)	.64	-.25	-.04	-.15	-.15	-.08	.24
SEPTICEMIA	(7)	.24	-.20	-.07	-.08	-.05	.52	.40
MENINGITIS	(8)	1.0	-.35	.05	-.05	-.17	-.09	.21
AGE	(9)	-.35	1.0	-.14	.15	.33	-.08	-.05
SEX	(10)	.05	-.14	1.0	.16	-.13	-.21	-.16
WHITE BLOOD COUNT	(11)	.05	.15	.16	1.0	.12	-.08	-.02
AGE X LOCAL INF.	(12)	-.17	.33	-.13	.12	1.0	.27	-.13
SYSTEMIC X LOCAL INF.	(13)	-.09	-.08	-.21	-.08	.27	1.0	.21
ETHANOL X SYSTEMIC	(14)	.21	-.05	-.16	-.02	-.13	.21	1.0
AGE X WHITE COUNT	(15)	-.25	.62	.04	.80	.29	-.10	-.05
LOCAL INF. X								
<u>H. influenzae</u>	(16)	-.07	-.13	.01	.11	.13	.48	-.05
AGE X SYSTEMIC INF.	(17)	.08	.12	-.16	-.02	.04	.27	-.13
WHITE COUNT X LOCAL	(18)	-.17	.08	-.10	.41	.71	.23	-.01
WHITE COUNT X ETHANOL	(19)	-.02	-.03	.12	.36	-.02	.03	.36
<u>N. meningitidis</u> X								
SEPTICEMIA	(20)	.28	-.03	-.13	-.11	-.11	-.06	.38
AGE X SEX	(21)	-.15	.32	.74	.22	.01	-.16	-.12
SEX X ETHANOL	(22)	-.07	.07	.34	.18	.22	-.07	-.05
AGE X <u>N. meningitidis</u>	(23)	.34	-.03	-.13	-.12	-.11	-.06	.45

CORRELATION COEFFICIENTS (CONT.)

VARIABLES	(15)	(16)	(17)	(18)	(19)	(20)	(21)
VACUOLATED NEUTROPHILS(1)	-.07	.01	-.16	.08	.18	.23	.08
ETHANOL ABUSE (2)	.06	-.08	.14	.05	.84	.22	.15
SYSTEMIC INFECTION (3)	-.22	.19	.73	-.11	.01	.29	-.20
LOCAL INFECTION (4)	.07	.34	.03	.80	.01	-.13	-.08
<u>H. influenzae</u> (5)	-.19	.69	.16	.16	-.10	-.06	-.12
<u>N. meningitidis</u> (6)	-.20	-.06	.16	-.15	-.01	.69	-.14
SEPTICEMIA (7)	-.15	.22	.79	-.07	.03	.32	-.16
MENINGITIS (8)	-.25	-.07	.08	-.17	-.02	.28	-.15
AGE (9)	.62	-.13	.12	.08	-.03	-.03	.32
SEX (10)	.04	.01	-.16	-.10	.12	-.13	.74
WHITE BLOOD COUNT (11)	.80	.11	-.02	.41	.36	-.11	.22
AGE X LOCAL INF. (12)	.29	.13	.04	.71	-.02	-.11	.01
SYSTEMIC X LOCAL INF. (13)	-.10	.48	.27	.23	.03	-.06	-.16
ETHANOL X SYSTEMIC (14)	-.05	-.05	-.13	-.01	.36	.38	-.12
AGE X WHITE COUNT (15)	1.0	-.06	.07	.34	.19	-.09	.36
LOCAL INF. X <u>H. influenzae</u> (16)	-.06	1.0	.25	.35	-.07	-.04	-.12
AGE X SYSTEMIC INF. (17)	.07	.25	1.0	-.05	.05	.30	-.08
WHITE COUNT X LOCAL (18)	.34	.35	-.05	1.0	.09	-.10	-.07
WHITE COUNT X ETHANOL (19)	.19	-.07	.05	.09	1.0	.04	.15
<u>N. meningitidis</u> X SEPTICEMIA (20)	-.09	-.04	.30	-.10	.04	1.0	-.09
AGE X SEX (21)	.36	-.12	-.08	-.07	.15	-.09	1.0
SEX X ETHANOL (22)	.14	-.05	-.11	.12	.63	-.04	.38
AGE X <u>N. meningitidis</u> (23)	-.09	-.04	.30	-.10	.06	.98	-.10

CORRELATION COEFFICIENTS (CONT.)

<u>VARIABLES</u>	(22)	(23)
VACUOLATED NEUTROPHILS(1)	.10	.23
ETHANOL ABUSE (2)	.63	.27
SYSTEMIC INFECTION (3)	-.15	.29
LOCAL INFECTION (4)	.17	-.13
<u>H. influenzae</u> (5)	-.08	-.06
<u>N. meningitidis</u> (6)	-.06	.69
SEPTICEMIA (7)	-.14	.31
MENINGITIS (8)	-.07	.34
AGE (9)	.07	-.03
SEX (10)	.34	-.13
WHITE BLOOD COUNT (11)	.18	-.12
AGE X LOCAL INF. (12)	.22	-.11
SYSTEMIC X LOCAL INF. (13)	-.07	-.06
ETHANOL X SYSTEMIC (14)	-.05	.45
AGE X WHITE COUNT (15)	.14	-.09
LOCAL INF. X <u>H. influenzae</u> (16)	-.05	-.04
AGE X SYSTEMIC INF. (17)	-.11	.30
WHITE COUNT X LOCAL (18)	.12	-.10
WHITE COUNT X ETHANOL (19)	.63	.06
<u>N. meningitidis</u> X SEPTICEMIA (20)	-.04	.98
AGE X SEX (21)	.38	-.10
SEX X ETHANOL (22)	1.0	-.04
AGE X <u>N. meningitidis</u> (23)	-.04	1.0