1985

The role of copper in the apparent aluminum toxicity of aquatic systems

Kirk Thomas O'Reilly
Portland State University

Follow this and additional works at: https://pdxscholar.library.pdx.edu/open_access_etds

Part of the [Biology Commons](#), and the [Plant Sciences Commons](#)

Let us know how access to this document benefits you.

Recommended Citation

O'Reilly, Kirk Thomas, "The role of copper in the apparent aluminum toxicity of aquatic systems" (1985).
Dissertations and Theses. Paper 3522.
<https://doi.org/10.15760/etd.5406>

This Thesis is brought to you for free and open access. It has been accepted for inclusion in Dissertations and Theses by an authorized administrator of PDXScholar. Please contact us if we can make this document more accessible: pdxscholar@pdx.edu.

AN ABSTRACT OF THE THESIS OF Kirk Thomas O'Reilly for the
Master of Science in Biology presented May 24, 1985.

Title: The role of copper in the apparent aluminum toxicity
of aquatic systems

APPROVED BY MEMBERS OF THE THESIS COMMITTEE:

[REDACTED]
John G. Rueter, Chairman

[REDACTED]
Richard R. Petersen

[REDACTED]
Byron E. Lippert

[REDACTED]
Eugene Enneking

The effect of variations in aluminum and copper concentrations on the growth rate and enzyme activity of the green alga Scenedesmus quadricauda was investigated. The goal was to determine which chemical species control the biological parameters. The computer program MINEQL (Westal et al 1976) was used to estimated chemical speciation. In the presences of both metals, algal growth rate and alkaline phosphatase activity could be correlated

to cupric ion activity. The activity of isolated bacterial alkaline phosphatase was found to be a function of both total copper concentration and cupric ion activity. A model was developed to predicted the effects on alkaline phosphatase of perturbations in aquatic chemistry.

This research has environmental significance because the concentration of aluminum increases in some lakes affected by acid rain. Aluminum is thought to be responsible for some of the observed toxic effect in these systems. The results of this research support the hypothesis that the effect of aluminum may be due to its interaction with the metal chemistry; the increase in aluminum leads to an increase in the toxic species of other metals such as copper.

THE ROLE OF COPPER IN THE APPARENT ALUMINUM
TOXICITY OF AQUATIC SYSTEMS

by

KIRK THOMAS O'REILLY

A thesis submitted in partial fulfillment of the
requirements for the degree of

MASTER OF SCIENCE
in
BIOLOGY

Portland State University

1985

TO THE OFFICE OF GRADUATE STUDIES AND RESEARCH:

The members of the Committee approve the thesis of
Kirk Thomas O'Reilly presented May 24, 1985.

John G. Rueter, Chairman

Richard R. Petersen

Byron E. Lippert

Eugene Enneking

APPROVED:

W. Herman Taylor, Head, Department of Biology

Jim E. Heath, Dean of Graduate Studies and Research

ACKNOWLEDGEMENT

I would like to thank the members of my committee; John Rueter who taught me how to think like a scientist, Richard Petersen who made Oregon lakes come alive, and Ed Lippert who introduced me to his friends the algae.

I also thank the members of my family, especially Brin and Ron, for their support, help, and suggestions during my nineteen plus years as a student.

E. Quentin Welch acted as my trusted assistant during the growth rate experiments described in Chapter II. I thank him for his help and friendship. I also appreciate the help Casson Stalling gave me in dealing with computers.

Finally, I would like to thank Kathryn Morrill for helping me keep sane by sharing my craziness.

TABLE OF CONTENTS

	PAGE
ACKNOWLEDGEMENTS.....	iii
LIST OF TABLES.....	vi
LIST OF FIGURES.....	vii
INTRODUCTION.....	1
CHAPTER I.....	3
BACKGROUND	
CHAPTER II.....	14
THE GROWTH RATE OF <u>SCENEDESMUS QUADRICAUDA</u> AS A FUNCTION OF ALUMINUM AND COPPER	
CHAPTER III.....	28
ALKALINE PHOSPHATASE ACTIVITY AS A FUNCTION OF THE AQUATIC CHEMISTRY OF ALUMINUM, COPPER AND ZINC	
CHAPTER IV.....	52
ALKALINE PHOSPHATASE ACTIVITY IN <u>SCENEDESMUS</u> : INTERACTIONS OF ALUMINUM AND COPPER	
CHAPTER V.....	64
THE USE OF A COMPUTER MODEL TO PREDICT THE INTERACTIONS OF COPPER AND ALKALINE PHOSPHATASE	
CHAPTER VI.....	78
CONCLUSION	

APPENDIX A.....	82
-----------------	----

COPPER AND LACTATE DEHYDROGENASE:
INHIBITION OF AN INTERNAL ENZYME

REFERENCES.....	89
-----------------	----

LIST OF TABLES

TABLE		PAGE
I	Modified FRAQUIL medium.....	16
II	Growth rate day ⁻¹ and pCu for concentration of aluminum and copper.....	19
III	Enzyme buffers tested.....	31
IV	Activity of alkaline phosphatase after twenty-four hours exposure to aluminum and copper.....	35
V	Chemical components, species, and stability constants used in MICROQL.....	37
VI	Activity of alkaline phosphatase at ten concentrations of aluminum and copper.....	38
VII	Reactivation of alkaline phosphatase after zinc addition.....	45
VIII	Activity of <u>Scenedesmus</u> alkaline phosphatase and cupric ion activity as a function of aluminum and copper concentration.....	57
IX	A comparison of experimental results with those predicted by MICROQL: zinc & pNPP	67
X	A comparison of experimental results with those predicted by MICROQL: copper.....	70

LIST OF FIGURES

FIGURES		PAGE
1.	The growth rate day ⁻¹ of <u>Scenedesmus quadricauda</u> as a function of aluminum and copper concentration.....	20
2.	The growth rate day ⁻¹ of <u>Scenedesmus quadricauda</u> as a function cupric ion activity (pCU).....	22
3.	The activity of the cupric ion as a function of total aluminum.....	24
4.	The growth rate day ⁻¹ of <u>Scenedesmus quadricauda</u> as a function of total aluminum at a set copper concentration.	26
5.	The activity of alkaline phosphatase in five buffers over twenty-four hours.....	32
6.	The activity of alkaline phosphatase as a function of aluminum concentration in the presence and absences of copper.....	40
7.	The activity of alkaline phosphatase as a function of cupric ion activity....	41
8.	A comparison of alkaline phosphatase activity at two zinc concentrations....	42
9.	Reactivation of alkaline phosphatase activity after zinc addition.....	46
10.	The activity of <u>Scenedesmus</u> alkaline phosphatase as a function of aluminum concentration in the presence and absences of copper.....	56
11.	The activity of <u>Scenedesmus</u> alkaline phosphatase as a function of copper concentration at four aluminum concentrations.....	58

LIST OF FIGURES
cont.

FIGURES		PAGE
12.	The activity of <u>Scenedesmus</u> alkaline phosphatase as a function of cupric ion activity.....	59
13.	The activity of the cupric ion as a function of total aluminum.....	61
14.	A comparison of experimental and predicted alkaline phosphatase activity at six zinc concentrations.....	72
15.	A comparison of experimental and predicted alkaline phosphatase activity at four pNPP concentrations.....	73
16.	A comparison of experimental and predicted alkaline phosphatase activity at seven copper concentrations.....	75

INTRODUCTION

The goal of this project has been to investigate the effect of variations in the concentration of aluminum and copper on the green alga Scenedesmus quadricauda. The hypothesis is that the growth rate of the alga and the activity of its enzymes are related to the concentration of the free cupric ion. The research has environmental significance because of the increase in the aluminum concentration of lakes affected by acid rain. Aluminum is thought to be responsible for some of the observed toxicity in these systems. I will present evidence that the effect of aluminum may be due to its interaction with the metal chemistry; the increase in aluminum leads to an increase in the toxic form of another metal such as copper.

The background will begin with a description of the environmental conditions which lead to this research. The concept of chemical speciation will be introduced. The background will cover the theory of speciation, its relationship to toxicity, and factors that control copper and aluminum speciation in the environment. This research is based on the understanding that the addition of one metal will affect the speciation of other metals in the system. The use of computer programs to estimate the results of metal interactions will also be discussed. The final purpose of the background is to describe the test

organism and enzyme system used to assay the effect of aluminum and copper.

The experiments performed to test the hypothesis are presented in chapters I through IV. The effect of variations in the concentration of aluminum and copper on the growth rate of Scenedesmus quadricauda was investigated (Chapter II). The relationship between the concentration of these metals and the activity of the enzyme alkaline phosphatase was then studied. Experiments were performed in_vitro with an isolated enzyme (Chapter III), and in_vivo with Scenedesmus cultures (Chapter IV). In Chapter V, a model is developed to predict the effect of changes in metal speciation.

CHAPTER I

BACKGROUND

Aluminum and Acid Rain

An environmental effect of acid precipitation is the increased mobility of aluminum (Driscoll et al 1984; LaZerte 1984). Aluminum, a common component of soil, is insoluble at the natural pH of rain. Acid rain leads to the dissolution of aluminum compounds. Although this process neutralizes the acidic input, it leads to an increase in dissolved aluminum (Johnson 1984). Increases in the concentration of this metal have been detected in streams, lakes and ground water (Driscoll et al 1984). Acidified lakes have been found to have aluminum concentrations 10 to 50 times higher than neutral lakes in the same region (Cronan and Schofield 1979).

The increase in aluminum is thought to be responsible for the loss of fish in lakes affected by acid rain (Dillon et al 1984). As acidity increases, toxic conditions may be produced by dissolved aluminum at pH levels that are not in themselves harmful (Cronan and Schofield 1979). A statistical survey suggested that the aluminum concentration is the primary chemical factor controlling the survival of fish (Schofield and Trojnar 1980). Although laboratory studies have shown that it is possible to kill fish (Baker and Schofield 1982) and phytoplankton

(Helliwell 1983) given high enough concentrations of aluminum, the mechanism of toxicity in the environment has not been demonstrated.

Metal Speciation

Metals can be present in a variety of chemical forms in natural waters. Each individual compound or complex is called a chemical species. The total concentration of a metal is the sum of the concentrations of its free and complexed species. The relative concentration of each species depends on the chemical interactions of all the reactive components of a system. The important factors controlling metal speciation is the relationship between the number of ligands available for metal complexation and the total concentration of metal.

There is chemical competition for complexation if two metals can reversibly bind to the same site. This competition can be represented by the equation:

An increase in the concentration of either metal will cause the free ion concentration of both metals to increase. The free metal activity is often reported as the negative log concentration or pMe. This convention is analogous to pH.

The interactions of metals in aquatic solutions can be modeled using computer programs (Westall et al 1976; Fontaine 1984). These programs are used to estimate the equilibrium concentration of the various chemical species. The programs used in this research are MINEQL (Westall et al 1976) and MICROQL (Westall 1979). These are basically the same program designed for mainframe and micro computers respectively. The input for these programs consists of the concentration of chemical components of a solution and the equilibrium or stability constants for all the possible interactions of the components. Components are defined in such a way that all possible species can be written as products of reactions involving only components. The computer performs a series of equilibrium calculations until the mass balance error is below an acceptable level. The output consists of the concentration of each species.

Although MICROQL is a valuable tool for determining chemical speciation in the laboratory, it does have limitations for environmental application. It can only be accurately used for solutions in which the exact concentration of each component is known. Also stability constants are necessary for all the possible chemical interactions. This information is not generally available for natural waters. The program assumes that equilibrium is reached between all of the species which may not be valid for a dynamic natural system (Fontaine 1984).

Copper and Aluminum Toxicity: Effect of Speciation

Copper is toxic to organisms of many taxonomic groups in aquatic ecosystems. Phytoplankton (Peterson et al 1984), zooplankton (Giesy et al 1983), and fish (Shaw and Brown 1973) have all been shown to be susceptible to copper toxicity. Natural levels of copper can inhibit algal growth. Huntsman and Sunda (1980) state there is 'increasing evidence for the toxicity of copper even in unpolluted water'. Steeman-Nielsen and Wiium-Andersen (1970) found copper to be poisonous at concentrations found in natural waters.

Chemical speciation is critical in determining the effect of metals in aquatic ecosystems. Only certain metal species interact with the biota. The activity of the free cupric ion (Cu^{+2}) is thought to be the primary factor determining the toxicity of copper to phytoplankton (reviewed in: Huntsman and Sunda 1980) and fish (Shaw and Brown 1973). Organic complexation decreases copper toxicity (Peterson et al 1984). Algal growth rate (Petersen 1982), enzyme activity (Rueter 1983), and nutrient uptake (Rueter and Morel 81) can all be related to the free cupric ion concentration.

The relative toxicity of different aluminum species is not well understood because chemical conditions have not been well controlled in aluminum toxicity studies. Some believe that the free metal ion is the toxic species

(Burrows 1977), while others claim $\text{Al}(\text{OH})_2^+$ is the most toxic form (Helliwell 1983). Organically complexed aluminum is essentially non-toxic (Johnson et al 1984).

Variations in the chemistry of a solution can lead to toxicity of a metal without an increase in the total concentration of that metal. Copper toxicity has been shown to increase by decreasing the concentration of natural complexing agents (Sunda and Lewis 1978). Petersen (1982) demonstrated that the addition of zinc can lead to copper toxicity in algal culture. Zinc competes and displaces some copper from the chelator present in the medium thus increasing the cupric ion concentration. Aluminum may exert its effect by similar mechanisms in natural systems.

Environmental Control of Metal Speciation: Humic Substances

The environmental application of the results of this research depends on the reversible complexation of metals in lakes. Due to the presence of a diverse array of organic and inorganic ligands in natural waters, a number of compounds could be important in the complexation capacity of an aquatic system (Perdue 1979). Both copper (Shaw and Brown 1973) and aluminum (LaZerte 1984) are naturally found as free metal ions, inorganic species, and as organic complexes with biological compounds.

Humic substances are a class of compounds which have been shown to be important in metal complexation (Gamble et al 1983). Terrestrial plant material is the primary source

of these compounds. The presence of various branched carbon chains prevents further degradation by micro-organisms. Humic substances are often the most prevalent organics in aquatic systems (Perdue 1979). These polymeric compounds are heterogeneous in terms of molecular weight, number of binding sites, and stability of metal complexes (Giesy et al 1983). Humics are effective chelators due to the presence of carboxyl, alcoholic, ketolic, and phenolic groups (Giesy 1983).

Although humic substances have been shown to form complexes with metals, the interaction can not be characterized with defined stability constants (Perdue and Lytle 1983; Giesy 1983; Saar and Weber 1982). This is partially due to the fact that humics are a class of different compounds, although attempts to separate them for individual study has proven to be unsuccessful (Perdue and Lytle 1983). Metal-humic interactions are best described by experimentally derived conditional stability constants and maximum binding capacities (Giesy et al 1983). Generally the derived values are only valid for humics from a particular environment. The stability constant is also influenced by pH due to proton-humic interactions (Perdue 1983).

Humic substances have been shown to be important in determining copper speciation and toxicity in natural waters (Giesy et al 1983, McKnight 1981, Huntsman and Sunda 1980). Copper can form stable complexes with a variety of

organic ligands. The complexation of copper with humics can be described by the equation:

Thus the activity of copper is determined by the concentration of humics, competing metals, and the pH. Giesy (1983), McKnight (1981), and Gatcher et al (1978) all found that a majority of the total dissolved copper in various lakes is associated with humics. Uncomplexed copper is generally the second most prevalent species. As discussed earlier, humic bound copper is unavailable to phytoplankton and does not lead to toxicity. Giesy (1983) and McKnight (1981) state that humic complexation is an important factor controlling copper toxicity to phytoplankton.

Metal ions compete with one another for binding sites; increasing the concentration of one metal increases the free ion concentration of others by displacement (Gamble et al 1983). Due to the chemical uncertainty of these compounds and differences in humic substances from different environments it is not possible to predict the exact outcome of competition between aluminum and copper in natural waters. A generalized order of metal-humic stabilities is $\text{Al} > \text{Cu}$ based on competition for binding sites (Giesy 1983). Giesy found that aluminum would displace copper from humics extracted from various sites in South Carolina. Kerndorff and Schnitzer (1980) allowed eleven metals to simultaneously interact with a humic solution. Although the results varied with pH, the

conditional binding of aluminum is similar to that of copper. Of the eleven metals tested both aluminum and copper were found to have intermediate affinity for humics. Although the displacement of copper by aluminum has not been shown in a natural system, it should occur given the similar affinities that the two metals have for humic complexation.

In this research, the synthetic chelators EDTA (Ethylenedinitrilotetraacetic acid) and NTA (Nitrilotriacetic acid) were used to model the actions of humic substances. Previous studies have shown this to be a useful approach. Davey et al (1973) found the results of algal bioassays in natural water could be calibrated by comparison to similar results in media containing EDTA. Gillespie and Vaccaro (1978) used a bacterial bioassay and found similarities in the effects of natural organics and EDTA on decreasing copper toxicity. Sunda and Lewis (1978) grew phytoplankton in media containing copper and various concentrations of humic substances and obtained the results predicted using synthetic chelator models. A complicating factor is that a high concentration of EDTA may be toxic to algae (Rueter, pers. comm.)

Scenedesmus quadricauda

The green alga Scenedesmus quadricauda (culture II, Freshwater Institute, Winnipeg, Canada) was used as a test organism in this research. It was chosen because of its

use in similar experiments (Petersen 1982). Its taxonomic position is (Bold and Wynne 1978):

Division- Chlorophycophyta

Order- Chlorellales

Family- Scenedesmaceae

Scenedesmus quadricauda is found as colonies containing up to four cells. The cell are joined laterally, and the terminal cells often have two spikes protruding from the ends. The number of cells per colony in laboratory cultures is influenced by the media used (Bold and Wynne 1978). Four and one cells per colony were the most common numbers observed in this research. Scenedesmus quadricauda are widely distributed in freshwater habitats.

Alkaline phosphatase

Metal toxicity can be expressed as an inhibition of any critical cellular process (Peterson et al 1984). I chose to measure the activity of alkaline phosphatase for a variety of reasons. Alkaline phosphatase has been shown to be a biochemical marker of metal toxicity (Rueter 1983). It is important in the environmental success of phytoplankton (Currie and Kalf 1984). The enzyme is associated with the surface of algae so it is in direct contact with the chemistry of the medium (Rueter 1983). Alkaline phosphatase is known to be a zinc metallo-enzyme (Coleman and Gettins 1983) which allows for modeling of

the interactions between the toxic metals and the zinc binding site.

Alkaline phosphatase is found in a wide variety of taxonomic groups. Isolated enzymes from various species have been studied and they share the same basic properties (Coleman and Gettins 1983). The enzyme is a dimer with a molecular weight of about 95,000. At least two zinc ions are associated with each dimer (Cohen and Wilson 1966). Alkaline phosphatase is a nonspecific phosphomonoesterase (Coleman and Gettins 83) which catalyzes the hydrolysis of orthophosphate from a wide variety of organic phosphate compounds.

Alkaline phosphatase activity can be critical in controlling the success of phytoplankton species (Currie and Kalft 1984a). Phosphorus is often the resource limiting phytoplankton growth in lakes (Goldman and Horne 1983). Alkaline phosphatase is required to utilize dissolved organic phosphate (DOP), which may be the primary source of phosphorus for phytoplankton (Currie and Kalft 1984a,b). Because of a faster uptake rate, bacteria utilize over 97% of the inorganic phosphate. If metal toxicity decreases the activity of alkaline phosphatase, it would lead to a decrease in phosphorus availability to phytoplankton.

Alkaline phosphatase has been shown to be a biochemical marker of copper toxicity (Rueter, 1983). This effect can be detected using a purified enzyme

solution, cell free algal extract, or active algal culture. The activity of the enzyme decreases in relation to an increase in the cupric ion concentration. The effect is thought to be due to the displacement of the native zinc ion by copper. Both metals form divalent cations with similar ionic radii and bonding geometries (Vallee 1958).

Background Summary

Acid rain leads to an increase in the concentration of aluminum in some lakes. My goal is to present evidence that the observed effect of increased aluminum may be due to the interaction of this metal with the aqueous chemistry of the system. The toxicity of a metal can be related to the concentration of a particular chemical species of the metal. In aquatic systems, the free metal ion is often the most toxic form. This form is often found in low concentration due to complexation with humic substances. Aluminum may exert its effect by competing for binding sites with more toxic metals such as copper. This would lead to an increase in the toxic form of copper without an increase in the total copper concentration. This hypothesis is tested by investigating the effect various concentrations of aluminum and copper have on algal growth rate and enzyme activity. Changes in these biological parameters should relate to variations in metal speciation.

Chapter II

The Growth Rate of Scenedesmus quadricauda as a function of aluminum and copper concentration

INTRODUCTION

The goal of this experiment is to determine the sensitivity of Scenedesmus quadricauda to aluminum and copper toxicity. The alga is grown in media containing various concentrations of aluminum and copper. The growth rate will be compared to the total concentration of each metal as well as the free ion activity of each metal. The hypothesis is that the growth rate should be related to the activity of the free cupric ion.

The design of this experiment is based on work done on the same organism by Petersen (1982). In his experiments, Petersen investigated the effect variations in zinc and copper concentration had on algal growth rate. Growth data was analyzed in terms of total concentration as well as the estimated chemical speciation. He found that in the presence of both metals, the observed toxicity could be related to the free cupric ion.

METHODS

Care must be taken with experiments on the effects of trace metals to prevent metal contamination. Nanopure water obtained from a Barnstead system was used in all

experiments. Plasticware was allowed to soak in 4% HCL for 24 hours and rinsed five times with Nanopure water immediately prior to use. The growth medium was a run through a column of Chelex 100 (Bio-rad lab.). This cation exchange material removes any metal contaminants present in the solution. All culture manipulations were performed in a laminar hood (Environmental Air Control Inc.). Clean plastic gloves were worn while handling the culture flasks.

Scenedesmus quadricauda (culture II, Freshwater Institute algal culture collection, Freshwater Institute, Winnipeg, Canada) was used as the test organism. The algae were grown in Fraquil (Morel et al, 1979), as modified by Petersen (1982). The chemical composition of the medium is listed in Table I. 1000x concentration stocks of the nutrients; NaSiO_4 , K_2HPO_3 , NaNO_3 and the major ions; CaCl_2 , MgSO_3 , and NaHCO_3 were prepared. A metal stock solution which contained 1×10^{-3} EDTA and 1000x concentration of the metals (excluding Cu and Al) listed in table I was also prepared. Two milliliters of each nutrient stock were diluted to one liter with Nanopure water. The major ions were similarly mixed. The nutrient mix and the ion solution were run through separate Chelex columns. 500 ml of each chelexed solution were combined and 1 ml of the metal stock was added to produce one liter of Fraquil. The medium was filter sterilized using a 0.45 μM filter (Nucleopore Co.).

TABLE I
MODIFIED FRAQUIL CULTURE MEDIA

chemical concentration component	concentration mol / L	chemical component	mol / L
Ca ⁺²	2.5 x10 ⁻⁴	CO ₃ ⁻²	1.5 x10 ⁻⁴
Mg ⁺²	1.5 x10 ⁻⁴	SO ₄ ⁻	1.5 x10 ⁻⁴
K ⁺	2.0 x10 ⁻⁵	Cl ⁻	5.2 x10 ⁻⁴
Fe ⁺³	4.5 x10 ⁻⁷	B(OH) ₄ ⁻	1.0 x10 ⁻⁷
Mn ⁺²	2.3 x10 ⁻⁸	MoO ₄ ⁻²	1.5 x10 ⁻⁹
Co ⁺²	2.5 x10 ⁻⁹	NO ₃ ⁻	1.0 x10 ⁻⁴
Zn ⁺²	1.0 x10 ⁻⁹	EDTA ⁻⁴	5.0 x10 ⁻⁶
NH ₃	1.3 x10 ⁻⁹	Al ⁺³	variable
H ⁺	variable	Cu ⁺²	variable

Concentration of the components of the media Fraquil (Morel et al 1979) as modified by Petersen (1982). H⁺ concentration (pH 5-8) was variable due to changes in the aluminum and copper concentrations.

Stock solutions of 1×10^{-1} M AlCl_3 and 5×10^{-2} M CuSO_4 were used as the source of these metals. 100 ml of medium at each experimental concentration was made in 250 ml Teflon flasks. These solutions were allowed to sit for twenty-four hours before experiments were performed so equilibrium between the metals and EDTA could be obtained.

The growth experiments were performed in 30 ml polycarbonate tubes. 20 ml aliquots of medium were inoculated with 0.2 ml of a stationary phase maintenance culture of Scenedesmus. Daily measurements were taken by placing each tube in a fluorometer (Turner Designs, Model 10) set to detect the fluorescence of chlorophyll a:

light source- bulb F4T5

excitation filter- c/s 5-60

emission filter- c/s 2-64

The increase in chl a fluorescence was used as the measurement of growth. The cultures were kept in an incubator (Enviroator Co.) at 18°C . The light was continuous and had an intensity of $100 \mu\text{ Einsteins m}^{-2} \text{ s}^{-1}$. The tubes were kept in clear plastic racks and shaken at 100 rpm. Each growth experiment lasted 7 days. The exponential phase of growth was determined by graphing the daily fluorescence on semi-log paper. Growth rate was determined using the equation:

$$u = (\ln F_2 - \ln F_1) / (\# \text{ days})$$

u = growth rate

F_1 = fluorescence on first day of exponential growth

F_2 = fluorescence on last day of exponential growth
= number of days between F_1 and F_2

The effect of aluminum and copper on algal growth rate was first determined individually. After the effective level of each metal was known, a matrix of metal concentrations were derived (Table II). The copper concentrations used in these experiments were 0, 3, 4, 4.5, and 5×10^{-6} M, while the aluminum concentrations were 0, 2, 3, 4, 5, 6, and 8×10^{-6} M. The growth rate was determined for thirty three concentration combinations of these metal concentrations (table II).

The concentrations of the chemical species at each set of aluminum and copper concentrations were estimated using the computer program MINIQL (Westall et al 1976). Critical equilibrium constants involving EDTA, aluminum, and copper were compared to other published values (Smith and Martell 1975).

RESULTS

Although individual cultures had growth rates as high as 0.93 day^{-1} , the average U_{max} for any concentration combination was 0.85. The control cultures with no additional metals always had growth rates about 0.2 units lower than cultures with low concentrations of aluminum or copper. Additional metal did not have a noticeable effect on growth rate until a threshold concentration is reached

TABLE II

GROWTH RATE DAY⁻¹ (upper) AND pCu (lower) AS
ESTIMATED BY MINIQL (Westall et al 1976),
FOR EACH CONCENTRATION OF
ALUMINUM AND COPPER

		Copper ($\times 10^{-6}$ M)				
		0	3	4	4.5	5
Aluminum ($\times 10^{-5}$ M)	0	0.65 13.0	0.85 11.42	0.77 10.83	0.81 10.11	0.08 8.20
	2	0.76 13.0	0.80 11.41	0.77 10.82	0.76 9.64	0.21 7.86
	3	N.D. 13.0	N.D. 11.38	0.77 10.79	0.58 9.61	0.04 7.46
	4	0.67 13.0	0.70 9.96	0.48 8.77	0.42 7.75	0.16 6.35
	5	0.68 13.0	0.46 6.97	0.23 6.50	0.33 6.28	0.0 6.08
	6	0.44 13.0	0.40 6.39	0.10 6.11	N.D. 5.98	N.D. 5.87
	8	0.08 13.0	0.16 6.03	0.15 5.84	0.0 5.76	0.0 5.68

N.D.- no data

Figure 1. The growth rate day⁻¹ of *Scenedesmus* as a function of the total concentration of aluminum and copper.

(Table II ;Figure 1). Above this level the growth rate decreases dramatically. The threshold concentration of both aluminum and copper depends on the concentration of the other metal.

The EC_{50} (effective concentration-50%) is the concentration of a metal at which the growth rate of the alga is fifty percent of maximum. The $Al-EC_{50}$ in the absence of copper is about 6×10^{-5} M. As the amount of copper increases, the $Al-EC_{50}$ decreases (Figure 1). At higher copper concentrations, less aluminum is required to inhibit growth. The $Al-EC_{50}$ can not be determined if the copper concentration is 5×10^{-6} M since the growth rate in the absence of aluminum is less than 50% of the control rate. The growth rate is slightly higher at this copper concentration if some aluminum is present.

At aluminum concentrations below 2×10^{-5} M, the $Cu-EC_{50}$ is between 4.5 and 5×10^{-6} M. If the aluminum concentration is raised to 3×10^{-6} M, 4.5×10^{-6} M copper leads to a 25% decrease in growth rate. The $Cu-EC_{50}$ is about 4×10^{-6} M if the aluminum concentration is 4×10^{-5} M. It drops to 3×10^{-6} M copper at 5×10^{-5} M aluminum. As the growth rate at 6×10^{-5} M aluminum is only about 50% of the control, it is not possible to determine an $Cu-EC_{50}$. At this aluminum concentration, the growth rate remains fairly constant until the copper concentration exceeds 3×10^{-6} M.

Figure 2. The growth rate day⁻¹ of *Scenedesmus* as a function of cupric ion activity (pCu). See the discussion for a description of A and B.

At the highest aluminum concentration tested, the growth rate was slightly higher at copper concentrations of 3 and 4×10^{-6} M than in the absence of copper.

A statistical correlation ($r^2=0.87$) was determined between the log concentration of the cupric ion (pCu) and the growth rate of Scenedesmus (Figure 2). Linear regression analysis revealed no such correlation between the growth rate and pAl ($r^2=0.46$), pH ($r^2=0.56$), or any of the other species. As the growth rate changed due to variations in the concentration of either metal, a relationship between the total concentration of a metal and the growth rate was not obtained.

DISCUSSION

Cupric ion activity is the chemical factor controlling the growth rate of Scenedesmus. Aluminum exerts its effect by competing with copper for organic binding sites. As the aluminum concentration increases, copper ions are displaced from EDTA and the cupric ion activity increases. The effect of competition between aluminum and copper for complexation is demonstrated in figure 3. At a set copper concentration, the activity of the cupric ion increases (pCu decreases) with aluminum concentration.

The results obtained agree with published results from a similar experiment with copper and zinc (Petersen 1982). He found that in the presence of copper, an increase in

Figure_3. The activity of the cupric ion increases with aluminum concentration. The copper concentration is set at 4×10^{-6} M copper.

zinc could lead to increased copper toxicity. The effect was thought to be due to the displacement of copper ions from EDTA. An important point is that although two metals may individually inhibit the growth of Scenedesmus, speciation must be known to determine the toxic agent in the presence of both metals.

Although a good correlation is obtained between growth rate and cupric ion activity, figure 2 shows two sets of data that appear to vary from remainder of the samples. The three data points labeled A are all cultures that contained the highest copper concentration (5×10^{-6} M). The total copper concentration may be important in determining toxicity at this level. Also, at 5×10^{-6} M the concentration of copper and EDTA are equal. EDTA's effectiveness as a buffer depends on the relative concentration of the metal and the chelator. If the buffering capacity is exceeded, the cupric ion activity could fluctuate and cause higher values than estimated by MINEQL.

The cultures labeled B all contained 2×10^{-5} M aluminum. The growth rate of these cultures was higher than predicted. A low concentration of aluminum may provide protection from copper toxicity. Evidence of a protective effect of aluminum was also found in experiments on the effect of aluminum and copper on the activity of alkaline phosphatase (Chapter III).

Figure_4. The growth rate day^{-1} of *Scenedesmus* as a function of total aluminum. The copper concentration is set at $4 \times 10^{-6} \text{ M}$ copper. Although an increase in aluminum does lead to a decrease in growth rate, this research has shown the cupric ion to responsible for the toxic effect.

Although there is a statistical relationship between the increase in aluminum and a loss of fish (Cronan & Schofield 1979), the information is not available to determine what changes in metal speciation may have occurred. Evidence of cause and effect can be misleading if the data is incomplete. Field data often consist only of changes in pH and total aluminum concentration. If a lake subjected to acid rain is modeled by a set of cultures with the same copper concentrations but different amounts of aluminum, toxicity does appear to increase with aluminum concentration (Figure 4). But the results of this experiment show that the toxicity is not due directly to aluminum but to changes in the aquatic chemistry.

CONCLUSION

The green alga Scenedesmus quadricauda was grown in thirty three different concentrations of aluminum and copper. The growth rate of the alga was compared to the total concentration of each metal, as well as, estimates of metal speciation. Growth rate was found to be related to the cupric ion activity. Aluminum alone has a minimal toxic effect. The results support the hypothesis that the observed effect of increased aluminum may be due to interactions with the aquatic chemistry of a system.

CHAPTER III

ALKALINE PHOSPHATASE ACTIVITY AS A FUNCTION OF THE AQUATIC CHEMISTRY OF ALUMINUM, COPPER, AND ZINC

INTRODUCTION

This research attempts to characterize the interactions between aluminum and copper in relation to the toxicity of these metals. The hypothesis predicts that metal speciation is more important than the total metal concentration. In this chapter, the inhibition of alkaline phosphatase is used as an indicator of toxicity. This enzyme was chosen because it is important in the phosphate nutrition of algae (Currie and Kalf 1984) and it has been shown to be a biochemical marker of metal toxicity (Rueter 1982).

The use of enzyme activity assays has benefits over cell growth rate experiments in metal toxicity studies. The chemistry of the media can be simplified without concern for the nutritional requirements of cells. Algae may release organic chelators which complicate metal speciation calculations (McKnight and Morel 1979). Another benefit is that enzyme studies take less time so more experiments can be performed.

Three types of experiments were performed in this investigation. Ten enzyme buffers solutions were tested to

determine their suitability in trace metal studies. The activity of alkaline phosphatase as a function of aluminum and copper concentration was determined. The results of those experiments suggested that zinc present in the buffer affects enzyme activity. The final set of experiments investigates if alkaline phosphatase inhibition due to zinc removal is reversible.

SECTION 1

Enzyme Buffer Solutions

METHODS

Assay of alkaline phosphatase activity- Alkaline phosphatase, isolated from E. coli (Sigma Co), was diluted to one unit / ml in 1×10^{-3} M Tris (pH 8.3) and stored at 5 °C. 0.1 ml of the enzyme-Tris solution was added per 10 ml of the experimental buffer solution. The final concentration of the enzyme was 1×10^{-2} units / ml. 2.5 ml of the enzyme-buffer was added to a 1 cm path length cuvette. Enzyme assays were started by adding 0.1 ml of 1×10^{-3} M n-Nitrophenylphosphate (pNPP)(Sigma Co). The hydrolysis of pNPP was followed at 410 nm in a spectrophotometer (Baush & Lomb, Spectronic 100) over four minutes. An absorbance change of 0.001 corresponded to a 1.4×10^{-7} M change in pNPP concentration.

Enzyme_buffer_solutions- The requirements for a suitable enzyme buffer include: the enzyme must have measurable activity in the solution; the enzyme must remain active over time; the pH must be maintained within the range of 7.5 to 8.0; it must permit free metal activities to be fixed at effective total metal concentrations; and the equilibrium constants between all the components and aluminum and copper must be known so estimates of metal speciation can be readily calculated.

Ten buffers were tested to determine their suitability in metal toxicity studies (Table III). Experimental buffers were prepared by diluting concentrated stock solutions. Buffers were allowed to sit at least four hours before assays were performed. Assays were performed over a twenty-four hour period.

RESULTS

The activity of the enzyme was higher in 1×10^{-3} M Tris (1.3×10^{-6} M pNPP / min) than in any of the subsequent buffers (Table III). The activity showed little change over twenty-four hours. A set of experiments were performed to compare the use of EDTA and Tris. Four solutions were tested: 1×10^{-3} M Tris, 1×10^{-3} M EDTA, 1×10^{-3} M Tris + 1×10^{-6} M EDTA, and 1×10^{-3} M Tris + 1×10^{-5} M EDTA. The activity of alkaline phosphatase appeared to be inhibited by EDTA (Figure 5).

TABLE III
 ENZYME BUFFERS TESTED

Chemical composition	Activity ($\times 10^{-7}$ M pNPP /min)		Comments
	time 0	24 hrs	
1×10^{-3} M tris	25.2	21.0	equilibrium constant not available for Al-tris
1×10^{-3} M tris 1×10^{-6} M EDTA	19.6	13.7	apparent inhibition due to EDTA
1×10^{-3} M tris 1×10^{-5} M EDTA	18.2	5.6	apparent inhibition due to EDTA
1×10^{-3} M EDTA	11.3	3.9	apparent inhibition due to EDTA
1×10^{-4} M NaHCO_3 1×10^{-5} M EDTA 1×10^{-8} M ZnCl	3.6	-. -	ionic strength too low
1×10^{-3} M NaHCO_3 1×10^{-5} M EDTA 1×10^{-8} M ZnCl	14.0	7.0	ZnCl does not prevent EDTA inactivation
1×10^{-3} M NTA	0.0	-. -	NTA solution is not a suitable buffer
1×10^{-3} M NaHCO_3 1×10^{-5} M NTA 1×10^{-8} M ZnCl	18.2	16.8	effective buffer
1×10^{-3} M NaHCO_3 1×10^{-5} M NTA 1×10^{-8} M ZnCl 1×10^{-3} M CaCl_2	17.6	5.6	activity not maintained
1×10^{-3} M NaHCO_3 1×10^{-5} M NTA	18.1	16.8	effective buffer

Figure 5. The activity of alkaline phosphatase ($\times 10^{-7}$ M pNPP / min) in five buffers over twenty-four hours.

◻ - 1×10^{-3} Tris, 1×10^{-6} EDTA

◐ - 1×10^{-3} Tris

◻ - 1×10^{-3} Tris, 1×10^{-5} EDTA

■ - 1×10^{-3} EDTA

⊕ - 1×10^{-3} NaHCO₃, 1×10^{-3} CaCl₂, 1×10^{-5} NTA, 1×10^{-8} ZnCl

Buffer AP-B contained 1×10^{-4} M NaHCO_3 , 1×10^{-5} M EDTA, and 1×10^{-8} M ZnCl_2 . The measured activity was 1.1×10^{-7} M pNNP / min. This is only 10 percent of the activity found in 1×10^{-3} M Tris. The activity increased about seven fold to 7×10^{-7} M pNNP / min if the concentration of NaHCO_3 was raised to 1×10^{-3} M. After three hours the activity in this solution decreased by 50 percent.

The decrease in activity was not noted when the EDTA was replaced by NTA. The activity of the enzyme in Buffer AP-D (1×10^{-3} M NaHCO_3 , 1×10^{-5} M NTA, and 1×10^{-8} M ZnCl_2) was about 18.2×10^{-7} M pNNP / min and remained constant for four days. Although Buffer AP-D appeared to be satisfactory, two variations were tested. In an attempt to lower the amount of aluminum and copper required to cause enzyme inhibition, Ca^{+2} was tested as a competing ion. Although the activity of alkaline phosphatase in Buffer AP-E (Buffer AP-D + 1×10^{-3} M CaCl_2) at time zero and two hours was similar to the activity in Buffer AP-D, it dropped by fifty percent at six hours. The activity continued to decrease with time (Figure 5). Buffer AP-F was a ZnCl_2 free version of AP-D. The activity of the enzyme in AP-F was similar to the results in AP-D. Buffer AP-D (1×10^{-3} M NaHCO_3 , 1×10^{-5} M NTA, and 1×10^{-8} M ZnCl_2) was chosen as the buffer to use in the aluminum and copper experiments.

SECTION 2

Aluminum and Copper

METHODS

Alkaline phosphatase was exposed to ten aluminum-copper concentration combinations. Assays were performed over twenty-four hours. The activity of the enzyme was compared to both the total concentration and calculated metal ion activities.

The experimental concentrations of aluminum and copper were 0, 2, 4, and 6×10^{-5} M. Ten different combinations of the two metals were tested (Table IV). In all cases, the total metal concentration did not exceed 6×10^{-5} M. Stock solutions of AlCl_3 (1×10^{-2} M) and CuSO_4 (5×10^{-2} M) were added to 100 milliliters of the buffer to make each concentration combination. The pH of each solution was measured (Orion Research, Ionalyzer 404) and adjusted to 7.7 with 0.1 M NaOH. The solutions were allowed to sit for 12 hour before the enzyme assay. 10 ml of each metal combination was placed in 30 ml polycarbonate tubes. At time zero, 0.1 ml of the enzyme-Tris solution was added to each tube. Assays were performed at time zero, 2 hours, 8 hours, and twenty four hours.

Copper is thought to inhibit alkaline phosphatase by displacing the native zinc ion. The result of competition between the two metals should depend on the relative activities of copper and zinc. The zinc concentration in

TABLE IV

ACTIVITY OF ALKALINE PHOSPHATASE AFTER
 TWENTY-FOUR HOURS EXPOSURE TO
 ALUMINUM AND COPPER

copper $\times 10^{-5}$ M	aluminum $\times 10^{-5}$ M	experimentally determined activity $\times 10^{-7}$ M pNPP / min			
0	0	17.2	17.6	15.8	15.8
2	0	14.0	12.6	13.2	10.2
4	0	5.6	6.0	10.6	8.4
6	0	3.6	3.6	4.2	3.2
0	2	17.6	16.8	16.2	16.8
2	2	7.0	6.0	6.4	6.4
4	2	6.4	4.6	4.6	4.6
0	4	16.2	15.8	14.0	14.8
2	4	6.4	6.4	8.4	8.4
0	6	14.8	15.8	13.5	10.2

the aluminum-copper buffers discussed above was raised from 1×10^{-8} M. to 1×10^{-6} M. After twenty-four hours exposure to increased zinc, the assays were repeated. The results obtained at the two zinc concentrations were compared.

The chemical speciation for each sample was estimated using a Pascal version of MICROQL (Westall 1985). The program was run on a Zenith 151 PC (Zenith data systems). A matrix (Table V) was set up for the component of the buffer. The stability constants used were obtained from the MINEQL program data file (Westall et al 1976).

RESULTS

Four data sets were available for each metal concentration combination (Table IV). The difference in activity between data sets was usually less than 2×10^{-7} M pNPP / min. An exception was the sample with 4×10^{-5} M copper; the data between the experimental runs differed by 5×10^{-7} M pNPP / min (range: $5.6 - 10.6 \times 10^{-7}$ M pNPP / min). The study was repeated for all the samples containing only copper. The activity determined at 4×10^{-5} M copper (4.9 & 7.3×10^{-7} M pNPP / min) suggests the lower activities were more accurate.

Individually aluminum and copper produced different effects on the activity of alkaline phosphatase (Table VI). The presence of copper led to an immediate inhibition of the enzyme. Linear regression analysis indicates that

TABLE V

CHEMICAL COMPONENTS, SPECIES, AND
STABILITY CONSTANTS USED
IN MICROQL

Components- NTA, Zn^{+2} , Al^{+3} , Cu^{+2} , H^+

<u>Species</u> -----	<u>K</u>	<u>Species</u> -----	<u>K</u>
NTA	0.0	NTA-H	10.41
NTA-H ₂	13.55	NTA-H ₃	15.72
Zn	0.0	ZnOH	-9.53
Zn-NTA	12.22	Zn-NTA ₂	14.11
Cu	0.0	CuOH	-7.84
Cu-NTA	14.32	CuOH-NTA	4.78
Cu-NTA ₂	17.01	Al	0.0
Al-NTA	14.53	AlOH-NTA	9.16
AlOH	-4.56	Al-NTA ₂	20.53
OH	-14.00		

TABLE VI

ACTIVITY OF ALKALINE PHOSPHATASE
AT TEN ALUMINUM-COPPER
CONCENTRATIONS

copper activity $\times 10^{-5}$ M	aluminum $\times 10^{-5}$ M	pCu^{+2}	pAl^{+3}	activity	
				time 0	24 hr
0	0	-. -	-. -	15.5	16.6
2	0	5.24	-. -	11.3	12.6
4	0	4.76	-. -	11.2	5.8
6	0	4.54	-. -	9.5	3.7
0	2	-. -	8.14	16.4	16.8
2	2	5.11	7.92	14.4	6.5
4	2	4.74	7.88	11.6	4.7
0	4	-. -	7.66	17.0	15.2
2	4	5.06	7.60	14.8	7.4
0	6	-. -	7.44	16.2	13.0

activity = $\times 10^{-7}$ M pNPP / min

the activity rate decreases 0.92×10^{-7} M pNPP / min per 1×10^{-5} M copper ($r^2=0.88$). The inhibition increased with time so that by twenty-four hours the enzyme activity compared to control is lower by 2.2×10^{-7} M pNPP / min per 1×10^{-5} M copper ($r^2=0.97$).

The experimental concentrations of aluminum did not immediately inhibit the activity of alkaline phosphatase. At time zero, enzyme activity appeared to increase slightly with aluminum concentration, but with a r^2 value of 0.55 this was not significant. After twenty-four hours, the activity of the enzyme decreases 0.4×10^{-7} M pNPP / min per 1×10^{-5} aluminum ($r^2=0.73$). This is only about one fifth the decrease caused by the same amount of copper.

The effect of increasing aluminum at a set copper concentration changed with time. For example, if the copper was 2×10^{-5} M, an increase of aluminum led to an activity increase of 0.8×10^{-7} M pNPP / min per 1×10^{-5} M at time zero ($r^2=0.85$). The aluminum may protect the enzyme from immediate copper inhibition. After twenty-four hours, an increase in aluminum led to an inhibition similar to an increase in copper. The activity dropped 1.14×10^{-7} M pNPP / min per 1×10^{-5} M aluminum (Figure 6). The inhibition is 400 percent greater than caused by aluminum alone.

An analysis of all the data indicates that alkaline phosphatase activity is related ($r^2=0.78$) to cupric ion

Figure 6. The activity of alkaline phosphatase ($\times 10^{-7}$ M pNPP / min) as a function of aluminum concentration in the presences and absences of copper.

■ - no copper

⊙ - 2×10^{-5} M copper

Figure 7. The activity of alkaline phosphatase ($\times 10^{-7}$ M pNPP / min) as a function of cupric ion activity (pCu).

Figure_8. A comparison of alkaline phosphatase activity ($\times 10^{-7}$ M pNPP / min) at two zinc concentrations.

concentration . If the questionable data discussed in the first paragraph of this section is ignored (activity of 10.6×10^{-7} M pNPP / min at 4×10^{-5} M copper), the r^2 value increases to 0.85 (Figure 6). The activity of alkaline phosphatase drops 10.4×10^{-7} M pNPP / min per one unit change in pCu (pCu = 5 ± 0.5).

The enzyme activity at various aluminum and copper concentrations was compared between two experiments with the same zinc concentration (1×10^{-8} M) and two experiments with a 100x difference in added zinc concentrations (Figure 8). At time zero, the relative activity of the enzyme exposed to copper was greater in the solution with higher zinc. The difference averaged about ten percent of the control rate. After twenty-four hours, the rate at all copper and aluminum concentrations was higher in the buffer with more zinc, but the relative inhibition caused by these metals was the same.

SECTION 3

Zinc Chemistry

The results obtained in section one and two suggested that the buffer solution used contained some zinc as a contaminant. Nitrilotriacetic acid (NTA), a component of the enzyme buffer, competes as a ligand with alkaline phosphatase for zinc (Cohen & Wilson 1966). If the zinc ion concentration of a solution is low enough, NTA can

TABLE VII
REACTIVATION OF ALKALINE PHOSPHATASE
AFTER ZINC ADDITION

Time	Zinc	Enzyme Activity	+10 min <--time after zinc addition-->	+1 hour	+4 hours
0	1×10^{-8}	17.9	-. -	-. -	-. -
0	1×10^{-7}	17.9	-. -	-. -	-. -
30 min	1×10^{-8}	3.9	11.6	12.3	13.3
30 min	1×10^{-7}	8.0	12.3	14.0	15.0
6 hrs	1×10^{-8}	0.0	10.9	13.7	15.4
6 hrs	1×10^{-7}	4.2	9.5	11.2	14.0
20 hrs	1×10^{-8}	0.0	10.9	14.0	13.7
20 hrs	1×10^{-7}	4.9	8.1	10.9	11.6

activity = $\times 10^{-7}$ M pNPP / min

Figure 9. Alkaline phosphatase was exposed to a low zinc (1×10^{-8} M) buffer. The activity decreased with time. At 30 minutes, 6 hours and 24 hours the concentration of zinc was increased to 1×10^{-8} M and the recovery of activity was monitored.

● — ● - activity in low zinc buffer
 ▲ — ▲ - activity after increase in zinc concentration

to 1×10^{-6} M. The recovery of activity was determined at ten minutes, one hour, and four hours after zinc addition (Figure 9).

RESULTS

The use of chelex treated Na_2CO_3 did alter the characteristics of the buffer. It was necessary to add zinc to the solution to maintain alkaline phosphatase activity. With a zinc concentration of 2×10^{-8} M, the enzyme activity dropped from 17.9×10^{-7} M pNPP / min at time zero to 3.9×10^{-7} M pNPP / min after thirty minutes. No activity was detected after six hours. In "non-chelexed" buffer, the activity was consistent for four days.

Reactivation of alkaline phosphatase could occur twenty hours after zinc was stripped from the enzyme (Table VII). The activity of enzyme exposed to 1×10^{-8} and 1×10^{-7} M zinc would increase if the zinc concentration was raised to 1×10^{-6} M (Figure 9). About seventy-five percent of the reactivation would occur within ten minutes of zinc addition. The activity continued to increase over four hours.

DISCUSSION

Tris is the recommended buffer for alkaline phosphatase assays because of its weak zinc binding strength. Although the activity was the greatest in Tris, a dependable equilibrium constant between the chelator and

aluminum is not available. EDTA was not used because the activity of alkaline phosphatase could not be maintained in the presence of this chelator. EDTA may act to inhibit the enzyme by competing for the zinc ion.

The combination of NaHCO_3 (1×10^{-3} M) and NTA (1×10^{-5} M) was the buffer chosen for the metal toxicity studies. The NaHCO_3 maintained pH (≈ 7.7) and set the ionic strength. NTA functioned as the organic complexing agent. The equilibrium between NTA and the metals zinc, aluminum and copper were available (Smith and Martell 1975), making it possible to estimate chemical interactions and metal speciation.

This research is based on the observation that an increase in aluminum concentration leads to loss of aquatic organisms. My hypothesis is that the effect of the aluminum is due to its interaction with the metal chemistry of the system. The results of this experiment indicate that the toxicity of aluminum is dependent on copper. In the absence of copper, the rate of alkaline phosphatase activity dropped 2.3 percent per 1×10^{-5} M rise in aluminum. The rate at 6×10^{-5} M aluminum was eighty-five percent of control after twenty-four hours. With copper present, the effect of increasing aluminum was similar to increasing copper. A copper concentration of 2×10^{-5} M magnified the effect of increasing aluminum by 400 percent (Figure 6).

Alkaline phosphatase activity was found to relate to both the total copper concentration and cupric ion activity. This may be due to the high copper concentration required to inactivate the enzyme. If the concentration of metal is greater than the concentration of chelator, the free metal concentration is proportional to the total metal.

Rueter (1983) found alkaline phosphatase activity to relate only to the cupric ion. He also found the effective concentration to be much less ($pCu = 10 \pm 2$ vs. 5 ± 0.5). Rueter performed his experiments in 1×10^{-3} M Tris. At that chelator concentration, the effects of total copper and the cupric ion could be separated since a greater change in total copper is necessary to change the activity of the cupric ion. Zinc contamination may be involved in the results obtained in each study (Rueter, pers. comm). The difference in results indicates the importance of the solution chemistry in trace metal studies.

An unexpected result was the apparent protective effect of aluminum at time zero. The activity level of the enzyme exposed to inhibitory levels of copper increased with aluminum concentration. This effect was no longer observed after two hours. A possible explanation is a kinetic effect. Both aluminum and copper can bind to a specific site on the enzyme. The binding of copper leads to inhibition while the binding of aluminum has little effect. At time zero, aluminum binding prevents copper

inhibition. At higher aluminum concentrations more of the enzyme is protected. Over time copper can displace the aluminum from the enzyme so the effect is not observed at two hours.

Copper may inhibit alkaline phosphatase by displacing the native zinc (Foy et al 1978). The results of this experiment do not support or refute this hypothesis. The inhibition appears to occur in two steps. There is an initial inhibition which is evident in the time zero assays (Table VI). There is also an additional slow decrease in activity over time. Alkaline phosphatase contains 2 to 6 zinc ions per molecule (Coleman and Gettins, 1983). The affinities of the binding sites for the zinc vary. The initial inhibition may be due to copper displacing a weakly bound zinc or binding to another site on the enzyme. The slow continued inactivation may be caused by the interaction of the copper with a more tightly bound zinc ion. If the inhibition is caused by displacement, increasing the concentration of zinc should protect the enzyme by increasing the probability of zinc binding. The observation that increasing the zinc concentration leads to greater enzyme activity, but does not protect the enzyme from copper inactivation makes it hard to predict whether the inactivation is due to the displacement of the native metal. These results suggest that there is more zinc in the buffer solution than expected. This is supported by the observation that the characteristics of the enzyme in the

buffer changed after the solution was passed through a cation exchange column.

Cohen and Wilson (1966) state that NTA will strip zinc from isolated alkaline phosphatase if the zinc concentration is low. In the chelex treated buffer, a zinc concentration of greater than 1×10^{-6} M is required to prevent NTA inactivation. The inhibition caused by zinc removal can be reversed for at least twenty hours if the zinc concentration is increased. The reversibility of zinc binding to alkaline phosphatase allows zinc to be modeled as a co-factor of the enzyme (Chap V).

CONCLUSION

The effect of aluminum on alkaline phosphatase activity is dependent on copper. In the presence of copper, an increase in aluminum concentration leads to enzyme inactivation. Aluminum alone has little effect. This supports the hypothesis that aquatic metal chemistry is important in determining the environmental effect of increasing aluminum.

The activity of the enzyme was related to both the total copper and cupric ion concentrations. The result varied from the prediction that free cupric ion activity was the most important chemical factor. This may be due to the experimental metal concentrations being greater than the chelator concentration.

CHAPTER IV

ALKALINE PHOSPHATASE ACTIVITY IN SCENEDESMUS CULTURE: INTERACTIONS OF ALUMINUM AND COPPER

INTRODUCTION

This research has shown that the aquatic chemistry of aluminum and copper influences the growth rate of Scenedesmus quadricauda (Chapter II) and the activity of isolated bacterial alkaline phosphatase (Chapter III). In this chapter the effect of the metals on the activity of alkaline phosphatase from Scenedesmus is investigated.

Algae synthesize alkaline phosphatase in response to phosphate limitation. The enzyme, a nonspecific phosphomonoesterase (Coleman and Gettins 1983), allows the cells to utilize dissolved organic phosphate (DOP) as a phosphorus source. Bacteria may be able to outcompete algae for the uptake of inorganic phosphate, making DOP the primary source of phosphorus for phytoplankton (Currie and Kalft 1984).

Alkaline phosphatase is found associated with the cell membrane (Kuenzler and Perras 1965). External enzymes have advantages in trace metal studies (Rueter 1983). Alkaline phosphatase is exposed directly to the chemistry of the medium. Internal enzymes may be protected from metal toxicity by the large number of potential binding sites in the cytoplasm.

METHODS

A modified version of Fraquil (Morel et al 1979) was prepared. The use of 1×10^{-6} NTA in place of EDTA allows comparison between the chemical interactions in the growth medium and the enzyme buffer solution (Chapter III). 3-phosphoglycerate(3PG) (1×10^{-7} M) was used as the sole phosphorus source to promote the synthesis of alkaline phosphatase. The medium was prepared in the manner described in the methods of the growth rate experiments (Chapter II), except the 3PG was added to the solution after it had been "chelexed".

The experimental aluminum concentrations were 0, 0.5, 1 and 2×10^{-5} M. At each aluminum concentration, four copper concentrations were tested (0, 4, 8, 12×10^{-6} M). 700 mL cultures of Scenedesmus quadricauda (culture II, Freshwater Institute algal culture collection, Freshwater Institute, Winnipeg, Canada) were grown in acid washed one liter polycarbonate flasks. After the culture had grow for 5 to 8 days, a 200 ml sample was removed and placed in a clean container. The pH was measured and recorded. $AlCl_3$ (1×10^{-2} M) was added to bring the aluminum concentration to the experimental level. The pH was measured again. If a change had occurred, the pH was corrected with 0.1 M NaOH. After 45 minute the pH was verified a final time. 10 ml samples of the culture were transferred to 30 ml

polycarbonate tubes. An hour after the addition of aluminum, copper (2×10^{-4} M CuSO_4) was added to the tubes. The algae were exposed to the copper for an additional hour before the assays were performed. Replicate tubes of aluminum free cultures were assayed as a control.

Five tubes could be assayed at one time. The first 10 ml sample was spiked with 0.2 ml of 1×10^{-3} M pNPP. After gentle mixing, 3 ml were filtered into a 1 cm path length spectrophotometer cuvette. The absorption at 410 nm was measured and recorded. This process was repeated for each of the five samples. After the initial reading of the fifth sample was recorded, a second round of measurements was begun. The exact time of each filtration was recorded. Approximately six minutes elapsed between repeated sampling of any single tube. After the absorption of each sample had been measured three times, the enzyme activity was calculated.

Because the number of cells in a culture varied, the absolute enzyme activity of the control samples was not consistent between experiments. The effect of metal addition is reported compared to the enzyme activity the two control assays. As the exact rate differed on different days, the results are given as percent of control activity.

The chemical speciation for each concentration combination was estimated using a Pascal version of MICROQL (Westall et al 1976; Westall 1985). Due to a limited memory

capacity, a condensed Fraquil data matrix was used (Table V). Components which were shown by MINIQL (Westall et al 1976) not to significantly affect metal speciation were excluded.

RESULTS

The observed effect of increasing aluminum changed as the concentration of copper increased (Figure 10). In a copper free culture, the addition of 2×10^{-5} M aluminum inhibited the activity of alkaline phosphatase by sixteen percent. This concentration of aluminum led to an activity decrease of 88 percent if the copper concentration was 8×10^{-6} M (Table VIII).

The EC₅₀ (metal concentration required to inhibit enzyme activity by 50 percent) for copper was determined by graphing the data (rate v. copper) and interpolating for copper concentration at 50% control activity (Figure 11). The copper EC₅₀ decreased as the aluminum concentration was increased.

The activity of the enzyme in the presence of both metals was found to be related ($r^2=0.89$) to the log cupric ion concentration (Figure 12). The effective range of pCu is between 7.7 and 5.7. In the absence of aluminum, there is not enough data to clearly state whether the cupric ion or total copper concentration controls toxicity. Enzyme activity did not relate to either total aluminum or pAl.

Figure 10. The activity of Scenedesmus alkaline phosphatase as a function of aluminum in the presence and absence of copper.

■ - no copper

○ - 8 x 10⁻⁶ M copper

TABLE VIII

ACTIVITY OF SCENEDESMUS ALKALINE PHOSPHATASE (upper)
AND CUPRIC ION ACTIVITY (lower) AS A FUNCTION OF
ALUMINUM AND COPPER CONCENTRATION

		Copper ($\times 10^{-6}$)			
		0	4	8	12

Aluminum ($\times 10^{-5}$)	0	100	93	67	0
		13.0	11.6	9.1	5.9
	0.5	105	105	39	0
		13.0	7.6	6.1	5.6
	1.0	93	62	10	0
		13.0	6.4	5.8	5.5
	2.0	84	70	8	0
		13.0	6.0	5.6	5.4

activity = percent of control

Figure 11. The activity of *Scenedesmus* alkaline phosphatase as a function of copper concentration at four aluminum concentrations.

● - no aluminum

○ - 5×10^{-6} M aluminum

□ - 1×10^{-5} M aluminum

■ - 2×10^{-5} M aluminum

Figure 12. The activity of *Scenedesmus* alkaline phosphatase as a function of cupric ion activity.

DISCUSSION

The concentration of copper was found to be important in determining aluminum toxicity. Aluminum alone had little effect on the activity of Scenedesmus alkaline phosphatase. In the presence of copper, increasing the concentration of aluminum led to a large decrease in activity.

The results support the hypothesis that chemical speciation is critical in determining metal toxicity. A strong correlation was found between enzyme activity and cupric ion concentration. The effect of aluminum concentration on the level of the cupric ion can be seen in Figure 13. At a set copper concentration, the level of the toxic copper species increases with aluminum concentration.

Although it has been shown that Scenedesmus alkaline phosphatase and the growth rate of the alga are both sensitive to cupric ion activity, I am not suggesting that the inhibition of the enzyme is the cause of growth rate inhibition. Alkaline phosphatase is acting as a model enzyme system. The decrease of growth may be due to the partial inhibition of many enzymes. The important point is that the effect of changes in metal speciation have been shown with an organism and with a single enzyme.

This experiment increases the number of alkaline phosphatase systems known to be sensitive to cupric ion

Figure_13. The activity of the cupric ion increases with aluminum concentration. The copper concentration is sent at $8 \times 10^{-6} \text{ M}$.

activity. Rueter (1982) found the enzyme to be an effective biochemical marker of toxicity in three increasingly complex environments. Comparable results were obtained in studies with the isolated enzyme, cultures of marine diatoms, and natural assemblages of freshwater phytoplankton.

The bioassay developed in this research could be used to quickly estimate the effect of changing metal concentrations in natural waters. A concentrated culture of Scenedesmus could be added to a water sample. The activity of the enzyme would be monitored before and after the addition of metals. If a portable spectrophotometer (e.g. Hach Co.) is available, this work could be performed in the field.

It has been shown that enzymes exposed to the environment are sensitive to changes in the metal chemistry. Similar effect may occur in other organisms. The observed sensitivity of fish to increased aluminum (Cronan and Schofield 1979) may be because of the number of important enzymes present in gill membranes.

CONCLUSION

The results obtained with Scenedesmus alkaline phosphatase were similar to those obtained with the isolated enzyme. In the presence of aluminum and copper,

enzyme activity is significantly correlated with cupric ion activity.

Aluminum alone was shown to have little effect on the enzyme. If copper was present in the medium, enzyme activity decreased as the aluminum concentration increased.

CHAPTER V

THE USE OF A COMPUTER MODEL TO PREDICT THE INTERACTION OF COPPER AND ALKALINE PHOSPHATASE

INTRODUCTION

This research has shown that metal speciation determines the effect of copper on alkaline phosphatase activity. Copper and zinc ions compete for the metal binding site of the enzyme (Foy et al 1978). Alkaline phosphatase can be considered as a ligand with different affinities for transition metals. The goal of this section is to model the interactions between the enzyme, zinc and copper in a way to predict the results of perturbations in aquatic chemistry.

THEORY

In chapters II through IV, the computer program MICROQL (Westall et al 1976) was used to model chemical interactions and estimate metal speciation. This chapter extends the use of MICROQL to model the actual interactions between alkaline phosphatase and the metals zinc and copper. This work is based on the assumption that the zinc binding site of the enzyme acts as a ligand. A knowledge of the outcome of competition between the metals for the binding site should allow one to predict the enzyme activity at a given metal concentration.

The concentration of the alkaline phosphatase-zinc-pNPP (E-Z-P) complex should be the factor controlling the measured enzyme activity (Lehninger 1982). A conditional stability constant for E-Z-P will be determined. An accurate formation constant is required to estimate the complex concentration. The formation of the alkaline phosphatase-zinc-pNPP complex depends on the interaction of both Zn^{+2} and pNPP with the enzyme. Kinetic experiments were performed in which the concentration of either zinc or pNPP was held constant at a saturating level, while the concentration of the other was varied. Using the Michaelis-Menten equation, the K for for both the substrate and the metal was determined. K is an indicator of the stability constant between an enzyme and another substance. If it is assumed that the binding of zinc and pNPP are sequential events:

1. Enzyme + Zn $\rightleftharpoons$ Enz-Zn
2. Enz-Zn + pNPP $\rightleftharpoons$ E-Z-P

the stability constants of the tricomponent complex should be the product of the enzyme-zinc and enzyme-pNPP constants (Rueter, pers. comm.). In other words, the log K of the E-Z-P complex should be the sum of the log K of each substance determined experimentally.

The stability constant between alkaline phosphatase and copper was determined in the second part of the experiment. Copper is thought to inhibit alkaline

phosphatase by displacing the zinc (Foy et al 1978). The two metals compete for the same site on the enzyme. An initial assumption was that the relative affinities of copper and zinc for the binding site should be similar to other organic ligands. Knowledge of the zinc stability constant should allow one to predict the copper constant by comparisons with tables of known values (Smith and Martell 1975). Although it was supported by the results, this assumption may not always be valid. With stability constants for copper and zinc, MICROQL could be used to predict the outcome of competition of metals for binding to alkaline phosphatase.

METHODS

The activity of alkaline phosphatase was assayed using the method described in Chapter III. The buffer (1×10^{-3} NaHCO_3 , 1×10^{-5} NTA) used in determining the K_m of pNPP contained 5×10^{-6} ZnCl_2 . Five substrate concentrations between 1.1×10^{-6} and 4.4×10^{-5} M were tested (Table IX). In determining the K value for zinc, the pNPP concentration was maintained at 4.4×10^{-5} M. The enzyme was exposed to low zinc for four hours to allow zinc equilibrium to be reached between the enzyme and NTA.

The chemical speciation for each zinc concentration was estimated using a Pascal version of MICROQL (Westall 1985). The program was run on a Zenith 151 PC (Zenith data

TABLE IX
A COMPARISON OF EXPERIMENTAL RESULTS
WITH THOSE PREDICTED BY MICROQL

Zn conc $\times 10^{-7}$	experimental enzyme activity		predicted enzyme activity	experimental percent Vmax		predicted percent Vmax
1	4.9	5.6	4.3	26	30	23
2	9.5	7.7	7.1	50	41	38
4	11.6	9.5	10.5	61	50	56
6	13.7	13.5	12.5	73	70	66
8	15.4	11.8	13.6	81	63	72
10	14.7	15.1	14.6	78	80	77
20	10.8	11.3	11.9	80	84	88

pNPP conc. $\times 10^{-6}$	experimental enzyme activity		predicted enzyme activity	experimental percent Vmax		predicted percent Vmax
1.1	5.0	5.0	7.6	27	27	40
2.2	9.8	10.5	10.8	52	56	58
4.4	13.0	--	13.8	73	--	73
9.6	14.0	14.7	16.2	74	78	86
44.0	18.2	--	16.9	96	--	90

systems). A matrix (Table V) was set up for the chemical components of the buffer. The stability constants used were obtained from MINEQL data files (Westall et al 1976). After the K was determined for pNPP and Zn^{+2} , the matrix was expanded to include pNPP and alkaline phosphatase as components. In addition to the enzyme-zinc-pNPP complex, another species created by matrix expansion was enzyme-H. It was assumed that proton binding to the enzyme is similar to NTA (see page 67) so the NTA-H formation constant was used.

The procedure for testing various formation constants of the E-Z-P complex was to set the pNPP concentration at 4.4×10^{-5} and the zinc concentration at 3.5×10^{-7} . Since these component concentrations gave an enzyme activity of one half V_{max} , the E-Z-P complex should be 50 percent of the total enzyme present. The calculated $-\log K$ for pNPP was 5.75 while it was 11 for Zn^{+2} , so the sum 16.75 was the first log formation constant tried for the E-Z-P complex. The E-Z-P to total enzyme ratio was determined for K values between 15.75 and 18.50. At $\text{pK}=17.85$, the program estimated a ratio of 0.53. As this compared favorably with experimental ratio of 0.50, a computer titration of zinc was performed. At each zinc concentration, the percent enzyme saturation was compared to the percent V_{max} determined experimentally. The zinc concentration was then set at 5×10^{-6} zinc and a computer titration of pNPP was

performed. Again the estimated enzyme saturation was compared to experimental results.

The experimentally determined binding constant between Zn^{+2} and alkaline phosphatase ($pK = 11$) was similar to the standard constant between Zn^{+2} and NTA ($pK = 10.66$, Smith and Martell 1975). This suggests the metal binding site of the enzyme may be similar to NTA. This information was used to predict that the binding copper to the enzyme should have a stability similar to Cu-NTA ($pK = 12.94$, Smith and Martell 1975). The MICROQL matrix was expanded to include the species Cu-enzyme. A computer titration was performed to predict the effect of increasing copper on the concentration of E-Z-P. The results were compared to enzyme activity determined experimentally. Six copper concentrations between 2×10^{-5} and 1×10^{-4} M were tested (Table X). The zinc concentration was maintained at 2×10^{-6} M, and the pNPP concentration used was 4.4×10^{-5} M.

The data were also used to get an improved estimate of the Cu-enzyme binding constant. The experimental conditions which gave an activity of fifty percent of V_{max} were set in the computer. The K was varied slightly and an estimation of enzyme saturation was calculated until an enzyme saturation of fifty percent was obtained.

TABLE X
A COMPARISON OF EXPERIMENTAL RESULTS
WITH THOSE PREDICTED BY MICROQL

Enz-Cu pK = 12.96

copper conc. $\times 10^{-5}$	experimental enzyme activity		predicted enzyme activity	experimental percent V _{max}		predicted percent V _{max}
1	17.2	17.2	18.0	92	91	95
2	13.8	16.6	10.6	73	86	56
4	7.4	4.9	5.5	39	26	29
5	7.1	5.7	4.5	38	30	24
6	1.4	2.5	3.8	6	13	20
8	1.0	2.1	2.8	5	11	15
10	1.1	0.7	2.3	6	4	12

Enz-Cu pK = 12.72

1	17.2	17.2	18.3	92	91	97
2	13.8	16.6	12.9	73	86	68
4	7.4	4.9	7.6	39	26	40
5	7.1	5.7	6.3	38	30	33
6	1.4	2.5	5.4	6	13	29
8	1.0	2.1	4.2	5	11	22
10	1.1	0.7	3.3	6	4	18

RESULTS

The K_m of pNPP in the experimental buffer was 1.9×10^{-6} M ($pK=5.75$). The pK of Zn^{+2} was 11. The first pK tested, 16.75, gave estimated enzyme $\frac{E-Z-P}{E+Z-P}$ ratio of 0.21. Lowering the pK decreased the relative concentration of the complexed enzyme, while increasing the pK had the reverse effect. At the experimental conditions that gave an activity of one half V_{max} , a pK of 17.85 led to an estimation of enzyme saturation of 0.53. Table IX and Figure 14 show the comparison of the experimental results with the calculated activities. A linear regression shows a relationship of 0.96 between the experimental results and the results of the simulated zinc titration. The relationship was 0.97 for the pNPP experiments.

The predicted results of copper addition and experimentally determined enzyme activities are presented in Table X. The Cu-NTA stability constant ($pK = 12.96$) is a good initial estimation for the interaction between copper and alkaline phosphatase. The model accurately predicts the effective range of copper concentrations. Copper concentrations below 1×10^{-5} M decreased the activity of the enzyme by less than ten percent (Table X). Alkaline phosphatase activity is inhibited if the level of copper increases further. The exact enzyme activity at a particular copper concentration was not accurately predicted. The model suggested lower activity at low copper levels ($< 5 \times 10^{-5}$ M

Figure_14. A comparison of experimental and predicted alkaline phosphatase activity ($\times 10^{-7}$ M pNPP / min) at six zinc concentrations.

● - experimental

+ - predicted

Figure 15. A comparison of experimental and predicted alkaline phosphatase activity ($\times 10^{-7}$ M pNPP / min) at four pNPP concentrations.

● - experimental

+ - predicted

copper) and greater enzyme activities at higher copper concentrations.

If the $pK = 12.72$, the model predicts fifty percent enzyme activity at the experimental copper EC_{50} (3×10^{-5} M). The results of the computer titration using this value are in Table X. The predicted activities are closer to the experimental results for the samples with $\leq 5 \times 10^{-5}$ M copper. The model still underestimates the effect of higher copper concentrations (Figure 16)."

DISCUSSION

MICROQL can be used to predict the effect of substrate and zinc concentration on the activity of alkaline phosphatase. The assumption that the stability constant of the enzyme-zinc-pNPP complex is a function of the zinc-enzyme and pNPP constants was supported. The difference between the sum of the pK s for zinc and pNPP (16.75) and the value which accurately predicted experimental results (17.85) differed by only six percent. The success of the model suggest that the binding of zinc is a reversible process. This idea is also supported by earlier results (Figure 9, Chapter III).

This model can be used to predict the effective range of cupric ion activity. Inhibition is the result of competition between zinc and copper for the metal binding site of alkaline phosphatase. MICROQL is effective in

Figure 16. A comparison of experimental and predicted alkaline phosphatase activity ($\times 10^{-7} \text{ M pNPP / min}$) at seven copper concentrations.

○ - experimental

■ - predicted

— - non-competitive inhibitor curve

estimating the outcome of metal-metal interaction if the stability constants between the components are known. This research has shown that satisfactory results are obtained if experimentally determined constants are used.

Predicted enzyme activity is greater than actual activity at high ($> 5 \times 10^{-5}$ M) copper concentrations. Figure 15 is a graph of activity as a function of copper concentration. A comparison of the predicted results with theoretical enzyme inhibition equations (Lehninger 1982) indicates the model assumes copper acts as a non-competitive inhibitor. The requirements of a non-competitive inhibitor are that its binding to the enzyme prevents product formation, it does not bind to the active site, and finally the inhibitor binds reversibly with the enzyme. MICROQL also assumes that all reactions are reversible. Although the formation of the copper-enzyme complex may be chemically reversible, inhibition has not been shown to be a reversible process.

The metal binding site of alkaline phosphatase has ligand characteristics similar to Nitrilotriacetic acid (NTA). The NTA-Me pK is 10.66 for zinc and 12.92 for copper (Smith and Martell 1975). The experimentally determined enz-Me pKs are 11 and 12.72 respectively. The structure of NTA is three acetic acids bound to a single nitrogen atom. The carboxyl groups are involved in metal complexation. This suggests the carboxyl groups of the

amino acids glutamic acid and/or aspartic acid are likely to be involved in the metal binding site of alkaline phosphatase.

CONCLUSION

This investigation used the computer program MICROQL to model the interactions between alkaline phosphatase and the metals zinc and copper. The goal is to be able to predict the effect of perturbations in the aquatic chemistry.

Experimental data and the computer program were used to determine the conditional stability of alkaline phosphatase with zinc, pNPP, and copper. The program accurately predicts enzyme activity at a given concentration of the substrate or native metal. The model also predicts the inhibitory range of cupric ion activity. The exact enzyme activity at a given copper concentration could not be predicted since the model assumes inhibition by copper to be reversible.

The conditional stability constants determined for alkaline phosphatase are similar to the known values for NTA. This suggests the metal binding site of the enzyme is similar to the chelator.

CHAPTER VI

CONCLUSION

The goal of this research was to determine how the chemical interaction between aluminum and copper effects the green alga Scenedesmus quadricauda. The hypothesis was that the growth rate of the alga and the activity of its enzymes should be related to the concentration of the free cupric ion. In the presence of both metals, increasing the aluminum concentration can lead to copper toxicity. The results support this hypothesis.

A correlation ($r^2 = 0.87$) was obtained between the growth rate of Scenedesmus and cupric ion activity. Similar correlations were obtained between the cupric ion and the activity of bacterial ($r^2 = 0.85$) and Scenedesmus ($r^2 = 0.89$) alkaline phosphatase. Correlations were not found between any other chemical species and the growth rate or activity of the Scenedesmus enzyme. The bacterial alkaline phosphatase was also correlated to the total copper concentration ($r^2 = 0.88$).

It was found that the computer program MICROQL (Westall et al 1976) can be used to model interactions of enzymes and metals in aqueous solutions. This supports the hypothesis that enzymes act as ligands. The model requires the determination of conditional stability constants between the enzyme and the metals. One can predict the effect of changing metal concentrations on enzyme activity.

An unexpected result was the apparent protective effect of low aluminum concentrations. The growth rate and enzyme activity were greater at a given cupric ion concentration if some aluminum was present (Figure 2, Table VI). This may have either a chemical or biological explanation. MICROQL may not accurately predict the interactions of dilute aluminum solutions. The apparent protective effect may actually be the comparison of the biological parameters to an overestimation of the cupric ion activity. It is also possible that aluminum does protect from copper inhibition. As discussed in Chapter III, aluminum ion may bind to sensitive sites on enzymes. This binding does not inactivate the enzyme, but delays or prevents copper from binding.

This research has shown that aluminum exerts a toxic response by influencing the metal speciation of a system. In terms of applying the results to environmental problems, this observation is more important than the actual effect on Scenedesmus. As the concentration of aluminum is likely to increase in lakes affected by acid rain, an understanding of the nature of toxicity is required for optimal management of the problem. This thesis has presented a mechanism of toxicity that has not been discussed in the literature.

The effect of acid rain on algae is not of much concern (Dillon et al 1984). Scenedesmus was chosen as a

test organism because of the ease of maintaining a chemically defined medium. The proposed mechanism should now be tested in more complicated systems. A study with zooplankton could determine if the effect of metal speciation is similar between plants and animals. The most important organisms affected by acid rain are fish. Although maintaining a chemically defined medium is difficult, a similar study with fish could be directly applied in the environment.

The results of this research suggest the type of lake that may be sensitive to increased aluminum. If changes in speciation are to be significant, the concentration of a toxic metal must be high relative to the number of complexation sites or ligands available. A lake that is low in dissolved organic material but with a high concentration of copper or other toxic metal would be susceptible to changes in aluminum concentration. Field studies in such a lake could be done with some of the techniques used in this research. Possible experiments include studies in which the metal concentration is varied in individual enclosures in lakes. This would allow the testing of the effect of metal interactions on natural populations. The use of alkaline phosphatase assays could be used to quickly estimate the interaction between the environmental chemistry and biochemistry in natural water samples.

SUMMARY

This research was performed because of the increase in aluminum in lakes impacted by acid rain. The result suggests that aluminum can exert its effect by interacting with the aquatic chemistry of the system. Aluminum alone was found to have little effect in all the systems investigated. But in the presence of copper, increasing aluminum concentration had a large effect. In these experiments, aluminum displaced copper ions from the chelators present in solution. The copper ion is the actual toxic agent. Of course, copper is not responsible for the loss of organisms in all lakes affected by acid rain. The exact toxic agent would depend on the relative concentration of all metals in a particular body of water. The important point is that the overall metal chemistry must be taken into account before a direct cause and effect between an increase in aluminum and toxicity can be established.

APPENDIX A

COPPER AND LACTATE DEHYDROGENASE: INHIBITION OF AN INTERNAL ENZYME

INTRODUCTION

An internal enzyme should show different sensitivity to metal toxicity than a enzyme exposed to the environment. Due to charged or polar nature of many organic molecules, the cytoplasm has numerous potential metal binding sites. The binding of metals to "safe" or unaffected sites decreases the free metal ion activity. A greater total metal concentration may be required to inhibit an internal enzyme. The time frame of metal inhibition is different for an internal and external enzyme (Rothstein 1959). External enzymes are immediately exposed to the changes in the chemistry of the environment. A metal must diffuse or be taken up into a cell before it can affect an internal enzyme.

My goal was to study the effects of aluminum and copper on an internal enzyme and compared the results to those obtained for alkaline phosphatase. I chose to try lactate dehydrogenase (LDH) because it is a zinc metallo-enzyme (Ochiai 1977) that should be present in algal cells. Unfortunately, a method could not be found that would allow for the extraction of the enzyme while maintaining defined conditions suitable for trace metal studies. This appendix

discusses the work that was performed with isolated LDH and describes the extraction techniques attempted.

METHODS

Isolated LDH Experiments- The enzyme lactate dehydrogenase (LDH) catalyzes the reaction:

LDH activity is measured by following the disappearance of NADH in a spectrophotometer at 340 nm.

The first goal was to determine the $V_{\max}$, K , and enzyme saturation concentration. Lactate dehydrogenase (Sigma Co), was diluted to 1.5 units per milliliters in 1×10^{-1} M Tris. Six concentrations of pyruvate (Sigma Co) were tested, 2.8×10^{-6} , 5.5×10^{-6} , 1.1×10^{-5} , 2.1×10^{-5} , and 3×10^{-5} M. Two milliliters of a 1×10^{-1} M Tris solution was added to a cuvette. To begin each assay, 0.1 ml of the LDH-Tris solution was added to the cuvette. The absorption of each sample was recorded every two minutes for ten minutes. This experiment was repeated using a 1×10^{-3} M buffer.

The effect of three concentrations of copper was determined. Copper (5×10^{-2} M CuSO_4) was added to 20 ml samples of 1×10^{-1} M Tris to bring the final concentration to 2.5×10^{-4} , 2.5×10^{-5} , and 2.5×10^{-6} M. Two milliliters of the LDH-Tris solution described above was added to each sample. Two milliliter subsamples were used in the enzyme

assays. 0.25 ml of the pyruvate and 0.25 ml of NADH were added to the cuvettes to begin the assays. The activity of the LDH was determined at time zero, 2 hours, 8 hours and 24 hours.

The effect of copper in a solution of 1×10^{-3} M Tris was then investigated. Six concentrations of copper between 1×10^{-8} and 1×10^{-3} M as well as a copper free control were tested. For each experimental concentration, 5×10^{-2} M CuSO_4 was added to ten milliliters of the Tris buffer. Twelve units of LDH were added to each sample. Assays were performed in 1×10^{-1} M Tris. A similar experiment was run testing the effects of four aluminum concentrations between 1×10^{-7} and 1×10^{-4} M.

LDH Extraction techniques- Standard biochemical enzyme isolation techniques (LeJohn and Stevenson 1975) could not be used to assay the activity of Scenedesmus lactate dehydrogenase. Metal speciation would not be maintained in a multi-step, multi-buffer process. The first goal was to determine if LDH activity could be analyzed if the cells were gently ground. Scenedesmus quadricauda was grown in enriched Fraquil ($2 \times \text{NaNO}_3$, K_2HPO_4). For each sample, 100 ml of a week old culture was centrifuged (International Centrifuge Co.) at 2000 rpm for five minutes. The pellet was transferred to a cold (5°C) mortar. Three milliliters of 50°C 1×10^{-1} M Tris were added and the cells were ground for two minutes. The mixture was centrifuged at 2000 rpm

for five minutes. Two milliliters of the supernatant was transferred to a cuvette. The change in absorption was monitored for five minutes. 0.25 ml of NADH was added and the sample was monitored for an additional five minutes. These steps were taken to determine the background change in NADH concentration. The enzyme activity assay was begun by adding 0.25 ml of the pyruvate solution. The change in absorption was monitored for thirty minutes.

LeJohn and Stevenson (1975) suggest that LDH can be extracted in a Tris(1×10^{-1} M)-acetate(2×10^{-3} M) buffer. This solution was used in the test of another extraction technique. 100 ml of culture was filtered through a 25mm glass fiber filter (Gelman Sciences, Type A/E). The filter and three milliliters of the Tris-acetate buffer were placed in a 15 ml glass tissue grinder (Wheaton Scientific). Some of the samples were ground using an electrical motor driven teflon pestle, while other were ground by hand using a ground glass pestle. Various grinding times were tested. After a sample had been ground, the mixture was centrifuged at 2000 rpm for five minutes. The LDH activity was assayed using the method described above.

As some enzymes are inactive or present in low concentration in the absence of their substrate, the effect of pre-exposure to pyruvate was tested. A stationary phase culture was split into two 100 ml samples. One milliliter of 2.27×10^{-4} M pyruvate was added to one of the samples.

After twenty-four hours both cultures were assayed for LDH activity. The filtration technique discussed above was used. In all cases the filters were hand ground for one minute.

Sullivan and Volconi (1976) explain how internal enzymes of algae can be assayed by dissolving the cell membrane. Two solvents were tested. In the first method, 0.001 v/v toluene in 1×10^{-1} M Tris was prepared. 100 ml of a stationary phase culture was centrifuged and the pellet was added to five milliliters of the Tris-toluene solution. The sample was mixed for five seconds in a vortex test tube mixer every 30 seconds for five minutes. The sample was then centrifuged (2000 rpm, five minutes). The toluene-Tris solution was removed by suction and replaced with five milliliters of a 1×10^{-1} Tris solution. Pyruvate and NADH (0.5 ml each) was added and the 2.5 ml of the sample was filtered into a spectrometer cuvette. After one hour the remainder of the sample was filtered and its absorption recorded. In the second method, two drops of the detergent Triton-x was added to five milliliters of a culture. After five minutes of gentle mixing, the sample was centrifuged (2000 rpm, five minutes). The Triton-x solution was replace with five milliliters of the Tris buffer. The above assay method was repeated.

RESULTS

Isolated LDH Experiments- The experiment using 1×10^{-1} M Tris indicates that a pyruvate concentration of 2×10^{-5} M was required to saturate the isolated lactate dehydrogenase. The diluted enzyme in the absence of copper lost twenty five percent of its activity in twenty four hours. In the presence of copper less of a decrease was noted.

LDH could be maintained in a 1×10^{-3} M Tris solution, but an assay could not be run at this buffer concentration. Table 11 shows the results of copper exposure. All the copper concentrations tested had some effect on enzyme activity. 1×10^{-8} M copper had little effect over the first eight hours, but after twenty four hours the activity was lower than the control rate. At time zero, the activity of the enzyme exposed to 1×10^{-7} M copper was only seventy five percent of the control. The activity continued to decrease with time. There was no activity after twenty four hours. At time zero the LDH exposed to 1×10^{-6} M copper had an activity of forty percent of the control. No activity was found at eight hours. At a copper concentration of 1×10^{-5} M, the time zero activity was twenty five percent of control, There was total inactivation at thirty minutes. Higher copper concentration caused an immediate inactivation of LDH. LDH activity did not depend on aluminum concentration. Samples exposed to low aluminum concentration had activities lower

than control, while those exposed to 1×10^{-5} and 1×10^{-4} M aluminum had activities higher than control after twenty four hours.

LDH Extraction techniques- None of the extraction methods tested were effective. Although activity could be determined in some samples, reproducible data was not obtained. Pre-exposure to pyruvate did have a noticeable effect. The measured change in absorption was 70-80 times greater than the pyruvate free sample, but the absorption increased instead of decreased as expected. This indicates another enzyme system was activated by the addition of pyruvate.

CONCLUSION

Internal and external enzymes should have different responses to metal toxicity. My goal was to investigate the effect of the aquatic chemistry of aluminum and copper on the activity of lactate dehydrogenase in Scenedesmus. Results obtained for this internal enzyme were to be compared to results using the external enzyme alkaline phosphatase. Experiments were performed in an effort to develop an extraction technique that would allow the reproducible measurement of LDH activity while maintaining conditions suitable for trace metal studies. None of the techniques attempted produced consistent results.

REFERENCES

- Baker, Joan and Schofield, Carl 1982. Aluminum toxicity to fish in acidic water. *Water, Air, Soil Poll.* 18. 289-309
- Bold, H. and Wynne, M. 1978. Introduction to the algae: structure and reproduction. Prentice-Hall Inc.
- Burrows, W.D. 1977. Aquatic aluminum: chemistry, toxicology, and environmental prevalence. *CRC Critical Reviews in Environmental Control*, 167-216
- Cohen and Wilson 1966. Measurements of zinc dissociation constants of alkaline phosphatase from E. Coli with zinc ion buffer. *Biochem.* 5(3) 904-909
- Coleman, Joseph and Gettins, Peter 1983. Molecular properties and mechanism of alkaline phosphatase. in *Zinc Enzymes*, Thomas Spiro ed., Wiley-interscience publication. Chap. 5
- Cronan, Christopher and Schofield, Carl 1979. Aluminum leaching response to acid precipitation: effects on high-elevation watersheds in the Northeast. *Science* 204(20) 304-305
- Currie, David and Kalft, Jaap 1984a. A comparison of the abilities of freshwater algae and bacteria to acquire and retain phosphorus. *L&O* 29(2) 298-310
- Currie and Kalft 1984b. The relative importance of bacterioplankton and phytoplankton in phosphorus uptake in freshwater. *L&O* 29(2) 311-321
- Davey, E., Morgan, M. and Erickson, S. 1973. A biological measure of the copper complexation capacity of seawater. *L&O* 18(6) 993-997
- Dillion, P., Yan, N. and Harvey, H. 1984. Acidic deposition: effects on aquatic ecosystems. *CRC Crit. Rev. Env. Control* 13(3), 197-194
- Driscoll, C. 1984. A procedure for the fractionation of aqueous aluminum in dilute acidic waters. *Intern. J. Environ. Anal. Chem.* 16 267-283
- Driscoll, C., Baker, J., Bisogni, J., and Schofield, C. 1984. Aluminum speciation in dilute acidic surface waters of the adirondack region of New York State. in *Geological aspects of acid deposition*. Bricker, O. ed Butterworth Publishers, Chap 4 55-75

- Fontaine, Thomas 1984. A non-equilibrium approach to modeling metal speciation in acid, aquatic systems: theory and process equations. *Ecological Modeling* 21. 287-313
- Foy, C., Chaney, R., and White, M. 1978. The physiology of metal toxicity in plants. *Annu. Rev. Plant Physiol.* 29: 511-566
- Gamble, Donald, Schnitzer, M., Kerndorff, H, and Langford, C. 1983. Multiple metal ion exchange equilibria with humic acid. *Geochimica et Cosm. Acta* 47 1311-1323
- Gatcher, Ron, Davis, Joan, & Mares, Antonio 1978. Regulation of copper availability to phytoplankton by macromolecules in lake water. *E,S,&T*, 12 1416-1421
- Giesy, J P 1983. Metal binding capacity of soft, acid, organic-rich waters. *Toxicology and environ. Chem.* 6 203-224
- Giesy, John, Newhall, Avis, and Leversee, Gordon 1983. Copper speciation in soft, acid, humic, waters: effects on copper bioaccumulation and toxicity to Simocephalus serrulatus. *Sci Tot. Environ.* 28 23-26.
- Gillespie, Paul and Vaccaro, Ralph 1978. A bacterial bioassay for measuring the copper-chelation capacity of seawater. *L&O* 23(3) 543-547
- Goldman, Charles and Horne, Alexander 1983. *Limnology*. McGraw-Hill. Chap 9.
- Haug, Alfred 1984. Molecular aspects of aluminum toxicity. *CRC Critical reviews in plant science.* 1(4) 345-373
- Helliwell, Stuart 1983. Speciation and toxicity of aluminum in a model fresh water. *Sci. & Tech. Letters* 4 141-144
- Huntsman, S. and Sunda, W. 1980. The role of trace metals in regulating phytoplankton growth: with emphasis on Fe, Mn, and Cu. in *Physiological Phytoplankton Ecology*, Morris, I. ed.
- Johnson, Noye 1984. Acid rain neutralization of by geological materials. in *Geological aspects of acid deposition*. Bricker, O. ed. Butterworth Publishers Chap 3 37- 53

- Johnson, N., Likens, G., Feller, M., & Driscoll 1984.
Acid rain and soil chemistry. *Science* 225 1424-1425
- Kerndorff, H., and Schnitzer, M., 1980. Sorption of metals on humic acid. *Geochimica Acta*. 44 1701-1708
- Kuenzler, E. and Perras, J. 1965. Phosphatases of marine algae. *Biol. Bull.* 128: 271-284
- LaZerte, B.D. 1984. Forms of aqueous aluminium in acidified catchments of central Ontario: a methodological analysis. *Can J Fish. Aquat. Sci.* 41 766-776.
- Lehninger, A. 1982. Principles of Biochemistry. Worth Publ.
- LeJohn, H.B. and Stevenson, R.M. 1975. D(-)-Lactate Dehydrogenase from Fungi. *Methods in Enzymology* vol XLI, W. Wood ed. Academic Press .
- McKnight, Diane 1981 . Chemical and biological process controlling the response of a freshwater ecosystem to copper stress: a field study of CuSO₄ treatment of Mill Pond Reservoir, Burlington, Massachusetts. *L & O* 26(3) 518-531
- McKnight, D. Chisholm, S. Harleman, D. 1983. CuSO₄ treatment of Nuisance blooms in drinking water reservoirs. *Environ. Management* 7(4) 311-320
- McKnight, D. and Morel, F. 1979. Release of weak and strong copper complexing agents by algae. *L&O* 24(5) 823-837
- Morel, F. Rueter, J. Anderson, D. and Guillard, R. 1979. Aquil: a chemically defined phytoplankton culture medium for trace metal studies. *J Phycol.* 15, 135-141
- Ochiai, E. 1977. Bioinorganic Chemistry: an introduction Allyn & Bacon Publ.
- Perdue, E.M. 1979. Solution Thermochemistry of Humic substances. In *Chemical Modeling in aqueous systems*. E.A. Jenne ed.. Am. Chem. Soc. symposium Chap 5, 99-114
- Perdue, Edward, and Lytle, Charles 1983. Distribution model for binding of protons and metal ions by humic substances. *ES&T* 17 (11) 654-660

- Petersen, Richard 1982. Influence of copper and zinc on the growth of a freshwater alga, Scenedesmus quadricauda: the significance of chemical speciation. ES&T 16 443-447
- Peterson, H., Healey, F.P., and Wagermann, R. 1984. Metal toxicity to algae: a highly pH dependent phenomenon. Can. J. Fish. Sci. 41, 974-979
- Rothstein, A. 1959. Cell membrane as site of action of heavy metals. Fed. Proc. 18, 1026-1038
- Rueter, John 1983. Alkaline phosphatase inhibition by copper: implications to phosphorus and use as a biochemical marker of toxicity. L&O 28(4) 743-748
- Rueter, J. and Morel 1981. The interaction between zinc deficiency and copper toxicity as it affects the silicic acid uptake mechanisms in Thalassiosira pseudonana. L&O 26(1) 67-73
- Saar, R. and Weber J., 1982. Fulvic acid: modifier of metal-ion chemistry. E,S, & T 16 (9)510-517
- Schofield, Carl and Trojnar, John 1980. Aluminum toxicity to brook trout (Salvelinus fontinalis) in acidified waters. in Polluted Rain. Toribara, T. ed. Plenum press 341-366.
- Shaw, T. and Brown V. 1973. The toxicity of some forms of copper to rainbow trout. Wat. Res. 377-382
- Smith and Martell 1975. Critical Stability Constants, vol 2 & 3. Plenum Press.
- Steemann-Nielsen and Wium-Andersen 1970. Copper ions as poisons in the sea and freshwater. Marine Bio. 6 93-97
- Sullivan, C. and Volcani, B. 1976. Role of silicon in diatom metabolism. VII. Silicic acid-stimulated DNA synthesis in toluene-permeabilized cells of Cylindrotheca fusiformis. Exptl Cell Res. 98 23-30
- Sunda, WG and Lewis, J. 1978. Effect of complexation by natural organic ligands on the toxicity of copper to a unicellular alga, Monochrysis lutheri. L&O 23(5), 870-876
- Vallee, B.L. 1959. Metal and enzyme interactions: correlation of composition, function, and structure. Physiol. Rev. 39, 443.

- Westall, J.C. 1985. MICROQL: a chemical equilibrium program in PASCAL. Oregon State Univ.
- Westall, J.C. 1979. MICROQL: a chemical equilibrium program in BASIC. Swiss Fed. Insit. Tech.
- Westall, J.C. , Zachary, J.L., and Morel, F.M. 1976. MINEQL: a computer program for the calculation of chemical equilibrium composition of aqueous systems. Mass. Inst. Tech., R.M. Parsons Lab. Tech Note 18