Portland State University

PDXScholar

Dissertations and Theses

Dissertations and Theses

6-28-1989

A Comparison of the Maladaptive Behaviors of Normal, Language Delayed, and Late Talking Toddlers

Denise Elaine James
Portland State University

Follow this and additional works at: https://pdxscholar.library.pdx.edu/open_access_etds

Part of the [Speech and Hearing Science Commons](#)

Let us know how access to this document benefits you.

Recommended Citation

James, Denise Elaine, "A Comparison of the Maladaptive Behaviors of Normal, Language Delayed, and Late Talking Toddlers" (1989). *Dissertations and Theses*. Paper 3894.
<https://doi.org/10.15760/etd.5764>

This Thesis is brought to you for free and open access. It has been accepted for inclusion in Dissertations and Theses by an authorized administrator of PDXScholar. Please contact us if we can make this document more accessible: pdxscholar@pdx.edu.

AN ABSTRACT OF THE THESIS OF Denise Elaine James for the
Master of Speech Communication: Speech and Hearing Sciences
presented June 28, 1989.

Title: A Comparison of the Maladaptive Behaviors of
Normal, Language Delayed, and Late Talking Toddlers.

APPROVED BY THE MEMBERS OF THE THESIS COMMITTEE:

Rhea Paul, Chair

Mary E. Gordon

Joseph S. Kaplan

People use language to communicate their needs and intentions, to express emotions, and to form relationships. It seems likely that a disruption in children's language development would have a negative impact on their social development. There is extensive research that shows that school age children with delayed language are "at risk" for increased maladaptive behaviors (Cantwell and Baker, 1977).

Whether this is also true for children in the earliest stages of language development is not yet known.

The questions this study sought to answer were: 1) Is there a significant difference in the severity and frequency of maladaptive behaviors seen in language delayed children, children who were "late talkers," or children with normal language? and 2) Is there a significant difference among the three subject groups in terms of which behaviors parents are the most concerned about?

The subjects in this study ranged in age from 18 to 41 months. They included 34 children with delayed language, 12 children who were "late talkers," and 29 children with normal language. The behavior of the subjects was assessed using two parental questionnaires, the Childhood Personality Scale (CPS) (Cohen, 1975), and the Behavior Checklist for Toddlers (BCT) (Rescorla, 1984).

The first question was analyzed by calculating the means for each area of behavior assessed and comparing the means among the three subject groups. To determine if there were significant differences among the three subject groups, t-tests were conducted at the .01 level of significance.

The second question was answered by determining what percentage of points, out of the total possible, parents assigned to each area of behavior. The behaviors were then ranked in order according to which areas of behavior parents assigned the highest percentage of points, and to which

areas of behavior they assigned the lowest percentage of points.

In answering the first question, the CPS showed the language delayed subjects to have significantly more problems than the normal language subjects in the areas of hyperactivity/attention deficit disorder and conduct. The BCT showed the language delayed subjects to have significantly more problems with hyperactivity/attention deficit disorder and relationships. No statistically significant differences were found either between the language delayed subjects and the "late talkers" or between the normal language subjects and the "late talkers."

In answering the second question, the CPS showed no differences among the three subject groups. The BCT found only minor differences among the three subject groups, with the parents of the normal language subjects expressing the most concern about their children's health, and the parents of both the language delayed children and the "late talkers" expressing the most concern about their children's conduct.

A COMPARISON OF THE MALADAPTIVE BEHAVIORS
OF NORMAL, LANGUAGE DELAYED,
AND LATE TALKING TODDLERS

by

DENISE ELAINE JAMES

A thesis submitted in partial fulfillment of the
requirements for the degree of

MASTER OF SCIENCE IN SPEECH COMMUNICATION:
SPEECH AND HEARING SCIENCE

Portland State University
1989

TO THE OFFICE OF GRADUATE STUDIES:

The members of the Committee approve the thesis of
Denise Elaine James presented June 28, 1989.

Rhea Paul, Chair

Mary E. Gordon

Joseph S. Kaplan

APPROVED:

Theodore G. Grove, Chair, Department of Speech Communication

C. William Savery, Interim Vice Provost for Graduate Studies
and Research

ACKNOWLEDGEMENTS

I would sincerely like to thank Dr. Rhea Paul whose knowledge and research expertise made this study possible. Her patience and support are much appreciated.

I would also like to thank the other members of my thesis committee, Mary Gordon, and Dr. Joseph Kaplan for their valuable contributions. I also owe Mary Gordon a special thank you for her guidance as my academic advisor.

Finally, I am very thankful for the love and support of my family. My parents have been a constant source of support as I have completed my education. I am also truly grateful to my husband David James, whose support, encouragement, and endless patience made it possible for me to complete graduate school.

TABLE OF CONTENTS

	Page
ACKNOWLEDGEMENTS	111
LIST OF TABLES	vi
LIST OF FIGURES	vii
CHAPTER	
I INTRODUCTION AND STATEMENT OF PURPOSE	1
Introduction	1
Statement of Purpose	4
Definition of Terms	6
II REVIEW OF THE LITERATURE	8
Behavior Problems of Children with Speech and/or Language Impairments	8
Studies Differentiating Children with Speech Disorders from Children with Language Disorders	13
Maladaptive Behaviors of Children with Impaired Language Only	15
Language Delays in Behaviorally Disturbed Children	16
Summary	19
III METHODS	21
Subject Selection	21
Instruments	24
Procedures	30
Data Analysis	31

Statistical Analysis	32
Reliability and Validity	33
Reliability	
Validity	
IV RESULTS AND DISCUSSION	
Results	34
Discussion	43
V SUMMARY AND IMPLICATIONS	
Summary	52
Implications	54
Research Implications	
Clinical Implications	
SELECTED BIBLIOGRAPHY	59
APPENDICES	62
Appendix A	62
Appendix B	65
Appendix C	69
Appendix D	72

LIST OF TABLES

TABLE		PAGE
I	Demographic Data for the Subjects on the Childhood Personality Scale: Means or Percentages	22
II	Demographic Data for the Subjects on the Behavior Checklist for Toddlers: Means or Percentages	23
III	<u>t</u> -Tests Indicating Differences Between the Normal (N) and Language Delayed (LD) Subjects on the Childhood Personality Scale	35
IV	<u>t</u> -Tests Indicating Differences Between the Normal (N) and Late Talking (LT) Subjects on the Childhood Personality Scale	36
V	<u>t</u> -Tests Indicating Differences Between the Language Delayed (LD) and Late Talking (LT) Subjects on the Childhood Personality Scale	37
VI	<u>t</u> -Tests Indicating Differences Between the Normal (N) and Language Delayed (LD) Subjects on the Behavior Checklist for Toddlers	38
VII	<u>t</u> -Tests Indicating Differences Between the Normal (N) and Late Talking (LT) Subjects on the Behavior Checklist for Toddlers	40

VIII	<u>t</u> -Tests Indicating Differences Between the Language Delayed (LD) and Late Talking (LT) Subjects on the Behavior Checklist for Toddlers	41

LIST OF FIGURES

FIGURE	PAGE
1. Order in Which Parents Ranked Their Concerns About the Behaviors Displayed by Their Children on the Childhood Personality Scale .	42
2. Order in Which Parents Ranked Their Concerns About the Behaviors Displayed by Their Children on the Behavior Checklist for Toddlers	44

CHAPTER I

INTRODUCTION AND STATEMENT OF PURPOSE

INTRODUCTION

Much research has been devoted to learning how the behavior of children with language disorders differs from that of their non-language disordered peers. Language is an integral part of being human. It is the single most important factor which separates man from animals. People use language to communicate their needs and intentions, to express emotions, and to form relationships. It seems likely that a disruption in children's language development would have a negative impact on their social development. The research to date shows that a relationship between language development and social development does exist. Several studies report that school-aged language delayed children are at risk for developing behavioral problems, including conduct disorders, attention deficit disorders, difficulty forming relationships, and other maladaptive behaviors such as soiling and crying easily (Cantwell & Baker, 1977). Whether or not this is also true for younger children is not yet known.

The speech-language pathologist may question the need to address the problem of maladaptive behaviors in the lan-

guage delayed child. After all, the role of the speech-language pathologist is to remediate speech and language disorders, not behavioral disorders. However, there are a number of important reasons why speech-language pathologists must be concerned with behavioral disorders. First, how well children communicate has a major impact on how they are treated by both adults and other children who come into contact with them. Laughton and Hasenstab (1986) wrote, "An inability to communicate with other children interferes with the establishment of peer relationships. This, in turn, inhibits further social and linguistic expression" (p.183). Adults also respond more warmly to children who interact appropriately, and who follow the rules of language use. In turn, the way people respond to children and the input they give them will subsequently affect further language development.

Another important reason for studying the behavior of language disordered children is the possibility that having a language disorder may cause children to develop maladaptive behaviors in order to compensate for their inability to communicate effectively. Laughton & Hasenstab (1986) hypothesized that maladaptive behaviors such as hyperactivity and withdrawal may be attempts to compensate for poor communication skills. They hypothesize that the hyperactivity so often seen in language disordered children may be an attempt to describe with actions what they have difficulty

putting into words. Poor language skills may also result in withdrawal. Withdrawal may occur because language delayed children need more time to process what has been said, or because they wish to escape from a communicative situation which is too taxing. If maladaptive behaviors do develop as a result of the language delay, teaching children to communicate effectively may be a crucial part of preventing the occurrence of such behaviors.

Third, the clinician must be prepared to deal with problem behaviors if progress is to be made. Children who are hyperactive or inattentive will waste much needed time in counterproductive activities such as squirming in their chairs, gazing out the window, etc. Knowing what maladaptive behaviors to expect from these children is the first step. Once it is known what behaviors to expect, the clinician can begin planning strategies for dealing with any difficulties that may arise because of them. If it is true that children who are language disordered display significantly more maladaptive behaviors, a routine part of any assessment should include assessing the child's behavior.

Finally, although this study is not designed to determine a causal relationship between maladaptive behaviors and language delays or disorders, it is hoped that it will provide some basis from which to formulate some hypotheses. For example, if children under 4 years of age do not display significantly more maladaptive behaviors than their

"normal" peers, it may be that the maladaptive behaviors develop as a result of the language delay. If maladaptive behaviors are a result of language delay, early language intervention might help to alleviate some of the problems.

In recent years there has been a growing trend toward early intervention for the language delayed child. Until recently, it was a commonly-held belief that young language delayed children, ages 18 to 36 months, are simply "late-bloomers" who will in time catch up with their peers. Parents were admonished not to worry and told to delay seeking intervention. Today the current trend is toward early intervention. It is felt that delaying intervention will simply allow language delayed children to lag further behind their peers. If language disorders are identified early, before maladaptive behaviors emerge, intervention could focus on remediating language without the added difficulty of dealing with a child who is behavior disordered. If on the other hand, maladaptive behaviors do accompany early language delay, intervention may be even more important to curtail problem behaviors as well as remediate the language delay.

STATEMENT OF PURPOSE

The purpose of this study was to determine if language delayed children between the ages of 18 and 41 months, or children who were "late talkers" display significantly more

maladaptive behaviors than their non-language delayed peers.

The questions this study sought to answer were:

1. Is there a significant difference in the severity and frequency of maladaptive behaviors seen in language delayed children, children who were "late talkers" or children with normal language?"

2. Is there a difference among the subject groups in terms of which behaviors parents are the most concerned with?

The hypothesis of this study is: The maladaptive behaviors of children 18 to 41 months old who have language delays, or who were "late talkers" will be significantly greater in terms of severity and frequency than the maladaptive behaviors of children of the same age who have normal language.

The null hypothesis is: The maladaptive behaviors of children 18 to 41 months old who have language delays, or who were "late talkers" will not differ significantly in terms of severity and frequency from the maladaptive behaviors of children of the same age who have normal language.

DEFINITION OF TERMS

The following operational definitions were used for this study:

Conduct - Problems in the area of conduct indicate a child whose behaviors violate the rights of others. Examples of problems with conduct include disturbing other children, defiance, and aggression.

Health - For the purpose of this study, the category of health is simply defined as problems with eating and sleeping.

Hyperactivity/Attention Deficit Disorder (Hyperactivity/ADD) - This category includes the behaviors overactivity, difficulty concentrating and short attention span. The child who displays these characteristics might have more trouble staying on task than other children of the same age. He may become easily frustrated.

Maladaptive Behaviors - Maladaptive behaviors are any behaviors which are considered negative by parents and others who come into contact with the child. This includes, but is not limited to, the behaviors in this section.

Mood - Problems with mood encompass such areas as anxiety and excessive sadness and includes more worries and fears than normal. The child may seem preoccupied and uninterested in playing.

Relationships - Problems in the area of relationships include shyness, avoidance of others, and poor eye contact.

The child with this difficulty may have problems forming relationships with his/her siblings and might display problems with attention seeking.

Stereotypical Behaviors - Stereotypical behaviors are movements or actions that are deliberate, repetitive and serve no useful purpose (American Psychiatric Association, 1980). In this study stereotypical behaviors include head banging, peculiar preoccupations and rocking.

CHAPTER II

REVIEW OF THE LITERATURE

There has been a great deal of interest in how children with speech and language disorders develop in terms of their social skills and emotions. It seems likely that if something as integral to being human as language is disrupted, this would have a negative impact on a person in a number of ways. In 1937, Orton stated that communicative disorders of all types would interfere with a person's social and emotional development. This statement has been upheld by a number of different research studies conducted over the years.

BEHAVIOR PROBLEMS OF CHILDREN WITH SPEECH AND/OR LANGUAGE IMPAIRMENTS

A number of studies conducted over the years have indicated that children with speech and language disorders are more likely than children without such disorders to display maladaptive behaviors. Cantwell and Baker (1977) conducted an extensive review of the literature and tentatively concluded that there is a higher incidence of psychiatric disorders in children who have speech and language disorders. They also concluded that the types of psychi-

atric disorders present in this population are not significantly different from those seen in the normal population. These findings were not considered conclusive, however, due to methodological problems such as differences in diagnosing and defining psychiatric disorders, sampling bias and inadequate sampling size, and failure to consider associative factors such as mental retardation and brain damage.

In an attempt to eliminate some of the methodological problems of the other studies, Cantwell et al. (1979) began a study which included 293 children, ages 1 year, 11 months to 15 years, 11 months from a community speech clinic. Each child was evaluated by a psychiatrist and an interview was conducted with each child's parents and teacher using modified versions of the Conners Parent and Teacher Questionnaire (Conners, 1973) and the Rutter Parent and Teacher Questionnaire (Rutter et al., 1970). A diagnosis of a speech and/or language disorder was made by a speech-language pathologist.

Cantwell et al. reported their findings for the first 100 subjects in 1979. The results of the psychiatric evaluation indicated that 53 out of the 100 subjects were diagnosed as having at least one psychiatric disorder according to the criteria of the Diagnostic and Statistical Manual of Mental Disorders (DSM-III) (The American Psychiatric Association, 1980). Attention deficit disorder was the most common, occurring in 19 children. This disorder includes a

developmentally inappropriate short attention span and poor concentration. Oppositional disorder, defined as an opposition to all authority, constant argumentativeness, and an unwillingness to respond to persuasion was the second most common psychiatric disorder. It occurred in 13 of the children. Twelve of the subjects had anxiety disorders; all were shy, 2 were overanxious, and 3 had separation anxiety disorders. Four of the subjects displayed conduct disorders including antisocial behavior which violates the rights of others. One had chronic depression, and 1 had a stereotyped movement disorder. They concluded that children with speech and/or language disorders were "at risk" for psychiatric illness, but a causal relationship was not determined.

Mattison, Cantwell and Baker (1980) later reported the findings from administration of the Conners (1972) and Rutter et al. (1970) parent and teacher questionnaires. They found that the items reported most frequently by parents of children with combined speech and language disorders were problems with attentional-motor items (e.g. easily frustrated, excitable, impulsive, restless, short attention span, easily distract-ed, and tantrums), conduct items (e.g., disobeys at home), health items (e.g., nightmares, problems getting to sleep and stomach aches), developmental and language items (e.g., hard to understand), mood items (e.g., angry or irritable), and relationship items (e.g., shy, fights with siblings, solitary and afraid of new

people). Teachers in the study reported much the same data as the parents, and it was determined that both agreed with the findings of the psychiatrist, particularly in the area of attentional-motor items and relationship items. (It must be noted that the study conducted by Mattison, Cantwell and Baker (1980) included 5 mentally retarded children, 4 hearing impaired children, 2 autistic children, and 1 child from a bilingual background).

In 1982(a), Baker and Cantwell published further findings on the same subjects. The parents of 40% of the language delayed children reported that their children's feelings were easily hurt, and 38% of the parents reported that their children were easily frustrated.

Baker and Cantwell further reported (1982b) that while only 29% of the children with a pure speech disorder (e.g., articulation disorder, voice disorder, fluency disorder) displayed some type of psychiatric illness (e.g., attention deficit disorders, avoidance disorders, oppositional disorders, separation anxiety disorders, adjustment disorders or affective disorders), 45% of the children with combined speech and language disorders, and a total of 95% of the children with a language disorder (e.g., expressive, receptive, or processing disorder no less than 6 months below age level) were diagnosed as psychiatrically ill.

Fifty-three percent of the children in the 1979 Cantwell et al. study were found to have "diagnosable

psychiatric disorders according to DSM-III criteria" (p. 499, Baker and Cantwell, 1987). In 1987 they published their findings on these children. The majority of differences and those which were most significant between the two groups were in the area of language development. They found that "The psychiatrically ill children showed significantly more disorders involving language, whereas the psychiatrically well children tended to have fewer disorders involving language and more disorders involving pure speech" (p. 507). Also, their expressive and receptive language deficits were more severe than those of the psychiatrically well children. The only developmental milestones which distinguished the ill children from the well children were the age at which the first word and the first sentence were spoken. The psychiatrically ill children were significantly delayed in both these areas. Cantwell and Baker (1987) found that speech and language are the factors "most significantly associated with psychiatric illness" (p. 508). They point out that this may mean that "speech and language factors may play a more direct role in the development of psychiatric disorders than has been previously hypothesized" (p. 508).

There have been other studies which support the finding that an association exists between speech and language disorders and behavioral disorders. Botelho (1986) cited the research of Beckey (1942) who observed children and reviewed teachers' reports to determine what factors were

related to delayed speech development (delayed speech development was not defined) and concluded that these children did not want attention, played alone, and cried easily. Botelho (1986) further reports that unlike Cantwell and Baker (1980), Beckey did not consider temper tantrums and thumb sucking to be frequently occurring behaviors in the delayed speech children.

STUDIES DIFFERENTIATING CHILDREN WITH SPEECH DISORDERS FROM CHILDREN WITH LANGUAGE DISORDERS

Studies have indicated that the type and severity of maladaptive behaviors observed in children are related to the type and severity of their communication deficit. Baker, Cantwell and Mattison (1980) compared the behavior of 46 children with "pure" speech disorders to that of 53 children with combined speech and language disorders and found that significant differences did exist. Children with combined speech and language disorders were rated significantly worse in terms of hyperactivity syndrome (e.g., excessive motor activity, attentional deficit, etc.), conduct disorders, establishing relationships and develop developmental phenomena (e.g., wetting bed/pants, crying easily, etc.).

Cantwell and Baker (1982a) concluded from their research that children with language disorders were much more likely to be psychiatrically ill than the children with pure speech disorders. Eighty-seven percent of the psychi-

atrically ill children had language disorders while only 36% of the psychiatrically well children had language disorders.

A study by Botelho (1986) supports these findings. Included in her study were 19 subjects between the ages of 5 and 11 years with speech and/or language impairments matched for age and sex with 19 subjects who had normal speech and language as determined by school records and a speech-language pathologist. Each group contained 14 boys and 5 girls. Behavior was evaluated using the Child Behavior Checklists (Achenbach & Edelbrock, 1983). Both parent and teacher questionnaires were used.

In comparing the speech and/or language impaired subjects to the normal subjects, Botelho (1986) found no significant differences. Upon dividing the boys into three groups, however, speech and language impaired, language impaired and speech impaired, she found significantly more maladaptive behaviors in both the speech and language impaired, and language impaired boys as compared to the speech impaired and normal boys. As with the boys, the girls displayed no significant differences between the speech and/or language impaired girls and the normal girls. They were not further subdivided into three groups because of the small number of female subjects.

The types of problems teachers identified more frequently in the speech and language, and language impaired boys included significantly lower school performance, inat-

tentiveness, aggressiveness, and more difficulty learning. These same differences were not reported by the parents, but Botelho (1986) speculated that the problems seen may be due to the more taxing language demands placed on children in school.

MALADAPTIVE BEHAVIORS OF CHILDREN WITH IMPAIRED LANGUAGE ONLY

Researchers have long suspected that language delays or disorders are related to behavioral and emotional problems. In 1959, Ingram remarked upon the large number of behaviorally disturbed children being seen by speech-language pathologists. Weiner (1968) also reported a high number of behaviorally disturbed children being seen by speech-language pathologists. The research which has been done to date supports these findings.

Stevenson and Richman (1978) conducted an epidemiological study of the language and behavior of 3-year old children. They took a random sample of 705 children. Using a test of expressive and receptive language, they found that 24 (3.1%) of the children had a language delay. They used a Behavior Screening Questionnaire to identify behavior problems and found that 101 (14.3%) of the children in the sample displayed behavior problems. Stevenson and Richman (1978) found that of the children with language delays, 59.1% had behavior problems. This is highly significant

when one considers that only 14.3% of the sample population displayed any behavioral problems. Also significant was the fact that 12.9% of the children identified as having behavior problems also were language delayed compared to 3.1% of the sample population who were language delayed.

In comparing the types of behavior problems seen in language delayed and non-language delayed children, Stevenson and Richman (1978) found few differences except that the behavior problems of the language delayed children were more severe. The types of problems they observed were most frequently in social relationships (e.g., dependency, relationships with siblings, relationships with peers). Also reported were problems with parental control, unhappy moods, poor appetite, overactivity, and difficulty concentrating.

LANGUAGE DELAYS IN BEHAVIORALLY DISTURBED CHILDREN

Other researchers have studied the language delays of children who display maladaptive behaviors. Some studies include children in child guidance centers (Chess & Rosenberg, 1974; Wylie, Franchak, & McWilliams, 1965), and others have studied behaviorally, emotionally disturbed children attending public schools (Camarata, Hughes, & Ruhl, 1988).

Wylie, Franchak, and McWilliams (1965) found that 15% of a total of 292 children in a community child guidance

clinic displayed what they referred to as "defective speech," including 27% who had articulation problems, 24% with delayed speech, 20% who stuttered, and 29% who had combined disorders. Five maladaptive behaviors were seen in the children with "defective speech" significantly more often than in the children with normal speech: Soiling, thumb-sucking, wetting, hyperactivity and involuntary movements. Only two of these behaviors, soiling and thumbsucking, were found to occur at a statistically significant level. It was noted by Wylie et al. that the fact that the children with impaired speech were significantly younger may have influenced their findings. The children with impaired speech ranged in age from 2 to 16 years with a mean age of 8 and a mode of 6. The children with normal speech included children aged 3 to 17 years, with a mean age of 10 and a mode of 9.

A study conducted by Chess and Rosenberg (1974) also looked at children in a child guidance center. They found that parents of 64% of the speech and language disordered children reported behavioral problems, including tantrums, disruptive behavior, difficulties in peer relationships, discipline problems, enuresis, separation anxiety, withdrawn behavior, hypochondriasm, and extreme sullenness. Hyperactivity was reported as a problem in 19% of the speech and language disordered subjects.

An interesting outcome in both the Wylie et al. (1965) study and the Chess and Rosenberg (1974) study is the incidence of children with articulation disorders versus children with language disorders. Normally, articulation comprises around 54% of all speech and language disorders (Spilka & Steer, 1951; Van Riper, 1954). Only 27% of the children in the child guidance centers studied by the researchers had articulation disorders. This may be an indicator that children with articulation disorders are less likely than children with language disorders to develop maladaptive behaviors.

Camarata, Hughes, and Ruhl (1988) conducted a study to determine if mild to moderately behavior disordered children have more significant language delays than children whose behavior is normal. Included in the study were 38 children identified as mild to moderately behavior disordered who were "enrolled at least part-time within special education classrooms in regular elementary schools" (p. 193). The subjects ranged in age from 8 years, 9 months to 12 years, 11 months. The Test of Language Development--Intermediate (TOLD--I) (Hammill & Newcomer, 1982) was used to assess the subjects' language development. It was administered with the following results: 27 (71%) of the subjects' standard scores fell two or more standard deviations below the mean of the normative data reported for the test on one or more subtests. Of the 11 remaining subjects, all but 1 scored

more than one standard deviation below the mean on one or more subtests. However, the intelligence quotients of the subjects in this study ranged from 67 to 126. Since the normative data on the TOLD--I is based on children with intelligence quotients in the normal range, Camarata and Hughes also reported separately for those subjects in their study who had normal IQ's. Of the remaining 21 subjects with normal IQ's, 20 received standard scores one or more standard deviations below the mean of the normative sample on one or more subtests.

SUMMARY

In surveying the literature, it is apparent that children with speech and/or language disorders are "at risk" for developing maladaptive behaviors. Several studies have shown a higher prevalence of behavior disorders in these populations (Cantwell & Baker, 1977). It has also been found that children with either combined speech and language disorders or language disorders display a higher incidence of maladaptive behaviors than do children who have only speech disorders. Cantwell and Baker (1982b) found that only 29% of children with pure speech disorders displayed maladaptive behaviors while 45% of the children with speech and language disorders and 95% of the children who had only a language disorder displayed some type of psychiatric illness.

However, while much is known about problem behaviors of language delayed or disordered children aged 3 and up, no research has focused strictly on children aged 3 and under. Cantwell et al. (1979) included subjects in their study who were as young as 1 year, 11 months old, but the study also included subjects as old as 15 years, 11 months. Wylie et al. (1965) studied children as young as 2 years old, but the mean age of the subjects was 8, and the mode 6.

The only study to date which has specifically focused on preschool children is that of Stevenson and Richman (1978) who did an epidemiological study of 3 year old children. They found that much like older children, 3 year old children who have language delays are significantly more likely than non-language delayed children to display maladaptive behaviors. However, it cannot be assumed that even younger children will follow the same pattern.

The importance of a study focusing on children in the earliest stages of language development must not be overlooked. It may be of real benefit in helping to solve the problem of the relationship between language delays or disorders and maladaptive behaviors. If nothing else, it will aid speech-language pathologists in deciding what areas to assess in the very young child.

CHAPTER III

METHODS

SUBJECT SELECTION

Three groups of subjects were used for this study: Language delayed, "late talkers," and children with normal language. The subjects were recruited for the study using media advertisements and referrals from local pediatric clinics. The language delayed (LD) group consisted of 34 children between the ages of 19 and 41 months. The determination of a language delay was made using the criteria outlined by Rescorla (1984). Children were considered language delayed if between the ages of 18 and 23 months they used less than 10 words, or if between 24 and 30 months of age their expressive vocabulary was less than 50 words and/or they used no two-word combinations. The second group, the "late talkers," consisted of 12 subjects between the ages of 19 and 41 months. The "late talkers" were those children initially classified as language delayed because their expressive vocabularies were less than 10 words at 18 to 23 months. However, these children's vocabularies "caught up," exceeding 50 words by 24 months of age. The third group, the normal language subjects, consisted of 33 children matched to the other two groups by age, sex ratio, and

socioeconomic status (SES). They were placed in the normal language group because their language exceeded the criteria outlined for language delay.

SES was determined using a four-factor scale developed by Myers and Bean (1968). SES scores on this scale range from 1 to 5 with one being the highest.

The subject data is reported on two separate tables as two behavioral questionnaires, (a) the Childhood Personality Scale (CPS) (Cohen, 1975), and (b) the Behavior Checklist for Toddlers (BCT) (Rescorla, 1984) were used in this study. The CPS was administered, on the average, 5 to 7 months later than the BCT. Therefore the mean age of the subjects is slightly higher on the CPS than it is on the BCT (See Tables I and II). Also, the number of subjects for the two behavioral questionnaires vary slightly as both questionnaires were not filled out by all parents.

TABLE I

DEMOGRAPHIC DATA FOR THE SUBJECTS ON THE CHILDHOOD PERSONALITY SCALE: MEANS OR PERCENTAGE.

Subject Group	N	Age in Months	Sex	SES
Language Delayed	33	32	73% Males 27% Females	2.97
Late Talkers	11	29	73% Males 27% Females	3
Normal	29	32	66% Males 34% Females	2.48

TABLE II

DEMOGRAPHIC DATA FOR THE SUBJECTS ON THE BEHAVIOR
CHECKLIST FOR TODDLERS: MEANS OR PERCENTAGE.

Subject Group	N	Age in Months	Sex	SES
Language Delayed	34	25	74% Males 26% Females	2.94
Late Talkers	12	25	75% Males 25% Females	3
Normal	28	26	67% Males 33% Females	2.47

A determination of the child's level of expressive vocabulary was made by having the parents fill out the Language Development Survey (LDS) (Rescorla, in press) (Appendix A). The LDS lists 350 common early words and parents are asked to circle the words their child says. The LDS has been shown by Rescorla to have excellent reliability, validity, sensitivity (i.e., the ability to correctly identify delayed children), and specificity (i.e., the ability to identify normal children as normal).

In order to be eligible to participate in the study, the children whose parents filled out the LDS had to meet a number of requirements. First, they had to have normal hearing as measured by passing a hearing screening at 25 dB in a sound field. Second, they had to have normal intelligence as determined by a score of 80 or above on the Mental Development Index of The Bayley Scales of Infant Development

(Bayley, 1969). Finally, subjects included in the study could show no signs of the disorders of autism, craniofacial or neuromotor dysfunction, as determined by the observation of a speech-language pathologist. All children who participated in this study passed the above screening measures.

INSTRUMENTS

The behavior of the subjects was assessed using two questionnaires filled out by the parents: (a) the CPS and, (b) the BCT. Each questionnaire requires the parent to read a word or sentence and assign a number designating the degree to which it describes their child.

The decision to use both questionnaires in this study was based upon several factors. First, both questionnaires were available at the time the data were collected. Second, the two questionnaires provide somewhat different information. The CPS requires the parents to judge how frequently a behavior occurs in their child. It also requires them to provide responses to very specific behavioral items such as "Turns his head away or looks down in an uncomfortable way when people pay attention to him." It was designed to assess psychiatric disorders in children from 18 months to 12 years of age. This makes it a useful tool if at later date a follow-up study is conducted using these subjects.

The BCT was designed specifically to assess the behavior of toddlers and requires the parent to make a more

general appraisal of their child's behavior. Rather than requiring parents to judge the frequency of certain behaviors their child displays, they are required to make a judgement of whether a behavior such as "shyness," or "moods" is "no problem," "some problem," or a "major problem". It may be that the items on the BCT are somewhat more relevant for toddlers. Also, it includes information on two areas of behavior that are not covered on the CPS, namely health and stereotypical behaviors. These are items that were covered in other studies and it was felt by this researcher that this information was important.

The CPS (Appendix B) has three separate sections, two of which were used for this study. The first section allows the individual to assign a score from 0 to 6, on how often each behavior is observed in the child (i.e., 0 = Never, 1 = Almost Never, 2 = Seldom, 3 = Half the Time, 4 = Frequently, 5 = Almost Always, 6 = Always). Eighteen items out of a total of 20 were used from the first section. The 2 items which were not used assessed language rather than behavior. Most statements in the first section refer to behaviors which are negative or maladaptive, but some refer to behaviors which are positive or adaptive. In order to use both the negative and positive statements, the scoring was reversed for the positive behaviors (i.e., a score of 0 became a score of 6; a score of 1, 5; a score of 2, 4; a score of 3 remained 3; a score of 4 became 2; a score of 5,

1; and a score of 6, 0.

The second section of the CPS has a statement in which the person is asked to assign a number indicating how closely that statement describes the child (e.g., 0 = Not At All, 1 = Just a Little, 2 = Pretty Much, 3 = Very Much). All 12 items from this section were used in the study.

The third section of the CPS was not used for the purpose of this study as it asked parents to make more of a general rating of their child's skills and did not cover specific behaviors.

The BCT (Appendix B) has only one section with a number scale from 0 to 2 (i.e., 0 = No Problem, 1 = Some Problem, 2 = Major Problem). The BCT has a total of 26 items. Only 25 of the items were used as one, "stuttering/stammering" assesses speech rather than behavior.

In order to facilitate comparisons across the two scales, as well as comparisons within the research literature, the items on the instruments were grouped into categories; on the CPS these categories include hyperactivity/ADD, immature relationships, conduct, and mood; on the BCT these categories include hyperactivity/ADD, immature relationships, conduct, mood, health and stereotypical behaviors.

On the CPS, Section 1, hyperactivity/ADD included the following items:

1. 'Loses interest in what he has started doing.
Goes from one thing to another.'
2. 'Jumps, runs and is on the move. Can't seem to be still for long.'
3. 'Persists in trying to do something, even if he has some small problems along the way.'*
4. 'Quickly shows his anger and frustration if he can't get something done that he's working on.'
5. 'Gives long attention to objects, toys or books that interest him.'*
6. 'Can get away from you quick as a flash when he wants to.'
7. 'Can pay attention to a long time to something.'*
8. 'Active, impossible to keep up with him.'

Hyperactivity on Section 2 of the CPS included:

9. 'Restless (overactive).'
10. 'Excitable, impulsive,'
11. 'Fails to finish things he starts,'
12. 'Fidgeting,'
13. 'Inattentive, distractible'
14. 'Demands must be met immediately; easily frustrated.'

A Hyperactivity score for the CPS was calculated by combining the scores for Sections 1 and 2.

On the BCT hyperactivity/ADD included:

1. 'Overactivity'
2. 'Concentration'
3. 'Is impulsive.'

The child's ability to form relationships and his social skills were also assessed. On Section 1 of the CPS this category included:

1. 'Turns his head away or looks down in an uncomfortable way when people pay attention to him.'
2. 'Shies away from getting attention.'
3. 'Tends to be resistant and unfriendly.'
4. 'Would rather be left alone if you try to play with him or talk to him.'
5. 'Smiles to a friendly person'*

Section 2 of the CPS did not include any relationship items.

On the BCT problems with relationships included:

1. 'Attention seeking'
2. 'Shyness'
3. 'Relationships with brothers/ sisters.'
4. 'Overly dependent'
5. 'Poor eye contact.'

The third category of behavior assessed was conduct.

On the CPS this included only two items in Section 2:

1. 'Disturbs other children.'
2. 'Temper outbursts.'

On the BCT conduct included these characteristics:

1. 'Difficult to manage.'
2. 'Aggression.'
3. 'Defiant.'

Mood was assessed using both sections of the CPS. On Section 1 problems with mood included:

1. 'Lies down, rests his head, or falls asleep instead of playing.'
2. 'Sits without doing anything unless another person tries hard to get him interested.'
3. 'Seems to have little zest for normal activities. Acts tired.'
4. 'Talks and acts happily and with excitement about things that interest him.'*

On Section 2 of the CPS mood included:

1. 'Cries'
2. 'Mood changes quickly.'
3. 'Temper outbursts.'
4. 'Anxious.'
5. 'Overly sad.'

On the BCT mood included:

1. 'Moods.'
2. 'Worries.'
3. 'Fears.'
4. 'Withdrawn, "spacey."'
5. 'Cries or laughs too easily.'

Health was assessed only on the BCT as the CPS has no items which fit in this category. It included only problems with:

1. 'Eating'
2. 'Sleeping'

Stereotypical behaviors were also assessed only on the BCT. This category included:

1. 'Echoes speech.'
2. 'Repetitive habits (flapping, twirling).'
3. 'Tics.'
4. 'Peculiar preoccupations.'
5. 'Rocks back and forth.'
6. 'Bangs head.'

*Scoring will be reversed

PROCEDURES

The behavioral questionnaires were filled out by either one or both parents together. The parents were given the questionnaire with the instructions to fill it out as they felt it best described their child. The CPS was given to

parents to fill out at home and they returned it by mail. The BCT was filled out by parents while they waited for language testing to be completed on their child.

DATA ANALYSIS

In order to compare the language delayed subjects, the "late talkers" and the normal language group, each child was assigned a score for each of the behavioral categories based upon the parent's rating. The scores were tallied using the coding sheets in Appendix C. The subject's scores are reported in Appendix D. The total score possible varied among the different behavioral categories. On the CPS the total possible score for hyperactivity/ADD was 66; the total possible for relationships was 30; the total possible for mood was 39; and the total possible for conduct was 6. On the BCT the total possible score for hyperactivity/ADD was 6; the total possible for relationships was 10; the total possible for mood was 10; the total possible for conduct was 8; the total possible for stereotypical behaviors was 12; and the total possible for health was 4.

After determining the mean score for each area of behavior for each of the three subject groups, the means were then compared to determine if a significant difference existed among the LD, "late talkers," and normal language. The two behavioral questionnaires, the CPS and the BCT were analyzed separately.

STATISTICAL ANALYSIS

The first question posed by this study was "Is there a significant difference in the severity and frequency of maladaptive behaviors displayed among the three subject groups?" To answer this question, t-tests were conducted to determine if there were any differences between either the normal language subjects and the LD subjects; the normal language subjects and the "late talkers"; or the LD subjects and the "late talkers." The t-tests were conducted at the .01 level of significance. A .01 level of significance was chosen as 30 t-tests were conducted and it was felt that using a lower level of significance would result in too high a percentage of Type I errors. The second question this study sought to answer was "Is there any difference among the three subject groups in terms of what types of behaviors parents are the most concerned about?" This question was answered by determining which areas of behavior parents assigned the highest percentage of total points possible to and to which area of behavior they assigned the lowest percentage of points possible. The areas of behavior were then ranked in order from highest to lowest percentage of points assigned.

RELIABILITY AND VALIDITY

Reliability

As a check on the researcher's scoring, 10% of the tests were rescored by a person who was trained as to how the tests were scored. The average percentage of agreement was 98%.

Validity

To check the validity of the categorization of the items on the two questionnaires into the factors of hyperactivity/ADD, relationships, mood, conduct, health and stereotypical behaviors, two other people categorized the items. The judges had no special background, but were trained as to what the behavioral terms in the study meant. The judges agreement with this author was, on the average, 90% on the BCT and 85% on the CPS.

CHAPTER IV

RESULTS AND DISCUSSION

RESULTS

The purpose of this study was to determine if differences exist in the severity, frequency, and type of maladaptive behaviors seen among language delayed children, children who were "late talkers," and normal language children aged 18 to 41 months as reported by parents. Two parent questionnaires were used, The Childhood Personality Scale (CPS) (Cohen, 1975) and The Behavior Checklist for Toddlers (BCT) (Rescorla, 1984).

The first question addressed was whether or not significant differences exist in the severity and frequency of maladaptive behaviors reported for LD children, children who were "late talkers," and normal language children. To answer this question, the means were compared for each behavioral category assessed on the two questionnaires for each of the three subject groups using t-tests. The differences among the subject groups are reported in Tables III, IV, and V for the CPS and in Tables VI, VII, and VIII for the BCT. Differences between the subject groups were considered significant at the .01 level.

On the CPS differences were found between the LD subjects and the normal subjects for both hyperactivity/ADD and conduct with the LD subjects displaying significantly more maladaptive behaviors in both these areas (See Table III).

TABLE III

t-TESTS INDICATING DIFFERENCES BETWEEN THE NORMAL (N)
AND LANGUAGE DELAYED (LD) SUBJECTS ON THE
CHILDHOOD PERSONALITY SCALE.

<u>Behavior</u>	<u>Mean</u> (SD)	<u>df</u>	<u>t</u>	<u>p</u> <
Hyperactivity/ ADD	N - 23.40 (9.29)	61	-2.911	<.010*
	LD - 30.26 (9.23)			
Relationships	N - 6.14 (4.71)	61	-1.900	.118
	LD - 8.70 (5.88)			
Mood	N - 5.07 (3.16)	55	-2.477	.034
	LD - 7.74 (5.30)			
Conduct	N - 1.36 (.95)	59	-2.748	<.010*
	LD - 2.18 (1.38)			

* Significant

The CPS showed no significant differences between the normal subjects and the "late talkers" or between the delayed subjects and "late talkers" (See Tables IV and V).

TABLE IV
t-TESTS INDICATING DIFFERENCES BETWEEN THE
NORMAL (N) AND LATE TALKING (LT) SUBJECTS
ON THE CHILDHOOD PERSONALITY SCALE.

<u>Behavior</u>	<u>Mean</u> (SD)	<u>df</u>	<u>t</u>	<u>p</u>
Hyperactivity/ADD	N - 23.40 (9.29)	24	-2.705	.024
	LT - 31.09 (7.50)			
Relationships	N - 6.14 (4.71)	21	-1.000	.999
	LT - 7.73 (4.40)			
Mood	N - 5.07 (3.16)	13	-2.638	.040
	LT - 9.50 (5.22)			
Conduct	N - 1.36 (.95)	15	-2.608	.038
	LT - 2.50 (1.32)			

TABLE V

t-TESTS INDICATING DIFFERENCES BETWEEN THE LANGUAGE
 DELAYED (LD) AND LATE TALKING (LT) SUBJECTS
 ON THE CHILDHOOD PERSONALITY SCALE.

<u>Behavior</u>	<u>Mean</u> (SD)	<u>df</u>	<u>t</u>	<u>p</u> <
Hyperactivity/ADD	LD - 30.26 (9.23)	23	-.300	.999
	LT - 31.09 (7.50)			
Relationships	LD - 8.70 (5.88)	25	.579	.999
	LT - 7.73 (5.40)			
Mood	LD - 7.74 (5.30)	17	-.964	.999
	LT - 9.50 (5.22)			
Conduct	LD - 2.18 (1.38)	19	-.683	.999
	LT - 2.50 (1.32)			

The BCT showed the LD subjects to have significantly higher scores than the normal subjects in the areas of hyperactivity/ADD and relationships. No significant differences were found between the normal and LD subjects in the areas of mood, conduct, health or stereotypical behaviors (See Table VI).

TABLE VI

t-TESTS INDICATING DIFFERENCES BETWEEN THE NORMAL (N)
AND LANGUAGE DELAYED (LD) SUBJECTS ON THE
BEHAVIOR CHECKLIST FOR TODDLERS.

<u>Behavior</u>		<u>Mean</u> (SD)	<u>df</u>	<u>t</u>	<u>p</u> <
Hyperactivity/ ADD	N - .30 (.65)		48	-3.151	<.010*
	LD - 1.18 (1.47)				
Relationships	N - .62 (.81)		51	-3.214	<.010*
	LD - 1.62 (1.60)				
Mood	N - .67 (.84)		56	-1.765	.158
	LD - 1.18 (1.42)				
Conduct	N - 1.17 (1.21)		62	-2.639	.020
	LD - 2.08 (2.38)				
Health	N - .63 (.75)		58	-.479	.999
	LD - .74 (.75)				
Stereotypical Behaviors	N - .33 (.80)		62	-1.007	.638
	LD - .53 (.75)				

* Significant

No significant differences were found between the normal subjects and "late talkers" or between the LD subjects and the "late talkers" on the BCT (Tables VII and VIII).

The second question addressed in this study was whether or not differences exist between the subject groups in terms of which behaviors parents are the most concerned with. In other words, for each subject group, to what behaviors did parents assign the highest scores, and to what areas of behavior did they assign lower scores? Since the categories of maladaptive behaviors do not contain equal numbers of points possible on either of the two questionnaires, a rank ordering was determined by dividing the total points possible on a particular category of behavior by the mean received by the subject group. On the CPS, no differences were found between the types of concerns displayed by the parents of each subject group. That is, parents of LD children, normal language children and "late talkers" all assigned the highest percentage of possible points to the category hyperactivity/ADD. The parents ranked concern about their child's conduct, second, immature relationships, third, and mood last. The results are displayed in Figure 1.

Some differences did occur in the parents concerns as outlined by the BCT. Parents of both the late talkers" and the LD subjects were most concerned with their child's

TABLE VII

t-TESTS INDICATING DIFFERENCES BETWEEN THE NORMAL (N)
AND LATE TALKING (LT) SUBJECTS ON THE
BEHAVIOR CHECKLIST FOR TODDLERS.

<u>Behavior</u>		<u>Mean</u>	<u>df</u>	<u>t</u>	<u>p</u> <
Hyperactivity/ ADD	N -	.30 (.65)	17	-2.158	.086
	LT -	.92 (.90)			
Relationships	N -	.62 (.81)	18	-1.658	.222
	LT -	1.17 (1.03)			
Mood	N -	.67 (.84)	26	.225	.999
	LT -	.58 (1.17)			
Conduct	N -	1.17 (1.21)	14	-1.502	.304
	LT -	2.08 (2.38)			
Health	N -	.63 (.93)	27	-.423	.999
	LT -	.75 (.75)			
Stereotypical Behaviors	N -	.33 (.80)	41	.903	.999
	LT -	.17 (.39)			

TABLE VIII

t-TESTS INDICATING DIFFERENCES BETWEEN THE LANGUAGE DELAYED
(LD) AND LATE TALKING (LT) SUBJECTS ON THE
BEHAVIOR CHECKLIST FOR TODDLERS.

<u>Behavior</u>	<u>Mean</u>	<u>df</u>	<u>t</u>	<u>p<</u>
Hyperactivity/ ADD	LD - 1.18 (1.47)	35	.718	.999
	LT - .92 (.90)			
Relationships	LD - 1.62 (1.60)	33	-1.115	.546
	LT - 1.17 (1.03)			
Mood	LD - 1.18 (1.42)	26	1.427	.324
	LT - .58 (1.17)			
Conduct	LD - 2.10 (1.62)	14	-.198	.999
	LT - 2.08 (2.38)			
Health	LD - .74 (.75)	21	-.058	.999
	LT - .75 (.39)			
Stereotypical Behaviors	LD - .53 (.75)	40	2.127	.074
	LT - .17 (.39)			

Figure 1. Order in which parents ranked their concerns about the behaviors displayed by their children on the CPS.

Key:

H = Hyperactivity, C = Conduct, R = Relationships, M = Mood

□ = LD Subjects, ■ = Late Talkers, ▲ = Normal Subjects

conduct while parents of the normal language children displayed the most concern for their child's health. However, parents of normal children did assign the second highest percentage of points possible to the area of conduct. The parents of the "late talkers" were also more concerned about their child's health than for any of the other behavioral items, ranking it second highest. For the LD subjects parents considered hyperactivity/ADD (number 2) to be more of a concern than health (number 3). Parents of the normal language subjects ranked mood as their third area of concern. Mood was ranked fifth by parents of both the LD subjects and the "late talkers." Parents of all three subject groups considered relationships to be their fourth area of concern. Hyperactivity/ADD was ranked as fifth by parents of the normal language subjects. Stereotypical behaviors were listed as the area of least concern by parents of all three subject groups. The results are displayed in Figure 2.

DISCUSSION

The two questionnaires, the CPS and the BCT both show the LD subjects display significantly higher scores in the area of hyperactivity/ADD than the normal subjects. This finding is in agreement with the 1978 study of British children conducted by Stevenson and Richman. They found that 3-year old language delayed children display significantly

Figure 2. Order in which parents ranked their concerns about the behaviors displayed by their children on the BCT.

Key:

H = Hyperactivity/ADD, R = Relationships, M = Mood,
 C = Conduct, He = Health, S = Stereotypical Behaviors
 □ = LD Subjects, ■ = Late Talkers, ▲ = Normal Subjects

higher scores in the areas of overactivity ($p < .01$) and poor concentration ($p < .01$). Hyperactivity and attention deficit disorder have also been found to occur more frequently in subjects diagnosed as childhood aphasics (Paul, Cohen, & Caparulo, 1983; Paul & Cohen, 1984), and in school-age children with less severe language delay (Wyllie, et al., 1965; Botelho, 1986).

The two questionnaires also were in agreement that no significant differences exist between the normal subjects and the "late talkers" nor between the LD subjects and the "late talkers." This finding indicates that the scores of the "late talkers" fall midway between the scores of the normal language and LD subjects in terms of their maladaptive behaviors.

There are two categories on which the two questionnaires differed; relationships and conduct for the normal and LD subjects. The CPS showed the LD subjects to have a significantly higher mean conduct score than the normal language subjects. Although the BCT showed the LD subjects to have more problems with conduct than the normal language subjects, the difference was not statistically significant.

Another area of difference is that the BCT showed the LD subjects to have a significantly higher mean for immature relationships than the normal language subjects. Again, while the CPS did not show a statistically significant difference between the normal and LD subject groups, the LD

group did receive a higher mean score. This researcher does not find it wholly unexpected that differences exist between the two questionnaires. In comparing the items on the CPS and the BCT, it can be seen that for several of the categories, the items on the two questionnaires are quite different, particularly in the areas of relationships and conduct. The conduct category on the CPS includes only two items, i.e., "disturbs other children" and "temper outbursts." The BCT has four items that comprise the category of conduct, i.e., "difficult to manage," "temper tantrums," "aggression" and "defiant."

This researcher cannot say with certainty which of the questionnaires more accurately reflects the actual behavior of the subjects. However, there are at least three possible reasons for the differences. First, on the CPS there are only two items categorized as conduct. It may be that if the CPS contained more items in the category of conduct the difference between the normal and delayed language subjects may not have been significant. Also, the items on the BCT are more relevant for toddlers than the items on the CPS. The BCT may be a more accurate reflection of the behavior of the children in the study. However, the possibility exists that the CPS is actually a more sensitive measure of the parent's true opinion. It may be that since the CPS simply requires the parent to state how frequently the behavior occurs, the parent is responding more accurately. A parent

might be reluctant to label an unacceptable behavior as a "major problem" as is the case if the parent assigns the full number of points (2) on the BCT.

The research literature seems to support the findings of the CPS, that is, that language delayed children do show significantly more problems with conduct than do children with normal language skills. This has been the case with older language delayed children. Stevenson and Richman (1978) found that 3-year old language delayed children are significantly more "difficult to control" than normal language children ($p < .001$). However, they found no differences between the two groups in the area of "tempers." Mattison et al. (1980) in their study of 99 children from a community speech and hearing clinic stated that the speech and language disordered children were rated severe by parents for such behaviors as tantrums, which supports the findings of the CPS in this study as "temper outbursts" was one of the two items labeled as conduct. Teachers rated this same group of children as having a problem with their attitude toward authority. Although the BCT did not find a statistically significant difference between the normal and LD subjects, it was filled out by parents only, not by teachers as was the case in the study conducted by Mattison et al. (1980). In comparing this study to that of Mattison et al. (1980) it must be remembered that the children in their study were in preschool or middle school.

The relationships category on the BCT showed the LD subjects to score significantly higher than the normal language subjects. The CPS did not show a statistically significant difference between the two groups in this area, although the language delayed subjects did, on the average, score higher. For both questionnaires, five items comprise the category immature relationships. However, the items on the two questionnaires are quite different. The CPS relationship items are very specific and seem to portray a child who is unfriendly and does not enjoy receiving attention (See Appendix B for a complete listing of the items). The items on the BCT tend to show a child who, while not unfriendly is shy, overdependent, and has some problems getting along with his/her brothers and sisters. It may be that while problems with relationships do occur more frequently for children with delayed language, the manner in which the problems are manifested are more in terms of overdependence rather than withdrawal and hostility.

Other research tends to support the finding of the BCT for immature relationships (i.e. delayed language subjects show significantly more problems with relationships than do their normal language peers). Stevenson and Richman (1978) found that language delayed children displayed significantly higher scores in the areas of problems with dependency ($p < .05$), relationships with siblings ($p < .01$), and relationships with peers ($p < .05$). An interesting fact is that

Stevenson and Richman (1978) found no difference between the two groups in the area of attention seeking. Attention seeking was one of the items on the BCT labeled as immature relationships. As a comparison of individual items was not done in this study, it cannot be stated with certainty if this one item differed significantly between the two groups. Baker et al. (1980) also found older language delayed children to show significantly more problems with relationships. Parents in that study named shy, fights with siblings, solitary and afraid of new people as being problem items for the combined speech and language impaired children.

No significant differences were found between either the "late talkers" and the normal language subjects or between the "late talkers" and the LD subjects. It may be that if more "late talkers" had been included in the study more significant differences would have emerged.

The second question addressed in this study was whether or not parents of the three subject groups differed in terms of what areas of behavior they were most concerned. It seems that parents of children in each group differ little on areas of most concern. No differences were found on the CPS and only minor differences were found on the BCT.

The only difference on the BCT between the "late talkers" and the LD subjects was that health and hyperactivity/ADD were reversed for these two groups. Health was ranked second and hyperactivity/ADD third by the parents of

the "late talkers." Parents of the LD subjects ranked their concern about their children's hyperactivity/ADD second and concern about health third. Somewhat different results occurred for the normal subjects with parents' scoring revealing health to be their area of most concern. They placed hyperactivity/ADD quite low on their list of concerns (fifth). This is not unexpected as one would expect parents of normal children to be more concerned about problems with eating and sleeping than anything else. Some difference in how parents ordered areas of concern may be due to the fact that health was not one of the categories assessed on the CPS. These findings were supported by the research of Baker and Cantwell (1982b) who found that parents of language disordered children complained more frequently about hyperactivity and developmental problems while parents of pure speech disordered children complained more frequently about somatic complaints.

It is interesting to note though, that parents of normal language subjects ranked hyperactivity/ADD fifth on the BCT when they ranked it first on the CPS. This difference may in part be due to the fact that two of the three items on the BCT have to do with the child's ability to concentrate and his/her attention span, while on the CPS there are more items which assess the child's level of activity. It may be that parents of normal children are more concerned with excess energy than they are with short attention span.

Hyperactivity/ADD also was ranked lower on the BCT than on the CPS by parents of both the "late talkers" and the LD subjects.

The finding from this research that little difference exists in the types of maladaptive behaviors displayed by language delayed and normal language children is supported by Cantwell and Baker (1977) and Stevenson and Richman (1978). It also has been found that the types of psychiatric disorders displayed by children with more severe handicaps (e.g. brain damage and intellectual retardation) do not differ markedly from the types of psychiatric disorders seen in the general population (Cantwell & Baker, 1977). In other words, the same types of maladaptive behaviors are seen in both normal and language delayed children, but language delayed children display more maladaptive behaviors than non-language delayed children.

CHAPTER V

SUMMARY AND IMPLICATIONS

SUMMARY

Human beings rely on language to share their thoughts and feelings, to express their needs and intentions. Without a shared language, man would be much the same as any other animal. It seems likely that if a children's language skills were impaired, it would have a negative impact on their social development. Several studies support the hypothesis that language delayed children are "at risk" for developing maladaptive behaviors (Cantwell & Baker, 1977). To date there have been several studies which have looked at the behaviors of speech and/or language delayed children. However, only one of these studies focused primarily on pre-school children. In 1978, Stevenson and Richman studied 3-year old children in the London area to determine if language delayed children showed more maladaptive behaviors than nonlanguage delayed children. They found that even language delayed children as young as 3 years old have significantly more maladaptive behaviors than their non-language delayed peers. However, it cannot be assumed from this study that even younger children will follow the same trend. Therefore, this study was designed to look at chil-

dren in the earliest stages of language development. The subjects ranged in age from 18 to 41 months.

The questions addressed by this study were:

1) Is there a significant difference in the severity and frequency of maladaptive behaviors seen in language delayed children, non-language delayed children, and children who were "late talkers" between the ages of 18 and 41 months as reported by parents? and; 2) Is there a difference among the subject groups in terms of which behaviors parents are the most and least concerned about. The means and standard deviations were computed for each category of behavior reported on the two parent questionnaires. Differences were determined by comparing the results between the three groups of subjects using t-tests and were considered significant at the .01 level.

The data was analyzed using t-tests because this researcher felt from looking at the raw data and from reviewing the literature that differences would exist among the three groups. The purpose of this study was to determine what those differences were. Differences were considered significant only at the .01 level because of the large number of t-tests conducted. It was felt that using a lower level of significance would result in too high a chance of Type I errors.

In looking at the results of this study, it can be seen that even very young language delayed children and

those who begin talking late are "at risk" for increased maladaptive behaviors. The language delayed children received significantly higher scores than the normal language children in the areas of hyperactivity (as reported both on the CPS and the BCT), conduct (CPS only) and relationships (BCT only).

IMPLICATIONS

Research Implications

There are several interesting findings from this study, but the need for further research still exists. A limitation of this study is that only 11 to 12 subjects were used in the groups of "late talkers." It would be useful to conduct a study with a larger sample of "late talkers." Perhaps if more "late talkers" had been included in this study, some significant differences would have been found between the "late talkers" and the other two subject groups.

Another area of further research would be to conduct a longitudinal study to determine whether or not changes in maladaptive behaviors are observed in the "late talkers" or the LD subjects over time and/or as their language improves. As there have been no studies conducted to date as to what types of maladaptive behaviors are displayed by children who begin talking late, it would be especially useful to know whether or not their behavior improves over time or whether it remains the same. If these behaviors were to improve as

the children's language improved it could then be hypothesized that the children's inability to express themselves clearly led to the maladaptive behaviors.

Since these maladaptive behaviors do manifest themselves so early, however, it seems likely that language is not a direct cause of the maladaptive behaviors but rather that both occur due to some other mutual cause, such as temperamental differences, inability to learn pro-social skills, inability to use models in the environment or interaction with parenting style. It may be that the temperaments of children with language disorders differ from those of children without language disorders. If differences exist in this area, it could affect how parents perceive their children, thereby influencing how they describe their behaviors. Also, a mismatch between the temperament of the child and the parent could influence the child's personality development. This factor was considered before beginning the study, but at the time, no measurements of temperament were available for children under 4 years of age.

Other researchers have hypothesized that both psychiatric disorders and speech and language disorders are due to some mutual cause. Cantwell and Baker (1977) hypothesized that both could be due to such common antecedents as intellectual retardation, deafness, or brain damage. These factors were taken into consideration in this study. All children who participated in the study had normal intelli-

gence and normal hearing, and none of the children in the study had brain damage. It may be that only an indirect link exists between speech and language disorders and behavior disorders (Rutter, 1972). It is only through further research that these questions will be answered.

A further limitation of this study was that neither the CPS nor the BCT have been evaluated for reliability or validity. The results of this study would perhaps be strengthened if that information were available.

Clinical Implications

This study has several important implications for the practicing speech-language pathologist. First, it provides support for the trend toward early intervention. Second, it provides some evidence that even children who begin talking late but then talk normally are still more likely to display more maladaptive behaviors than their normal language peers. Third, it points out the need for a multidisciplinary team approach to early childhood intervention. In looking at the findings from this research, it can be seen that even very young language delayed children are more likely than their normal peers to display inappropriate behaviors. In looking at the research to date, it can be hypothesized that since older language delayed children display significantly more maladaptive behaviors than their normal language peers that these behaviors are not going to just disappear. It is important that speech-language pathologists share with other

professionals the knowledge that even very young language delayed children are more likely to develop maladaptive behaviors. As the same types of behaviors are observed in older language delayed children (Cantwell & Baker, 1977) it does not seem that simply ignoring the problem makes it go away. This leads to the issue of whether or not speech-language pathologists should be concerned about the child who begins to talk late.

In the past, parents were often told not to worry if their children did not say their first word until two or even three years of age. However, this study gives reason for some concern. Although no significant differences were found between the normal language children and the "late talkers," there was a definite trend for the "late talkers" to display higher scores in almost all areas assessed. This indicates that while the "late talkers" do not have behavioral problems as severe as those of the LD children, they are still "at risk" and should not be considered "normal." If nothing else, it might be reassuring for parents of children who are "late talkers" to know that certain differences in behavior should not be totally unexpected for their child.

Finally, this study underscores the need for a multidisciplinary team approach in early language intervention. The speech-language pathologist must be prepared to see more maladaptive behaviors in language delayed children and must

have knowledge of how to deal with those behaviors. Botelho (1986), in discussing the findings for her research, made the point that speech-language pathologists and psychologists must work together to find the best possible treatment plan for language delayed children. She wrote:

A team approach may be most effective in working with the child, as the speech-language pathologist can provide information to the school psychologist/counselor to help them understand the language component of the behavior problem and they can provide the speech-language pathologist with appropriate techniques to deal with the behavioral problems to minimize interference with learning.
(p. 65)

In looking at the combination of behavioral problems and language disorders seen even in children less than 3-years old, the importance of a multidisciplinary approach cannot be overstressed. It is difficult to determine as of yet, whether or not language disorders lead to behavioral disorders or whether the relationship between the two is less direct. The best approach therefore seems to be one that would allow the speech-language pathologist and the psychologist to combine their unique areas of knowledge to provide the best possible intervention for these children.

SELECTED BIBLIOGRAPHY

- Achenbach, T.M., and Edelbrock, C. (1983). Manual for the Child Behavior Checklist and Revised Child Behavior Profile. U.S.A: Queen City Printers, Inc.
- American Psychiatric Association Task Force on Nomenclature and Statistics (1980). Diagnostic and Statistical Manual of Mental Disorders, (DSM--III) 3rd ed. Washington, D.C.: American Psychiatric Association.
- Baker, L., and Cantwell, D.P. (1987). Factors associated with the development of psychiatric illness in children with early speech/language problems. Journal of Autism and Developmental Disorders, 17, 499-510.
- Baker, L., and Cantwell, D.P. (1982a). Developmental, social and behavioral characteristics of speech and language disordered children. Child Psychiatry and Human Development, 12, 195-206.
- Baker, L., and Cantwell, D.P. (1982b). Psychiatric disorders in children with different types of communication disorders. Journal of Communication Disorders, 15, 113-136.
- Baker, L., Cantwell, D.P., and Mattison, R.E. (1980). Behavior problems in children with pure speech disorders and in children with combined speech and language disorders. Journal of Abnormal Child Psychology, 8, 245-256.
- Bayley, N. (1969). The Bayley Scales of Infant Development. Atlanta: The Psychological Corporation.
- Beckey, R.E. (1942). A study of certain factors related to retardation of speech. Journal of Speech Disorders, 7, 223-249.
- Botelho, J.S. (1986). A Comparison of Behavioral Problems Between Speech and /or Language Impaired Children and Normal Children. Thesis submitted at Portland State University.

- Camarata, S.M., Hughes, C.A., and Ruhl, K.A. (1988). Mild/moderate behaviorally disordered students: A population at risk for language disorders. Language, Speech and Hearing Services in the Schools, 19, 191-200.
- Cantwell, D.P., and Baker, L. (1980). Psychiatric and behavioral characteristics of children with communication disorders. Journal of Pediatric Psychology, 2, 161-178.
- Cantwell, D.P., and Baker, L. (1977). Psychiatric disorder in children with speech and language retardation: Archives of General Psychiatry, 34, 583-591.
- Cantwell, D.P., Baker, L., and Mattison, R.E. (1979). The prevalence of psychiatric disorder in children with speech and language disorder: An epidemiological study. American Academy of Child Psychiatry Journal, 18, 450-461.
- Chess, S., and Rosenberg, M. (1974). Clinical differentiation among children with initial language complaints. Journal of Autism and Childhood Schizophrenia, 4, 99-109.
- Cohen, D. (1975). The Childhood Personality Scale. Washington, D.C.: The National Institute of Mental Health.
- Conners, C.K. (1973). Rating scales for use in drug studies with children,. Psychopharmacology Bulletin, 24-59.
- Hammill, D., and Newcomer, P. (1982). Test of Language Development--Intermediate. Austin, TX: PRO-ED.
- Ingram, T.T.S. (1959). A description and classification of the common disorders of speech in children. Archives of Disorders in Children, 36, 444-467.
- Laughton, J., and Hasenstab, M.S. (1986). The Language Learning Process: Implications for Management of Disorders. Maryland: Aspen Publishers, Inc.
- Mattison, R., Cantwell, D.P., and Baker, L. (1980). Behavior problems in children with speech and language retardation. Child Psychiatry and Human Development, 10, 246-257.
- Myers, J. K., and Bean, L. L. (1968). A Decade Later: A Follow-Up of Social Class and Mental Illness. New York: Wiley & Sons.

- Orton, S.T. (1937). Reading, Writing, and Speech Problems in Children. New York: W.W. Norton Co., Inc.
- Paul, R., and Cohen, D.J. (1984). Outcomes of severe disorders of language acquisition. Journal of Autism and Developmental Disorders, 14, 405-421.
- Paul, R., Cohen, D.J., and Caparulo, B.K. (1983). A longitudinal study of patients with severe developmental disorders of language learning. Journal of the American Academy of Child Psychiatry, 22, 525-534.
- Rescorla, L. (In press). The language development survey: A screening tool for delayed language in toddlers.
- Rescorla, L. (1984). The Behavior Checklist for Toddlers: Language Delay at Two. Paper presented at the National Conference of the American Academy of Child Psychiatry. Boston, MA.
- Rutter, M. (1972). Psychiatric causes of language retardation, in M. Rutter (ed.) The Child with Delayed Speech. London: William Heinemann, Ltd.
- Rutter, M., Graham, P., and Yule, W. (1970). A Neuro-Psychiatric Study in Children. Suffolk, England: Lavenham Press, Ltd.
- Spilka, B., and Steer, M.D. (1951). Incidence of speech and hearing deficiencies in the general population and schools of the U.S. Lafayette, IN: Purdue Speech and Hearing Clinic Staff.
- Stevenson, J., and Richman, N. (1978). Behavior, language and development in three-year-old children. Journal of Autism and Childhood Schizophrenia.
- Van Riper, C. (1954). Speech Correction Principles and Methods. (3rd edition). Englewood Cliffs: Prentice-Hall, Inc.
- Weiner, P.S. (1969). The emotionally disturbed child in the speech clinic: Some considerations. Journal of Speech and Hearing Disorders, 33, 158-167.
- Wylie, H.L., Franchak, P., and McWilliams, B.J. (1965). Characteristics of children with speech disorders seen in a child guidance center. Perceptual Motor Skills, 20, 1101-1107.

APPENDIX A
LANGUAGE DEVELOPMENT SURVEY

The Language Development Survey

<u>FOOD</u>	<u>ANIMALS</u>	<u>ACTIONS</u>	<u>HOUSEHOLD</u>	<u>PERSONAL</u>	<u>CLOTHES</u>
apple	bear	bath	bathtub	brush	belt
banana	bee	breakfast	bed	comb	boots
bread	bird	bring	blanket	glasses	coat
butter	bug	catch	bottle	key	diaper
cake	bunny	clap	bowl	money	dress
candy	cat	clean	chair	paper	gloves
cereal	chicken	close	clock	pen	hat
cheese	cow	come	crib	pencil	jacket
coffee	dog	cough	cup	penny	mittens
cookie	duck	cut	door	pocketbook	pajamas
crackers	elephant	dance	floor	tissue	pants
drink	fish	dinner	fork	toothbrush	shirt
egg	frog	doodoo	glass	umbrella	shoes
food	horse	down	knife	watch	slippers
grapes	monkey	eat	light		sneakers
gum	pig	feed	mirror	<u>PEOPLE</u>	socks
hamburger	puppy	finish	pillow	aunt	sweater
hotdog	snake	fix	plate	baby	
icecream	tiger	get	potty	boy	<u>OUTDOORS</u>
juice	turkey	give	radio	daddy	flower
meat	turtle	go	room	doctor	house
milk		have	sink	girl	moon
orange	<u>BODY</u>	help	soap	grandma	rain
pizza	<u>PARTS</u>	hit	sofa	grandpa	sidewalk
pretzel	arm	hug	spoon	lady	snow
soda	belly	jump	stairs	man	star
soup	bottom	kick	table	mommy	street
spaghetti	chin	kiss	telephone	own name	sun
tea	ear	knock	towel	pet name	tree
toast	elbow	look	trash	uncle	
water	eye	love	tv	Ernie etc	<u>PLACES</u>
	face	lunch	window		church
<u>TOYS</u>	finger	make		<u>VEHICLES</u>	home
ball	foot	nap		bike	hospital
balloon	hair	outside		boat	library
blocks	hand	pattycake		bus	McDonalds
book	knee	peekaboo		car	park
bubble	leg	peepee		motorbike	school
crayons	mouth	push		plane	store
doll	neck	read		stroller	zoo
present	nose	ride	<u>ACTIONS</u>	train	
slide	teeth	run	(Cont.)	trolley	
swing	toe	see	take	truck	
teddybear	tummy	show	throw		
		sing	tickle		
		sit	walk		
		sleep	want		
		stop	wash		

MODIFIERS

allgone
all right
bad
big
black
blue
broken
clean
cold
dark
dirty
down
good
happy
heavy
hot
hungry
little
mine
more
open
pretty
red
shut
stinky
that
this
tired
up
wet
white
yellow
yucky

OTHER

A, B, C etc
away
booboo
byebye
curse words
here
hi, hello
in
me
meow
my
myself
nightnight
no
off
on
out
please
Sesame Street
scuse me
shut up
thank you
there
under
welcome
what
where
why
woofwoof
yes
you
yumyum
1, 2, 3 etc

Please list any other words
your child uses here:

Does your child combine two
or more words in phrases?
(e.g. more cookie, car
bye bye, etc)
yes_____ no_____

Please list below THREE of
your child's longest and best
sentences or phrases:

APPENDIX B

BEHAVIORAL ASSESSMENT QUESTIONNAIRES

DEPARTMENT OF HEALTH, EDUCATION, AND WELFARE
NATIONAL INSTITUTE OF MENTAL HEALTH

CHILDHOOD PERSONALITY SCALE (CPS)--R
(Factor Scale)

____ Father

____ Mother

Family Name:

Name of Child:

Age:

Date:

INSTRUCTIONS: The purpose of this questionnaire is to get a picture of each child's personality as he or she typically has been for the last two months. Some of the sentences may describe the child very well. Other sentences will not be at all like this child. There are seven columns after each sentence. For each sentence, check the column that is most true of this child's personality and the way he acts.

This is an example of the type of question and how to answer it: Happy to sing when there are guests in the house.

Never	Almost Never	Seldom	Half The Time	Fre- quently	Almost Always	Always
0	1	2	3	4	5	6

If this child always sings when there are guests, you should check column "6". If he never sings for company, check column "0". If he sings sometimes but not always, check the box between "0" and "6" that best describes him.

	Never	Almost Never	Seldom	Half The Time	Fre- quently	Almost Always	Always
	0	1	2	3	4	5	6
H Loses interest in what he has started doing. Goes from one thing to another							
H Jumps, runs, and is on the move. Can't seem to be still for long							
R Turns his head away or looks down in an uncomfortable way when people pay attention to him.							
M Lies down, rests his head, or falls asleep instead of playing							
M Talks with delight. Gets pleasure out of almost everything							
H Persists in trying to do something, even if he has some small problems along the way							
H Quickly shows his anger and frustration if he can't get something done that he's working on							
R Shies away from getting attention. Moves away from people							
M Sits without doing anything unless another person tries hard to get him interested							
Will talk or babble to you about his toys, clothes, and what he is doing							
H Gives long attention to objects, toys, or books that interest him							
H Can get away from you "quick as a flash" when he wants to							

	Never	Almost Never	Seldom	Half The Time	Fre- quently	Almost Always	Always
	0	1	2	3	4	5	6
R Smiles to a friendly person							
M seems to have little zest for normal activities. Acts tired							
M Talks and acts happily and with excitement about things that interest him							
H Can pay attention for a long time to something							
H Active, impossible to keep up with him							
R Tends to be resistant and unfriendly							
R Would rather be left alone if you try to play with him or talk to him							
Is a talkative child who expresses himself in language or near language							

Listed below are items about children's behavior or problems they sometimes have. How much do you think your child has been bothered by this problem at this time?

	Not at all	Just a little	Pretty much	Very much
	0	1	2	3
H Restless (overactive)				
H Excitable, impulsive				
C Disturbs other children				
H Fails to finish things he starts (short attention)				
H fidgeting				
H Inattentive, distractible				
H Demands must be met immediately; frustrated				
M Cries				
M Mood changes quickly				
C Temper outbursts				
M Anxious				
M Overly sad				

In comparison with other children his or her own age, how would you describe your child?

	Better than most 1	Average 2	Slower than most 3
language skills			
physical skills			
social skills			
emotional maturity			
problem solving ability			

BEHAVIOR CHECKLIST FOR TODDLERS

CHILD'S NAME _____ PARENT'S NAME _____

DATE _____

		NO PROBLEM	SOME PROBLEM	MAJOR PROBLEM
He	Eating	_____	_____	_____
He	Sleeping	_____	_____	_____
H	Overactivity	_____	_____	_____
H	Concentration	_____	_____	_____
R	Attention Seeking	_____	_____	_____
M	Moods	_____	_____	_____
M	Worries	_____	_____	_____
M	Fears	_____	_____	_____
C	Difficult to Manage	_____	_____	_____
C	Temper Tantrums	_____	_____	_____
C	Agression	_____	_____	_____
R	Shyness	_____	_____	_____
M	Withdrawn	_____	_____	_____
S	Echoes speech	_____	_____	_____
S	Repetitive habits (flapping, twirling)	_____	_____	_____
R	Relationships with brothers/sisters	_____	_____	_____
R	Overly Dependent	_____	_____	_____
S	Tics	_____	_____	_____
	Stuttering/Stammering	_____	_____	_____
R	Poor eye contact	_____	_____	_____
M	Cries or laughs too easily	_____	_____	_____
H	Is impulsive	_____	_____	_____
C	Defiant	_____	_____	_____
S	Peculiar Pre- occupations	_____	_____	_____
S	Rocks back and forth	_____	_____	_____
S	Bangs head	_____	_____	_____

DOES YOUR CHILD HAVE PROBLEMS WHICH CONCERN YOU IN ANY OTHER
AREAS OF BEHAVIOR OR DEVELOPMENT?

APPENDIX C
CODING FORMS

CHILDHOOD PERSONALITY SCALE

SCORE FORM

NAME _____

CODER _____

AGE _____

DATE _____

HYPERACTIVITY		MOOD		RELATIONSHIPS	
Question #	# of Points	Question #	# of Points	Question #	# of Points
H ₁		M ₁		R ₁	
H ₂		M ₂		R ₂	
H ₃ *		M ₃		R ₃	
H ₄		M ₄ *		R ₄	
H ₅		M ₅		R ₅ *	
H ₆		M ₆		TOTAL	
H ₇ *		M ₇			
H ₈		M ₈		CONDUCT	
H ₉		M ₉		Question #	# of Points
H ₁₀				C ₁	
H ₁₁				C ₂	
H ₁₂					
H ₁₃					
H ₁₄					
TOTAL		TOTAL		TOTAL	

* Reverse Scoring

BEHAVIOR CHECKLIST FOR TODDLERS

SCORE FORM

NAME _____

CODER _____

AGE _____

DATE _____

HYPERACTIVITY		RELATIONSHIPS		MOOD	
Question #	# of Points	Question #	# of Points	Question #	# of Points
H ₁		R ₁		M ₁	
H ₂		R ₂		M ₂	
H ₃		R ₃		M ₃	
H ₄		R ₄		M ₄	
		R ₅		M ₅	
TOTAL		TOTAL		TOTAL	

STEREOTYPICAL BEHAVIORS		CONDUCT		HEALTH	
Question #	# of Points	Question #	# of Points	Question #	# of Points
S ₁		C ₁		He ₁	
S ₂		C ₂		He ₂	
S ₃		C ₃		He	
S ₄		C ₄			
S ₅					
S ₆					
TOTAL		TOTAL		TOTAL	

APPENDIX D
SUBJECT DATA

CHILDHOOD PERSONALITY SCALENORMAL

	H	R	M	C
12	7	5	2	0
14	17	11	6	1
27	32	0	2	1
32	26	20	11	1
36	14	5	7	1
39	17	8	5.5	2
40	25	5	3	1
50	28	2	1	1
55	11	3	1	0
56	30	8	6	2
58	29.5	6	9.5	2.5
59	22	7	7	1
63	26	6	6	1
72	37	12	14	2
78	14	0	1	0
81	33	1	4	1
113	29	9	6	3
126	9	6	4	0
128	32	4	3	4
129	11	1	2	1
130	30	5	3	1
131	38	6	4	2
132	20	4	4	1
133	8	4	4	2
138	34	19	3	3
139	21	4	10	2
141	20	7	8	1
144	22	3	5	1
150	36	7	5	1
<u>TOTAL</u>	678.5	178	147	39.5
<u>MEAN</u>	23.40	6.14	5.07	1.36
<u>S. D.</u>	9.29	4.71	3.16	.95

BEHAVIOR CHECKLIST FOR TODDLERSNORMAL

	H	R	M	S	C	He
12	0	1	0	1	1	1
14	0	0	0	0	0	1
27	0	0	1	0	2	0
32	1	2	1	0	3	2
36	0	0	0	0	0	1
39	0	0	0	1	2	0
40	0	0	0	0	0	0
41	0	0	0	0	0	0
50	0	0	0	0	1	0
55	0	0	0	0	0	0
56	0	2	1	1	1	4
58	0	0	3	0	3	0
59	0	0	1	0	0	1
63	1	1	2	1	1	0
72	2	1	2	4	2	1
78	0	0	0	0	2	1
81	2	0	0	0	4	1
113	0	0	1	0	0	0
126	0	1	0	0	0	0
128	0	1	0	0	2	0
129	0	0	1	0	0	0
130	0	0	0	0	0	0
131	2	1	1	1	3	2
132	0	2	0	0	2	0
133	0	2.5	2	0	0	1
138	0	1	0	0	1	2
139	0	2	1	0	1	0
141	0	0	0	0	1	0
144	1	1	1	1	3	0
150	0	0	2	0	0	1
<u>TOTAL</u>	9	18.5	20	10	35	19
<u>MEAN</u>	.30	.62	.67	.33	1.17	.63
<u>S. D.</u>	.65	.81	.84	.80	1.21	.93

CHILDHOOD PERSONALITY SCALEDELAYED

	H	R	M	C
1	22	12	5	1
6	55	14	23	3
7	41	11	10	5
15	23	6	10	1
19	33	19	6	2
29	39	4	13	3
52	32	1	8	2
53	25	1	2	0
54	27	8	5	2
57	28	18	13	1
83	38	5	5	3
85	24	9	8	3
87	25	6	6	3
88	30	8	3	0
89	45	7	10	2
90	29	13	5	2
91	20	1	2	3
92	11	13	2	1
93	24	2	6.5	1
94	28	18	1	2
98	35	7	4	2
102	43	4	8	1
103	11	10	11	1
105	37	7	4	2
107	22	8	7	3
111	33	21	9	6
112	25	9	7	1
114	31	16	12	2
115	36	4	5	2
116	40	1	3	5
119	23.5	4	4	1
142	32	2	15	4
145	31	18	23	2
<u>TOTAL</u>	998.5	287	255.5	74
<u>MEAN</u>	30.26	8.70	7.74	2.18
<u>S. D.</u>	9.23	5.88	5.30	1.38

BEHAVIOR CHECKLIST FOR TODDLERSDELAYED

	H	R	M	S	C	He
1	0	0	0	0	0	0
6	5	3	2	0	3	1
7	3	2	1	1	3	0
15	2	1	0	0	0	1
19	3	4	4	2	3	1
29	4	0	1	2	5	1
52	1	0	3	0	2	0
53	0	0	1	0	0	0
54	2	0	0	0	4	0
57	0	3	2	1	1	0
83	0	5	1	0	4	1
85	0	2	3	0	3	1
87	0	0	0	0	2	1
88	2	3	4	1	0	1
89	2	2	3	0	3	2
90	0	0	0	1	0	1
91	0	2	1	2	2	0
92	0	3	1	0	1	1
93	1	3.5	4	2	2	0
94	0	0	1	0	1	2
98	1	0	0	0	1	0
100	1	1	0	0	1	0
102	5	1	0	1	7	2
103	0	2	0	1	0	2
105	0	0	0	0	2	0
107	0	0	0	0	1	1
111	0	3	2	1	2	2
112	1	0	0	0	1	1
114	0	2	0	0	2	2
115	2	1	0	1	4	0
116	1	4	2	0	3	0
119	2	2.5	0	0	2.5	1
142	0	0	0	0	2	0
145	2	5	4	2	4	0
<u>TOTAL</u>	40	55	40	18	71.5	25
<u>MEAN</u>	1.18	1.62	1.18	.53	2.10	.74
<u>S. D.</u>	1.47	1.60	1.42	.75	1.62	.75

CHILDHOOD PERSONALITY SCALELATE TALKERS

	H	R	M	C
4	37	13	13	3
9	23	3	2	2
26	35	15.5	9.5	1
51	26	2	4	4
86	27	5	6	2
95	39	3	10	3.5
97	25	7	4	2
101	34	10	12	1
109	45	8	20	5
122	30	7	13	3
184	21	11.5	11	1
<u>TOTAL</u>	342	85	104.50	27.50
<u>MEAN</u>	31.09	7.73	9.50	2.50
<u>S.D.</u>	7.50	4.40	5.22	1.32

BEHAVIOR CHECKLIST FOR TODDLERSLATE TALKERS

	H	R	M	S	C	He
4	2	2	0	0	6	2
9	1	0	0	0	0	2
26	2	2	1	1	0	0
51	0	0	0	0	0	0
60	0	2	0	0	2	1
86	0	0	0	0	0	0
95	2	2	0	0	4	1
97	0	0	0	1	2	1
101	2	1	0	0	3	0
109	1	3	4	0	7	0
122	1	1	1	0	2	1
184	0	1	1	0	1	1
<u>TOTAL</u>	11	14	7	2	25	9
<u>MEAN</u>	.92	1.17	.58	.17	2.08	.75
<u>S.D.</u>	.90	1.03	1.17	.39	2.38	.75

KEY:

H-Hyperactivity/Attention Deficit Disorder

R-Relationships

M-Mood

C-Conduct

S-Stereotypical Behaviors

He-Health