

LOWER COLUMBIA RIVER AQUATIC NONINDIGENOUS SPECIES SURVEY 2001-2004

Final Technical Report: Appendices

Mark D. Sytsma

Center for Lakes and Reservoirs
Portland State University
Portland OR 97207

Jeffery R. Cordell

Wetland Ecosystem Team
University of Washington
Seattle WA 98195

John W. Chapman

Hatfield Marine Science Center
Oregon State University
Newport OR 97365

Robyn C. Draheim

Center for Lakes and Reservoirs
Portland State University
Portland OR 97207

Prepared for the
UNITED STATES COAST GUARD
and the
UNITED STATES FISH AND WILDLIFE SERVICE

October 2004

Appendices: Contents

APPENDIX A: TAC.....	1
APPENDIX B: SPECIES LIST	2
Kingdom: Monera.....	3
Division: Cyanophycota.....	3
Cyanobacteria	3
Division: Bacillariophyta.....	4
Division: Chlorophyta.....	9
Division: Phaeophycophyta	11
Division: Chrysophyta	11
Division: Pyrrophytophyta	12
Division: Magnoliophyta	12
Kingdom: Animalia	42
Phylum: Porifera	42
Phylum: Ciliophora.....	42
Class: Ciliatea	42
Phylum: Cnidaria	43
Class: Anthozoa	43
Class: Hydrozoa	44
Phylum: Ectoprocta.....	45
Class: Phylactolaemata	45
Phylum: Entoprocta	47
Phylum: Nemertea	48
Class: Enopla	48
Phylum: Annelida	48
SubClass: Oligochaeta	48
Subclass: Polychaeta.....	57
Errant Polychaetes	57
Sedentary Polychaetes	61
Phylum: Mollusca	69
Class: Gastropoda	72
Class: Bivalvia	82
Phylum: Arthropoda.....	88
Subphylum: Crustacea	88
Infraclass: Cirripedia.....	88
Class: Ostracoda.....	90
Subclass: Copepoda	90
Suborder: Cladocera.....	96
Class: Malacostraca	97
Peracarida – Cumacea.....	97
Peracarida – Tanaidacea	100
Peracarida – Isopoda	101
Peracarida – Amphipoda.....	109
Peracarida – Mysida.....	125
Decapoda.....	127

Phylum: Chordata	131
Subphylum: Vertebrata	131
Superclass: Osteichthyes.....	131
Class: Amphibia.....	161
Class: Reptilia	162
Order: Testudines.....	162
Class: Mammalia	163
Order: Rodentia.....	163

APPENDIX A: TAC

Technical Advisory Committee Participants

Name	Affiliation
Jim Athearn	US Army Corps of Engineers
Jim Carlton	Williams College
Andy Cohen	San Francisco Estuary Institute
Sebastian Degens	Port of Portland
Mark Dybdahl	Washington State University
Rich Everett	US Coast Guard
Steve Fend	USGS – Menlo Park
Terry Frest	Deixis Consultants
Jon Graves	Oregon Graduate Institute
Kathy Hamel	Washington Department of Ecology
Leslie Harris	Los Angeles County Museum of Natural History
Paul Heimowitz	U.S. Fish and Wildlife Service (formerly of) Oregon Sea Grant
Denny Lassuy	US Fish and Wildlife Service
Henry Lee II	USEPA Coastal Ecology
Claudia Mills	University of Washington
Annette Olson	University of Washington
Blaine Parker	Columbia River Inter-Tribal Fish Commission
Jennifer Parsons	Washington Department of Ecology
Greg Ruiz	Smithsonian Environmental Research Center
Scott Smith	Washington Department of Fish and Wildlife
David Strayer	Institute of Ecosystem Studies (formerly of) Lower Columbia River Estuary Program
Bruce Sutherland	Columbia River Steamship Operators Association
Jim Townley	Columbia River Estuary Studies Taskforce
Matthew VanEss	Oregon Department of Fish and Wildlife
David Ward	(formerly of) U.S. Fish and Wildlife Service
Erin Williams	

APPENDIX B: SPECIES LIST

Guide to the format of this section

- Species arranged by Phylum/Division, then Class and/or other relevant taxonomic breakdown
- List of all species compiled from literature review and the field surveys

Family	<i>Species Name</i>	LCRANS = present in survey, LIT= present in literature review	Origin
--------	---------------------	---	--------

- Species Descriptions
Species Name, Author
Synonyms (if applicable)
Source of Information (LCRANS, LIT)
Origin (i.e. Introduced, Cryptogenic or Native)
Descriptive paragraph

Kingdom: Monera
Phylum: Cyanophycota

Kingdom: Monera

Division: Cyanophycota

Cyanobacteria

There are 124 freshwater genera of cyanobacteria or blue-green algae reported from North America, however this division is in a state of taxonomic flux (Sheath and Wehr 2003). Cyanobacteria can be important in surface blooms, often toxic, in nutrient rich waters. All three genera below are widely distributed (Komarek 2003).

Nostocaceae	
<i>Anabaena</i> sp.	LCRANS, LIT
<i>Aphanizomenon flosaquae</i>	LIT
<i>Aphanizomenon</i> sp.	LIT
Oscillatoriaceae	
<i>Lyngbya</i> sp.	LIT
<i>Oscillatoria</i> sp.	LCRANS
<i>Phormidium</i> sp.	LCRANS
<i>Spirulina</i> sp.	LIT

Kingdom: Plantae

Division: Bacillariophyta

Phytoplankton species are the most common of all groups found in ballast water entering eastern Pacific ports (Carlton and Geller 1993, Levings et al. 2004, Cohen 1998).

Achnantheaceae	
<i>Achnanthes deflexa</i>	LIT
<i>Achnanthes lemmermannii</i>	LIT
<i>Achnanthes suchlandtii</i>	LIT
<i>Achnanthes</i> sp.	LIT
<i>Karayevia clevei</i>	LIT
<i>Planothidium hauckianum</i>	LIT
<i>Planothidium lanceolatum</i>	LIT
<i>Planothidium peragalli</i>	LIT
<i>Rossithidium linearis</i>	LIT
Achnanthidiaceae	
<i>Achnanthidium minutissimum</i>	LIT
Amphipleuraceae	
<i>Frustulia rhomboids</i>	LIT
Asterolampraceae	
<i>Asteromphalus heptactis</i>	LIT
Aulacoseiraceae	
<i>Aulacoseira ambigua</i>	LIT
<i>Aulacoseira distans</i>	LIT
<i>Aulacoseira granulata</i>	LCRANS, LIT
<i>Aulacoseira granulata f. spiralis</i>	LCRANS
<i>Aulacoseira islandica</i>	LIT
<i>Aulacoseira italica</i>	LCRANS
Bacillariaceae	
<i>Bacillaria paxillifer</i>	LCRANS
<i>Cylindrotheca closterium</i>	LCRANS, LIT
<i>Cylindrotheca gracilis</i>	LIT
<i>Cymbellonitzschia diluviana</i>	LIT
<i>Hantzschia amphioxys</i>	LCRANS
<i>Hantzschia distinctepunctata</i>	LIT
<i>Hantzschia marina</i>	LIT
<i>Nitzschia accuminata</i>	LIT
<i>Nitzschia acicularis</i>	LIT
<i>Nitzschia amphibian</i>	LIT
<i>Nitzschia capitellata</i>	LIT
<i>Nitzschia dissipata</i>	LIT
<i>Nitzschia frustulum</i>	LIT
<i>Nitzschia holsatica</i>	LIT
<i>Nitzschia linearis</i>	LIT

<i>Nitzschia longissima</i>	LCRANS
<i>Nitzschia palea</i>	LIT
<i>Nitzschia paleacea</i>	LIT
<i>Nitzschia parvula</i>	LIT
<i>Nitzschia pungens</i>	LCRANS
<i>Nitzschia recta</i>	LIT
<i>Nitzschia seriata</i>	LIT
<i>Nitzschia sigma</i>	LCRANS, LIT
<i>Nitzschia sigmoidea</i>	LIT
<i>Nitzschia subhybrida</i>	LIT
<i>Nitzschia sublinearis</i>	LIT
<i>Nitzschia vermicularis</i>	LCRANS
<i>Nitzschia vitrea</i>	LCRANS
<i>Nitzschia</i> sp.	LIT
<i>Tryblionella angustata</i>	LIT
<i>Tryblionella apiculata</i>	LIT
<i>Tryblionella hungarica</i>	LIT
<i>Tryblionella victoriae</i>	LIT
Berkeleyaceae	
<i>Berkeleya rutilans</i>	LIT
Biddulphiaceae	
<i>Eucampia zodiacus</i>	LCRANS
Catenulaceae	
<i>Amphora angusta</i>	LIT
<i>Amphora coffaeiformis</i>	LIT
<i>Amphora micrometra</i>	LIT
<i>Amphora ovalis</i>	LCRANS, LIT
<i>Amphora perpusilla</i>	LIT
<i>Amphora sabyii</i>	LIT
<i>Amphora tenerrima</i>	LIT
<i>Amphora</i> sp.	LCRANS
Chaetocerotaceae	
<i>Bacteriastrum delicatulum</i>	LIT
<i>Bacteriastrum hyalinum</i>	LCRANS
<i>Chaetoceros convolutes</i>	LIT
<i>Chaetoceros decipiens</i>	LCRANS, LIT
<i>Chaetoceros didymus</i>	LIT
<i>Chaetoceros radicans</i>	LIT
<i>Chaetoceros</i> sp.	LCRANS, LIT
Cocconeidaceae	
<i>Cocconeis klamathensis</i>	LIT
<i>Cocconeis placentula</i>	LCRANS, LIT
Coscinodiscaceae	
<i>Coscinodiscus apiculatus</i>	LIT
<i>Coscinodiscus centralis</i>	LIT
<i>Coscinodiscus curvatulus</i>	LIT

<i>Coscinodiscus excentricus</i>	LIT
<i>Coscinodiscus hantzschii</i>	LCRANS
<i>Coscinodiscus perforatus</i>	LIT
<i>Coscinodiscus radiatus</i>	LCRANS
<i>Coscinodiscus</i> sp.	LIT
Cymbellaceae	
<i>Cymbella affinia</i>	LIT
<i>Cymbella cuspidate</i>	LIT
<i>Cymbella elginis</i>	LCRANS, LIT
<i>Cymbella</i> sp.	LIT
<i>Encyonema minutum</i>	LIT
<i>Placoneis gastrum</i>	LIT
<i>Placoneis placentula</i>	LIT
Diploneidaceae	
<i>Diploneis fasca</i> var. <i>pelagica</i>	LCRANS
<i>Diploneis puella</i>	LIT
<i>Diploneis smithii</i>	LIT
<i>Diploneis subovalis</i>	LIT
<i>Diploneis</i> sp.	LCRANS
Eupodiscaceae	
<i>Odontella longicuris</i>	LCRANS, LIT
<i>Odontella aurita</i>	LIT
Fragilariaceae	
<i>Asterionella formosa</i>	LCRANS, LIT
<i>Asterionella gracillima</i>	LCRANS
<i>Asterionella japonica</i>	LIT
<i>Asterionellopsis glacialis</i>	LCRANS
<i>Diatoma hiemale</i> var. <i>mesodon</i>	LIT
<i>Diatoma tenue</i>	LIT
<i>Diatoma tenue</i> var. <i>elongatum</i>	LIT
<i>Diatoma vulgare</i>	LCRANS, LIT
<i>Diatoma vulgare</i> var. <i>breve</i>	LIT
<i>Fragilaria capucina</i>	LCRANS, LIT
<i>Fragilaria crotonensis</i>	LCRANS, LIT
<i>Fragilaria oceanica</i>	LIT
<i>Meridion circulare</i>	LIT
<i>Staurosira contruens</i>	LIT
<i>Synedra ulna</i>	LCRANS, LIT
<i>Synedra delicatissima</i>	LCRANS
Gomphonemataceae	
<i>Gomphonema acuminatum</i>	LCRANS
<i>Gomphonema</i> sp.	LCRANS
<i>Reimeria sinuata</i>	LIT
Heliopeltaceae	
<i>Actinoptychus senarius</i>	LCRANS
<i>Actinoptychus splendens</i>	LCRANS

Hemidiscaceae	
<i>Actinocyclus ehrenbergii</i>	LCRANS
Lauderiaceae	
<i>Lauderia annulata</i>	LCRANS
Lithodesmiaceae	
<i>Ditylum brightwellii</i>	LCRANS
Melosiraceae	
<i>Melosira italica</i>	LCRANS
<i>Melosira nummuloides</i>	LCRANS
<i>Melosira varians</i>	LCRANS
Naviculaceae	
<i>Amphiprora gigantea</i> var <i>sulcata</i>	LCRANS
<i>Navicula elegans</i>	LCRANS
<i>Navicula</i> sp.	LCRANS
Pinnulariaceae	
<i>Pinnularia</i> sp.	LCRANS
Pleurosigmataceae	
<i>Gyrosigma</i> sp.	LCRANS
<i>Pleurosigma fasciola</i>	LCRANS
<i>Pleurosigma</i> sp.	LCRANS
Rhizosoleniaceae	
<i>Proboscia alata</i>	LCRANS
<i>Rhizosolenia setigera</i>	LCRANS
Skeletonemaceae	
<i>Skeletonema costatum</i>	LCRANS
<i>Skeletonema tropicum</i>	LCRANS
Stephanodiscaceae	
<i>Cyclotella comta</i>	LCRANS
<i>Cyclotella meneghiniana</i>	LCRANS
<i>Stephanodiscus hantzschii</i>	LCRANS
Surirellaceae	
<i>Surirella caproni</i>	LCRANS
<i>Surirella linearis</i>	LCRANS
Tabellariaceae	
<i>Tabellaria fenestrata</i>	LCRANS
Thalassionemataceae	
<i>Thalassionema nitzschioides</i>	LCRANS
Thalassiosiraceae	
<i>Thalassiosira lineatus</i>	LCRANS
<i>Thalassiosira pacificia</i>	LCRANS
<i>Thalassiosira punctigera</i>	LCRANS

Division: Chlorophyta

Green Algae

The division Chlorophyta includes both planktonic forms and macroalgal species as well as marine, estuarine and freshwater species. Filamentous green algae can often form free-floating mats or may be intertwined with other algal masses attached to hard surfaces (Shubert 2003). Macroalgae were not actively collected and identified during the LCRANS survey.

Chlorococcaceae		
<i>Schroederia setigera</i>	LCRANS	Cryptogenic
Dictyosphaeriaceae		
<i>Dictyosphaerium pulchellum</i>	LCRANS	Cryptogenic
Hydrodictyaceae		
<i>Pediastrum integrum</i>	LCRANS	Cryptogenic
<i>Pediastrum</i> sp.	LCRANS	
Scenedesmaceae		
<i>Actinastrum hantzschii</i>	LCRANS	Cryptogenic
<i>Scenedesmus longispina</i>	LCRANS	Cryptogenic
<i>Scenedesmus</i> sp.	LCRANS	
Ulvaceae		
<i>Enteromorpha intestinalis</i>	LIT	Native
Ulva	LCRANS, LIT	
Volvocaceae		
<i>Eudorina elegans</i>	LCRANS	Cryptogenic
<i>Eudorina</i> sp.	LIT	

CHLOROCOCCACEAE

Schroederia setigera (Schroeder) Lemmermann

Synonyms: *Ankistrodesmus setiguris*, *Reinschiella setigera*

LCRANS

Origin Cryptogenic

Freshwater planktonic alga. Widely reported and common in the plankton of North America (Shubert 2003). Also found in Europe and Asia.

DICTYOSPHAERIACEAE

Dictyosphaerium pulchellum Wood, 1872

LCRANS

Origin Cryptogenic

Kingdom: Plantae
Division: Chlorophyta

Colonial form. This genus is common but not considered abundant in North America (Shubert 2003).

HYDRODICTYACEAE

Pediastrum integrum Naeg.

LCRANS

Origin Cryptogenic

The genus is found in all regions of North America (Shubert 2003).S

SCENEDESMACEAE

Actinastrum hantzschii Lagerheim, 1882

LCRANS

Origin Cryptogenic

Colonial alga. Genus is widely reported from North America, common in ditches, ponds, bogs and lakes (Shubert 2003).

Scenedesmus longispina Meyen

LCRANS

Origin Cryptogenic

The most commonly reported genus of coccoid green algae worldwide (Shubert 2003).

ULVACEAE

Enteromorpha intestinalis (L.) Link

Synonyms: *Ulva intestinalis*

LIT

Origin: Native

Found on rocks in the high to mid tidal zone in protected bays and estuaries from Alaska to Mexico (Abbott and Hollenberg 1976). The genus *Enteromorpha* is cosmopolitan.

VOLVOCAACEAE

Eudorina elegans

LCRANS

According to Shubert (2003) *Eudorina elegans* is among the most frequently encountered species of green algae.

Kingdom: Plantae
Division: Chlorophyta

Division: Phaeophycophyta

The brown algae

Macroalgae were not actively collected and identified during the LCRANS survey but *Fucus distichus* was noted because of its abundance at Trestle Bay and Baker Bay sites.

Fucaceae		
<i>Fucus distichus</i>	LCRANS, LIT	Native

FUCACEAE

Fucus distichus Linnaeus 1767
LCRANS, LIT
Native

Found attached to rocks in the upper to mid-intertidal zone from northern Washington State to Point Conception, California (Abbott and Hollenberg 1976). Dominant macrophyte in the intertidal zone in Trestle Bay and Baker Bay.

Division: Chrysophyta

Silicaflagellates

There are 72 genera of silicaflagellates known from inland habitats in North America, freshwater species are typically associated with standing bodies of water (Sheath and Wehr 2003). The skeletons of silicoflagellates usually comprise 1-2% of the siliceous component of marine sediments; making them much less abundant than diatoms. Marine species can contribute to blooms and are widely distributed throughout the world's oceans (McCartney 1993).

Dictyochaceae		
<i>Dictyocha fibula</i>	LCRANS	Cryptogenic

DICTYOCHACEAE

Dictyocha fibula Ehrenb.
LCRANS
Origin: Cryptogenic

Marine species, also known from the eastern Atlantic.

Kingdom: Plantae
Division: Pyrrophytophyta

Division: Pyrrophytophyta

The dinoflagellates

Dinoflagellates are typically a minor component of the phytoplankton and at times form dense blooms – usually in the presence of high levels of nitrates and phosphates (Sheath and Wehr 2003).

Ceratiaceae		
<i>Ceratium hirundiella</i>	LCRANS	Cryptogenic
Protopteridinaceae		
<i>Protopteridinium depressum</i>	LCRANS	Cryptogenic

CERATIACEAE

Ceratium hirundiella

LCRANS

Cryptogenic

Freshwater dinoflagellate, found throughout North America, distributed worldwide.

PROTOPTERIDINACEAE

Protopteridinium depressum

LCRANS

Cryptogenic

Marine dinoflagellate, distributed worldwide

Subkingdom: Tracheobionta

Division: Magnoliophyta

Aquatic vascular plants include a variety of lifeforms including submersed and emergent, free-floating and rooted species. Submersed species are restricted to shallow water, low current-velocity sites due to light and scouring effects. Emergent species occur are common on islands in the lower Columbia River. Emergent species are typically included in wetland delineation work, however, submersed species are often overlooked. LCRANS sampling focused on cataloging introduced submersed species, although introduced emergent species were noted when observed. Submersed and emergent species were included in the literature review.

Alismataceae		
<i>Alisma</i> spp.	LCRANS	
<i>Alisma triviale</i>	LIT	Native
<i>Sagittaria cuneata</i>	LIT	Native
<i>Sagittaria latifolia</i>	LIT	Native#
<i>Sagittaria</i> spp.	LIT	
Apiaceae		
<i>Angelica lucida</i>	LIT	Native
<i>Heracleum maximum</i>	LIT	Native
<i>Hydrocotyle ranunculoides</i>	LCRANS, LIT	Native
<i>Lilaeopsis occidentalis</i>	LIT	Native
<i>Oenanthe sarmentosa</i>	LIT	Native
<i>Sium suave</i>	LIT	Native
Araceae		
<i>Lysichiton americanus</i>	LIT	Native
Asteraceae		
<i>Achillea millefolium</i>	LIT	Cryptogenic
<i>Aster</i> spp.	LIT	
<i>Aster subspicatus</i>	LIT	Native
<i>Bidens cernua</i>	LIT	Native
<i>Boltonia asteroides</i>	LIT	Cryptogenic
<i>Canadanthus modestus</i>	LIT	Native
<i>Cotula coronopifolia</i>	LCRANS, LIT	Introduced
<i>Helenium autumnale</i>	LIT	Native
<i>Senecio triangularis</i>	LIT	Native
Azollaceae		
<i>Azolla mexicana</i>	LCRANS	Native
Boraginaceae		
<i>Myosotis laxa</i>	LIT	Native
Cabombaceae		
<i>Cabomba caroliniana</i>	LCRANS, LIT	Introduced
Callitrichaceae		
<i>Callitriche stagnalis</i>	LCRANS, LIT	Introduced
<i>Callitriche verna</i>	LCRANS, LIT	Native
<i>Callitriche</i> spp.	LIT	
Ceratophyllaceae		
<i>Ceratophyllum demersum</i>	LCRANS, LIT	Native
Chenopodiaceae		
<i>Salicornia depressa</i>	LIT	Native
Clusiaceae		
<i>Hypericum scouleri</i>	LIT	Native
Commelinaceae		
<i>Murdannia keisak</i>	LIT	Introduced*
Crassulaceae		
<i>Crassula aquatica</i>	LIT	Native

Cyperaceae		
<i>Carex lyngbyei</i>	LIT	Native
<i>Carex obnupta</i>	LIT	Native
<i>Carex</i> spp.	LCRANS, LIT	
<i>Eleocharis minima</i>	LIT	Native [#]
<i>Eleocharis palustris</i>	LIT	Native
<i>Eleocharis</i> spp.	LCRANS, LIT	
<i>Schoenoplectus americanus</i>	LIT	Native
<i>Schoenoplectus tabernaemontani</i>	LIT	Native
<i>Schoenoplectus maritimus</i>	LIT	Native
<i>Schoenoplectus robsutus</i>	LIT	Native [#]
<i>Scirpus microcarpus</i>	LIT	Native
<i>Scirpus</i> spp.	LIT	
Dictyosphaeriaceae		
<i>Dictyosphaerium</i> sp	LIT	
Equisetaceae		
<i>Equisetum fluviatile</i>	LIT	Native
Fabaceae		
<i>Lathyrus palustris</i>	LIT	
<i>Lupinus</i> sp.	LIT	
<i>Trifolium</i> spp.	LIT	
<i>Vicia nigricans</i> ssp. <i>gigantea</i>	LIT	Native
Haloragaceae		
<i>Myriophyllum aquaticum</i>	LCRANS, LIT	Introduced
<i>Myriophyllum sibiricum</i>	LCRANS	Native
<i>Myriophyllum spicatum</i>	LCRANS, LIT	Introduced
Hydrocharitaceae		
<i>Egeria densa</i>	LCRANS, LIT	Introduced
<i>Elodea canadensis</i>	LCRANS, LIT	Native
<i>Elodea nuttallii</i>	LCRANS, LIT	Native
<i>Vallisneria americana</i>	LCRANS, LIT	Introduced
Iridaceae		
<i>Iris pseudacorus</i>	LCRANS, LIT	Introduced
Isoetaceae		
<i>Isoetes tenella</i>	LIT	Native
Juncaceae		
<i>Juncus balticus</i>	LIT	Native
<i>Juncus effusus</i>	LIT	Native
<i>Juncus filiformis</i>	LIT	Native
<i>Juncus nevadensis</i>	LIT	Native
<i>Juncus oxymersis</i>	LIT	Native
Juncaginaceae		
<i>Triglochin maritimum</i>	LIT	Native
Lamiaceae		
<i>Mentha arvensis</i>	LIT	Native
<i>Mentha aquatica</i>	LIT	Introduced

<i>Mentha aquatica x spicata</i>	LIT	Introduced
<i>Mentha</i> spp.	LIT	
<i>Prunella vulgaris</i>	LIT	Native
Lemnaceae		
<i>Lemna minor</i>	LCRANS, LIT	Native
Liliaceae		
<i>Veratrum californicum</i>	LIT	Native
Lythraceae		
<i>Lythrum salicaria</i>	LCRANS, LIT	Introduced
Menyanthaceae		
<i>Nephrrophyllidium crista-galli</i>	LIT	Native
Najadaceae		
<i>Najas</i> sp.	LIT	
Nymphaeaceae		
<i>Nymphaea odorata</i> spp. <i>odorata</i>	LCRANS, LIT	Introduced
Onagraceae		
<i>Epilobium ciliatum</i> ssp. <i>watsonii</i>	LIT	Native
<i>Ludwigia uruguayensis</i>	LIT	Introduced
Orchidaceae		
<i>Platanthera dilatata</i> var. <i>dilatata</i>	LIT	Native
Plantaginaceae		
<i>Littorella</i> sp.	LIT	
<i>Plantago lanceolata</i>	LIT	Introduced
Poaceae		
<i>Beckmannia syzigachne</i>	LIT	Cryptogenic
<i>Deschampsia caespitosa</i>	LIT	Cryptogenic
<i>Distichlis spicata</i>	LIT	Native
<i>Elymus glaucus</i>	LIT	Native
<i>Glyceria striata</i>	LIT	Native
<i>Hordeum brachyantherum</i>	LIT	Native
<i>Lolium arundinacea</i>	LIT	Introduced
<i>Phalaris arundinacea</i>	LCRANS, LIT	Introduced
<i>Spartina</i> spp.	LIT	Introduced*
Polygonaceae		
<i>Polygonum hydropiperoides</i>	LIT	Native
<i>Polygonum</i> spp.	LCRANS, LIT	
Pontederiaceae		
<i>Eichhornia crassipes</i>	LIT	Introduced*
Potamogetonaceae		
<i>Potamogeton crispus</i>	LCRANS, LIT	Introduced
<i>Potamogeton epihydrus</i>	LCRANS	Native
<i>Potamogeton foliosus</i>	LIT	Native
<i>Potamogeton friesii</i>	LIT	Native
<i>Potamogeton natans</i>	LCRANS	Native
<i>Potamogeton pectinatus</i>	LCRANS	Native
<i>Potamogeton pusillus</i>	LCRANS	Native

Kingdom: Plantae
 Division: Magnoliophyta

<i>Potamogeton richardsonii</i>	LCRANS, LIT	Native
<i>Potamogeton zosteriformis</i>	LCRANS	Native
<i>Potamogeton</i> spp.	LCRANS	
Ranunculaceae		
<i>Caltha asarifolia</i>	LIT	Native
<i>Ranunculus</i> spp.	LI	
Rosaceae		
<i>Argentina anserina</i>	LIT	Native
Rubiaceae		
<i>Galium</i> sp.	LIT	
<i>Galium trifidum</i> ssp. <i>columbianum</i>	LIT	Native
Ruppiaceae		
<i>Ruppia maritima</i>	LCRANS, LIT	Cryptogenic
<i>Ruppia</i> spp.	LIT	
Salicaceae		
<i>Salix hookeriana</i>	LIT	Native
Scrophulariaceae		
<i>Castilleja ambigua</i> ssp. <i>ambigua</i>	LIT	Native
<i>Gratiola ebracteata</i>	LIT	Native
<i>Gratiola neglecta</i>	LIT	Native
<i>Limosella aquatica</i>	LCRANS, LIT	Native
<i>Mimulus guttatus</i>	LIT	Native
Sparganiaceae		
<i>Sparganium erectum</i>	LIT	Introduced*
<i>Sparganium angustifolium</i>	LCRANS	Native
Typhaceae		
<i>Typha angustifolia</i>	LCRANS, LIT	Introduced
<i>Typha latifolia</i>	LIT	Native
<i>Typha</i> spp.	LCRANS, LIT	
Zannichelliaceae		
<i>Zannichellia palustris</i>	LCRANS, LIT	Native
Zosteraceae		
<i>Zostera japonica</i>	LCRANS, LIT	Introduced
<i>Zostera marina</i>	LIT	Native
<i>Zostera</i> sp.	LCRANS, LIT	

= likely mis-identification * = unsuccessful establishment

ALISMATACEAE

Alisma triviale

American water plantain

Syn: *A. brevipes*, *A. plantago-americanum*, *A. plantago-aquatica*, *A. subcordatum*

LIT

Origin: Native

Kingdom: Plantae
Division: Magnoliophyta

Alisma triviale is native to the California floristic province, i.e. from the dry regions of the Great Basin and the Mojave Desert to the Pacific coast to Canada (Hickman 1993). Also found in Southeastern US, Eurasia, eastern Africa, and perhaps Australia.

Sagittaria cuneata Sheldon duck potato, arrowhead, wapato
Syn: *Sagittaria arifolia*
LIT
Origin: Native – probably misidentified

Native to California, Pacific Northwest to Southern Canada (Hickman 1993). Only found east of Cascades in Oregon and Washington (Ecology 2003). May be confused with *S. latifolia* below. May also be confused with *Alisma* spp., *Valisneria* sp., or *Sparganium* spp. which all have ribbon-like underwater leaves but it is unlikely to be confused with other plants when the arrowhead shaped leaves are present (Ecology 2003).

Sagittaria latifolia Willd. duck potato, arrowhead, wapato
Syn: *S. chinensis*, *S. esculenta*, *S. longirostra*, *S. obtuse*, *S. ornithorhyncha*, *S. planipes*, *S. pubescens*, *S. viscosa*
LIT
Origin: Native

Native to California, Pacific Northwest to Southern Canada (Hickman 1993). Unlike *S. cuneata*, *S. latifolia* is common on Pacific coast, and in central, and eastern United States. In Washington it is distributed primarily west of Cascades and the Columbia River Gorge. See above for notes on similar species.

APIACEAE

Angelica lucida L. seacoast angelica
Syn: *Coelopleurum actaeifolium*, *Coelopleurum qmelinii*, *Coelopleurum lucidum*, *Coelopleurum lucidum* ssp.
LIT
Origin: Native

Native to the Pacific coasts of North America and Siberia. Used for medicinal purposes by some Eskimo communities. Also found in coastal areas of Northeastern North America.

Heracleum maximum Bartr. cowparsnip
Syn: *H. lanatum*, *H. sphondylium* var. *lanatum*, *H. sphondylium* ssp. *montanum*
LIT
Origin: Native

Native to North America. Used as a wetland indicator species.

Kingdom: Plantae
Division: Magnoliophyta

Hydrocotyle ranunculoides L.f. floating marsh pennywort
LCRANS, LIT
Origin: Native

Native to Washington, Oregon and British Columbia. On the rare-plant list for Washington and B.C. *Hydrocotyle ranunculoides* is considered an aggressive invader in Australia, the U.K., and parts of Africa.

Lilaeopsis occidentalis Coult. & Rose western grasswort
LIT
Origin: Native

Distributed along the West coast of North America from California to British Columbia (Hickman 1993).

Oenanthe sarmentosa K. Presl ex DC. water parsley
LIT
Origin: Native

Western N. America - British Columbia to California.

Sium suave Walter hemlock water parsnip
Syn: *S. cicutifolium*, *S. floridanum*, *S. suave* var. *floridanum*
LIT
Origin: Native

Native to North America, distributed across the northern states and south to Texas. (Hickman 1993).

ARACEAE

Lysichiton americanus Hultén & St. John western skunk cabbage
Syn: *Lysichiton americanum*, *L. camtschatcensis*
LIT

Native to Western North America (Hickman 1993)..

ASTERACEAE

Achillea millefolium L. western yarrow, milfoil
Syn: *Achillea borealis*, *Achillea lanulosa*
LIT
Origin: Cryptogenic

Kingdom: Plantae
Division: Magnoliophyta

There are both native and introduced phases of *Achillea millefolium* in North America. Introduced and native phases differ primarily in chromosome number and are difficult to distinguish morphologically. Native and introduced phases hybridize. The intricate pattern of morphologic, geographic, and ecologic variation within the species has frustrated all efforts to organize an intraspecific taxonomy on a circumboreal or even a strictly North American basis (Alekssoff, 1999).

Aster subspicatus Nees.
LIT
Origin: Native

Douglas aster

Bidens cernua L.
LIT
Origin: Native

nodding beggartick

Boltonia asteroides
LIT
Origin: Cryptogenic

boltonia aster, white doll's daisy

Patchy distribution east of the Rockies by this native North American daisy indicated that it may have been introduced to the western U.S. According to the USDA database this is *Boltonia asteroides* var. *recognita* (USDA - NRCS 2004).

Canadanthus modestus
Syn: *Aster modestus*
LIT
Origin: Native

Canada aster, giant mountain aster

Native to the Pacific Northwest and Canada, this species is not widespread in the U.S. (USDA - NRCS 2004).

Cotula coronopifolia L.
LCRANS, LIT
Origin: Introduced

brass buttons

Endemic to South Africa, *Cotula coronopifolia* is now also found in North America. On the Pacific Coast the species has become established from British Columbia to California. Its presence on the San Francisco Peninsula was reported in 1878. The introduction of *C. coronopifolia* to California is believed to have been via ship ballast (Cohen and Carlton 1995) and may have been spread by shipping up and down the West Coast.

Helenium autumnale L.
Syn: *Helenium grandiflorum*
LIT
Origin: Native

common sneezeweed

Kingdom: Plantae
Division: Magnoliophyta

Distributed throughout the U.S. *Helenium autumnale* var. *grandiflorum* is most likely to be the species reported in previous literature.

Senecio triangularis Hook. arrowleaf ragwort
LIT
Origin: Native

Native to western North America (Hickman 1993).

AZOLLACEAE

Azolla mexicana Schlecht. & Cham. ex K. Presl Mexican water-fern
LCRANS
Origin: Native

Distribution: Western North America and northern South America. Other similar species of water-fern are found nearly worldwide (Ecology 2003).

BORAGINACEAE

Myosotis laxa Lehm. smallflowered forget-me-not
LIT
Origin: Native

May be confused with *Myosotis scorpioides*, common European forget-me-not (Hickman 1993).

CABOMBACEAE

Cabomba caroliniana Gray fanwort, Carolina fanwort
LCRANS, LIT
Origin: Introduced

Native to North and South America's eastern subtropical-temperate zones, *Cabomba caroliniana* is now found in Europe, Asia and Australia (Ecology 2001). Though the species is native to the southeastern United States it has been introduced to the northeastern US and Oregon. The attractive foliage of *C. caroliniana* has made it popular with the aquarium trade since the 1890's. Still popular, the species has been commercially available for some time. The introduction of *C. caroliniana* has been attributed to discarded aquarium plants. Though the species can reproduce sexually, vegetative fragments are the primary mode of reproduction and dispersal. Once established, *C. caroliniana* can threaten recreational use, navigation and the habitat of native species. This species is considered invasive (Les and Mehrhoff 1999).

Kingdom: Plantae
Division: Magnoliophyta

CALLITRICHACEAE

Callitriche stagnalis Scop.

European pond water-starwort

LCRANS, LIT

Origin: Introduced

Globally widespread, *Callitriche stagnalis* is found in Europe, northern Africa, Asia, Australia and North America. Once introduced to North America, many early collections of the species occurred in coastal areas of the United States. It has been hypothesized that the species initial establishment was in or near seaports, introduced by improper disposal of shipping ballast. The first documented specimens of *Callitriche stagnalis* found in Oregon were collected from an unspecified coastal location in 1871 and Clatsop County in 1902. By the turn of the century, *Callitriche stagnalis* had become a popular plant for aquariums, facilitating the establishment of inland populations via discarded plants. *Callitriche stagnalis* is a prolific seed producer and seeds are possibly the species primary mode of dispersal. The spread of *Callitriche stagnalis* has been comparatively slow, it is not a particularly aggressive colonizer but it will displace native species once it establishes itself. (Philbrick et al. 1998).

Callitriche verna L.

vernal water-starwort, spiny water-starwort

Syn: *Callitriche palustris*

LCRANS, LIT

Origin: Native

C. verna is found throughout the Northern hemisphere and is considered circumboreal.

CERATOPHYLLACEAE

Ceratophyllum demersum L.

coontail, hornwort

LCRANS, LIT

Origin: Native

Ceratophyllum demersum occurs across the entire U.S. and throughout most of Canada (IFAS 2004).

CHENOPODIACEAE

Salicornia depressa Standl.

low saltwort

Syn: *Salicornia europaea*, *Salicornia maritima*, *Salicornia virginica*

LIT

Origin: Native

Kingdom: Plantae
Division: Magnoliophyta

According to the Washington Flora Project *S. virginica* may be the best name for this plant. It is distributed along the Pacific, Gulf and Atlantic coasts of the U.S.

CLUSIACEAE

Hypericum scouleri Hook Scouleri's St. Johnswort
Syn: *H. formosum* ssp. *Scouleri*, *H. formosum* var. *nortoniae*
LIT
Origin: Native

Native to western North America, *Hypericum scouleri* is a well-known medicinal plant.

COMMELINACEAE

Murdannia keisak (Hassk.) Hand.-Maz. Asian spiderwort
Syn: *Anelimia*
LCRANS, LIT
Origin: Introduced – not established

Origin: Introduced throughout the Pacific Northwest and the Southeastern U.S., *Murdannia keisak* is associated with rice culture in East Asia where it is a native plant. According to the Virginia Native Plant Society (2004) it was probably first brought to South Carolina or Louisiana in rice imported for growth in this country. In the United States, it is now found in all eastern coastal states from Delaware to Louisiana, and in Kentucky and Tennessee. The aggressive nature of this plant has now been clearly displayed by its ability to establish itself in freshwater wetlands and crowd out native vegetation by forming a solid mat of vegetation. Even in its native region, this species is a troublesome weed. Not only does it produce thousands of very small seeds, it can reproduce vegetatively. It was found in a freshwater tidal marsh on Lois Island in the Columbia River estuary. The island was resurveyed by Portland State University and Washington Department of Ecology in November, 1997 and again during LCRANS but no *M. keisak* was found.

CRASSULACEAE

Crassula aquatica (L.) Schoenl Water pygmy weed
Syn: *Tillaea aquatica* L. H&C
LIT
Origin: Native

Crassula aquatica is native to North America but is considered a rare or threatened species in many states (Rook 2002). It grows in a variety of location types including

Kingdom: Plantae
Division: Magnoliophyta

vernal pools, ponds and the edges of lakes, and may also be found in salt marshes (Hickman 1993).

CYPERACEAE

Carex lyngbyei Hornem.

Lyngby's sedge

Syn: *Carex cryptocarpa*, *Carex cryptochlaena*

LIT

Origin: Native

A tidal wetland species, *Carex lyngbyei* is common in Pacific Northwest marsh communities. It is native to the west coast and ranges from the central coast of California to Alaska (Hickman 1993).

Carex obnupta Bailey

slough sedge

Syn: *Carex magnifica*

LIT

Origin: Native

Carex obnupta is native to the west coast of North America. It can be found along the Pacific Coast from California to British Columbia. It grows in bogs, marshes, wet meadows, ditches and the edges of rivers and lakes. It is very common in areas where fresh and salt water meet but is confined to lower elevations. Hickman 1993 considers it to be a horticultural variety.

Eleocharis minima

hairgrass, small spike rush

Syn: *Eleocharis bicolor*, *Eleocharis uncialis*

LIT

Origin: Native

May be a misidentification as this species may be confused with other *Eleocharis*. The USDA distribution map does not show this species in Oregon or Washington but considers it to be native to North America (USDA - NRCS 2004). It is used as cool-water aquarium plant.

Eleocharis palustris (L.) Roemer & J.A. Schultes

common spike rush

Syn: *Eleocharis mamillata*, *Eleocharis perlonga*, *Eleocharis smallii*, *Eleocharis xyridiformis*

LIT

Origin: Native

A native species, *Eleocharis palustris* is found widely throughout North America (USDA - NRCS 2004).

Kingdom: Plantae
Division: Magnoliophyta

Schoenoplectus tabernaemontani (K.C. Gmel.) Palla softstem bulrush
LIT
Origin: Native

A native sedge *Schoenoplectus tabernaemontani* is distributed throughout North America (USDA - NRCS 2004).

Schoenoplectus americanus (Pers.) Volk. ex Schinz & R. Keller chairmaker's bulrush
syn: *Scirpus americanus*
LIT
Origin: Native

This native sedge is can be found throughout much of North America with the exception of the great lakes region (USDA - NRCS 2004).

Schoenoplectus maritimus (L.) Lye cosmopolitan bulrush
Syn: *Scirpus maritimus*
LIT
Origin: Native

Schoenoplectus maritimus, a native bulrush, can be found throughout much of North America.

Schoenoplectus robustus (Pursh) M.T. Strong sturdy bulrush
Syn: *Scirpus robutus*
LIT

Although it is native to North America this record may represent a mis-identification as the USDA has no record of this species occurring in OR. There are, however, many other species of sedges that may be confused with this one.

Scirpus microcarpus J.& K. Presl paniced bulrush
Syn: *Scirpus rubrotinctus* Fern.
LIT
Origin: Native

Native to North America, this sedge is distributed throughout much of the U.S. except the Southeast.

EQUISETACEAE

Equisetum fluviatile L. water horsetail

Kingdom: Plantae
Division: Magnoliophyta

LIT

Origin: Native

An ancient plant with a circumboreal distribution, *Equisetum fluviatile* commonly grows in dense colonies along shorelines or in shallow water. Most often confused with marsh horsetail (*E. palustre*).

FABACEAE

Lathyrus palustris L.

marsh pea

LIT

Origin: Native

Native to much of the U.S. *Lathyrus myrtifolius* is a state listed threatened and/or endangered species along much of the east coast (USDA - NRCS 2004).

Vicia nigricans* ssp. *gigantea (Hook.) Lassetter & Gunn.

giant vetch

LIT

Origin: Native

Native to Western N. America , *Vicia nigricans* ssp. *gigantea* is found from Alaska to California (Hickman 1993).

HALORAGACEAE

Myriophyllum aquaticum (Vell.) Verdc. parrot feather watermilfoil

LCRANS, LIT

Origin: Introduced

Myriophyllum aquaticum is sold primarily for aquatic gardens, but sometimes also for aquarium use. Since 1996, sale of parrot feather has been banned in Washington because it is an aggressive invader that rapidly takes over lakes and ponds. Parrotfeather is a native of South America that grows well in Pacific Northwest waters. It is distributed throughout much of North America and Hawaii (USDA - NRCS 2004). According to the Washington Department of Ecology all of the parrot feather plants in the United States are female, so no seeds are produced. However, the plant spreads readily through fragmentation of the stems and rhizomes (Ecology 2001).

Myriophyllum sibiricum Komarov

shortspike watermilfoil

Syn: *Myriophyllum exalbescens* Fern.

LCRANS

Origin: Native

Kingdom: Plantae
Division: Magnoliophyta

Though considered native to northern North America and Eurasia, *Myriophyllum sibiricum* may possibly be a circumboreal species that has increased in range (Ecology 2001, Aiken 1981). It is distributed throughout North America except in the southeastern U.S.

***Myriophyllum spicatum* L.**
LCRANS, LIT
Origin: Introduced

spike watermilfoil

Once commonly sold as an aquarium plant, *Myriophyllum spicatum*, is native to Europe and Asia. It was introduced to North America many years ago and is now found over much of the United States (Ecology 2001). *M. spicatum* can be found in lakes, ponds, shallow reservoirs and low energy areas of rivers and streams as well as in the brackish waters of protected tidal creeks and bays. This species is considered a serious pest in waterbodies that have experienced disturbances such as nutrient loading, intense plant management, or abundant motorboat use (Nichols 1994). Milfoil is rapidly spread from lake to lake on boat trailers. Milfoil forms very dense mats of vegetation on the surface of the water interfering with recreational activities such as swimming, fishing, water skiing, and boating and clogging water intakes used in power generation and irrigation (Ecology 2001). The vast, dense mats can rob oxygen from the water by preventing the wind from mixing the oxygenated surface waters to deeper water.

HYDROCHARITACEAE

***Egeria densa* Planch.**
LCRANS, LIT
Origin: Introduced

Brazilian waterweed

Native to South America, *Egeria densa* has also become established in Europe, Japan, Australia and North America (Ecology 2001). For decades *Egeria densa* has been commercially cultivated and sold for use in water gardens and aquariums. Due to its popularity it is now found throughout the United States, apparently dispersed by improper aquarium disposal and cultivated escapees. Populations of this species occurring in North America are staminate therefore no seeds are produced. The primary mode of reproduction is asexual via vegetative fragments. Recreational boating and other activities in infested water bodies contribute to the vegetative dispersal of *Egeria densa* (Les and Mehrhoff 1999). In 1944 *Egeria densa* was found in Oregon (Cohen and Carlton 1995). Officials now consider *Egeria densa* to be one of the greatest threats to Oregon's water bodies. Silver Lake County, in Washington State spends over one million dollars a year to control *Egeria densa* (Ecology 2001). It is also illegal to sell *Egeria densa* in Washington State (Ecology 2001). Not only does *Egeria densa* displace native species, it clogs waterways and impedes navigation (Cohen and Carlton 1995). *Egeria densa* is currently considered a highly invasive species with increasing populations (Les and Mehrhoff 1999).

Kingdom: Plantae
Division: Magnoliophyta

Elodea canadensis Michx. Canadian waterweed, common elodea
Syn: *Anacharis canadensis*, *Elodea brandegeae*, *Elodea ioensis*, *Elodea linearis*, *Elodea planchonii*, *Philotria canadensis*, *Philotria linearis*
LCRANS, LIT
Origin: Native

Origin: Native aquatic plant distributed throughout North America. Because it is a popular aquarium plant it has been widely exported around the world, subsequently introduced and is now considered a noxious weed in parts of Europe, Australia, Africa, Asia, and New Zealand (Ecology 2001). Often confused with *Elodea nuttallii* and *Egeria densa*.

Elodea nuttallii Planch.) St. John western waterweed
Syn: *Anacharis nuttallii* Planch., *Anacharis occidentalis* (Pursh) Victorin, *Elodea columbiana* St. John., *Elodea minor* (Engelm. ex Caspary) Farw., *Elodea occidentalis* (Pursh) St. John, *Philotria angustifolia* (Muhl.) Britt. ex Rydb, *Philotria minor* (Engelm. ex Caspary) Small, *Philotria nuttallii* (Planch.) Rydb., *Philotria occidentalis* (Pursh) House, , *Udora verticillata* var. *minor*
LCRANS, LIT
Origin: Native

Occurs in the Northwest and California, but is more common in the eastern U.S. , *E. nuttallii* can be found in lakes, rivers, ponds and ditches. Unlike *E. canadensis*, *E. nuttallii* prefers fresh to slightly brackish water (Ecology 2001).

Vallisneria Americana Michx tapegrass, water celery
LCRANS, LIT
Origin: Introduced

Vallisneria americana is an aquatic perennial indigenous to eastern North America. The species is now also found in Asia, Australia, Central America and the Caribbean. In the Pacific Northwest *Vallisneria americana* was introduced to provide habitat for wildlife and fish. The species is not an aggressive colonizer and does not cause many of the problems associated with other introduced aquatic plants. Therefore *V. americana* is not considered a pest species (Ecology 2001).

IRIDACEAE

Iris pseudacorus yellow flag iris, water flag
LCRANS, LIT
Origin: Introduced

Kingdom: Plantae
Division: Magnoliophyta

A perennial wetland plant with attractive yellow flowers, *Iris pseudacorus* was brought to and cultivated in eastern North America during the early to mid 19th century. By the 1860's its escape from cultivation was reported. Native to Europe, it is now found throughout the United States and Canada (Cohen and Carlton 1995). Though it is invasive, *I. pseudacorus* is still offered commercially and is widely cultivated. Large, floating seeds are water dispersed. Rhizomes may also be broken off and can float downstream to establish new populations. Due to its competitiveness, *I. pseudacorus* populations are increasing. Once established, native species are displaced and the plant can become a nuisance. Little work has been done on effective removal of yellow flag; glyphosate application is somewhat effective, manual removal may be more effective but may result in highly disturbed habitat.

ISOETACEAE

Isoetes tenella Léman spiny-spore quillwort
Syn: *Isoetes setacea*, *Isoetes muricata*, *Isoetes echinospora*, *Isoetes braunii*
LIT
Origin: Native

Origin: Native distribution: from Newfoundland to British Columbia, south to Pennsylvania and California (Rook 2002).

JUNCACEAE

Juncus balticus Willd. Baltic rush, wire grass
LIT
Origin: Native

Distributed throughout North America (USDA - NRCS 2004).

Juncus effusus L. common rush
LIT
Origin: Native

Distributed throughout North America, may be one or more of four var. possibilities (USDA - NRCS 2004).

Juncus filiformis L. thread rush
LIT
Origin: Native

Distributed throughout the western US and in the Great Lakes region (USDA - NRCS 2004).

Kingdom: Plantae
Division: Magnoliophyta

Juncus nevadensis S. Wats.

Sierra rush

LIT

Origin: Native

Native to the western U. S., there are four varieties found in Oregon and Washington (USDA - NRCS 2004).

Juncus oxymeris Engelm.

pointed rush

LIT

Origin: Native

Origin: Nativespecies with a distribution limited to the west coast of North America (USDA - NRCS 2004).

JUNCAGINACEAE

Triglochin maritimum L.

seaside arrowgrass

Syn: *Triglochin maritima* L., *Triglochin elatum* Nutt.

LIT

Origin: Native

Distributed throughout most of the U.S. except the gulf and mid Atlantic states (USDA - NRCS 2004), also found in Europe and Asia, *Triglochin maritimum* may be a circumpolar species complex.

LAMIACEAE

Mentha aquatica L.

water mint

LIT

Origin: Introduced

Found primarily along the eastern coast from Nova Scotia to South Carolina, but also occurring in most of the inland eastern states and throughout the central and western United States. Water mint is native to Europe and is often sold as a water garden plant. Was probably brought to North America with European immigrants who valued it for its medicinal and herbal uses.

Mentha arvensis L.

wild mint

LIT

Origin: Native

Kingdom: Plantae
Division: Magnoliophyta

This is the only native species of *Mentha* found in the U.S., the rest are all introduced. This plant is very common and used for culinary purposes. Some states such as Nebraska consider this an invasive wetland plant. With a temperate distribution Hickman 1993 considers this to be naturalized from Europe but native tribal records indicated widespread use of this plant (<http://www.wsdot.wa.gov/environment/culres/ethbot/m-p/Mentha.htm>).

Mentha x piperita L. (pro sp.) *aquatica* × *spicata* peppermint
LIT
Origin: Introduced

Origin: Introduced hybrid of two nonnative Eurasian mint species *Mentha aquatica* x *M. spicata*, this plant is popular herb. Peppermint is found throughout much of North America. (USDA - NRCS 2004).

Prunella vulgaris L. common selfheal
LIT
Origin: Native

Prunella vulgaris is native to the continental U.S. but is considered an invasive native in the Northeast and in the Great Plains states (USDA - NRCS 2004).

LEMNACEAE

Lemna minor L. common duckweed
LCRANS, LIT
Origin: Native

Lemna minor is distributed throughout much of the temperate and subtropical regions of the world including North America, Eurasia, Australia, and New Zealand. It may be confused with other duckweeds as well as *Azolla mexicana*. Natural duckweed mats are likely to be a mixture of species.

LILIACEAE

Veratrum californicum Dur. California false hellebore, corn lily
LIT
Origin: Native

Native to North America west of the Rockies, there are two varieties of *Veratrum californicum* found in the Pacific Northwest. Traditional uses of *V. californicum* include its use as a contraceptive, the whole plant should be considered highly toxic (The Compleat Botanica).

Kingdom: Plantae
Division: Magnoliophyta

LYTHRACEAE

Lythrum salicaria L. purple loosestrife, salicaire, spiked loosestrife
LCRANS, LIT
Origin: Introduced

Origin: Introduced throughout much of North America, this species is considered a serious pest, is listed as a nuisance and/or noxious weed in many states, and is banned from sale in most U.S. states (USDA - NRCS 2004). Purple loosestrife disrupts wetland ecosystems by displacing native plants and animals. Economic impacts are high in agricultural communities when irrigation systems are clogged or when wetland pastures are lost to grazing.

MENYANTHACEAE

Nephrhyllidium crista-galli (Menzies ex Hook.) Gilg deercabbage
Syn: *Fauria crista-galli*
LIT
Origin: Native

Native to Oregon and Washington (in the Olympic Mountains and North Cascades) north to British Columbia and Alaska (USDA - NRCS 2004).

NYMPHAEACEAE

Nymphaea odorata* ssp. *odorata American white waterlily
LCRANS, LIT
Origin: Introduced

Nymphaea odorata is native to eastern North America. It consists of two subspecies *N. odorata* ssp. *odorata* and ssp. *tuberosa* (Paine) Wiersema & Hellquist. The two subspecies are widespread in the eastern, central, and mid western United States. *N. a odorata* ssp. *odorata* has been introduced into several western and northwestern states (Weirsema 1997). It is considered a class c nox weed in Washington.

ONAGRACEAE

Epilobium ciliatum Raf. ssp. ***watsonii*** (Barbey) Hoch & Raven fringed willowherb
Syn: *Epilobium adenocaulon*, *Epilobium americanum*, *Epilobium brevistylum*, *Epilobium californicum*, *Epilobium delicatum*, *Epilobium ecomosum*, *Epilobium ursinum*
LIT
Origin: Native

Kingdom: Plantae
Division: Magnoliophyta

Native to the Pacific Northwest and California (USDA - NRCS 2004). *Epilobium ciliatum* has a nearctic distribution.

Ludwigia uruguayensis (Camb.) Hara Uruguayan primrose-willow, water primrose
LIT
Origin: Introduced

Ludwigia uruguayensis is a perennial herb with bright yellow, showy flowers and willow-like leaves that can be found creeping along the shoreline, floating on the water surface, or growing upright. It is a non-native species originally from South America and has been introduced into Europe and northern North America. Water primrose spreads by seeds and by plant fragments. It is easily dispersed by shipping, waterfowl, and human activity. It is also sold as an ornamental species. In Washington water primrose has established in the drainage canals in the Longview/Kelso area. It has been in the area for about 25 years. There is a herbarium specimen dated 1956, from the "Longview Toll Bridge" (Ecology 2001).

There has been some confusion in the past as to the origin of *L. uruguayensis*. Some authors consider this a species complex native to both South America and the Southern U.S. Jennifer Parsons of the Washington Department of Ecology and one of the taxonomic advisors to the LCRANS survey considers this whole complex to be weedy and non-native to the Pacific Northwest.

ORCHIDACEAE

Platanthera dilatata (Pursh) Lindl. ex Beck **var. *dilatata*** scentbottle
LIT
Origin: Native

Native to the northern U.S. and the western states, *Platanthera dilatata* var. *dilatata* is a rare orchid that inhabits soggy soil, bogs, marshes, meadows, fens and prefers full sun (USDA - NRCS 2004).

PLANTAGINACEAE

Plantago lanceolata L. narrowleaf plantain
LIT
Origin: Introduced

An introduced weed, *Plantago lanceolata*, is native to Europe, has been spread throughout the continental U.S., Alaska, Hawaii and Puerto Rico and thrives in many other temperate climates. *P. lanceolata* is commonly found along roadsides, railroads and other disturbed habitats. The leaves of many *Plantago* spp. have medicinal uses and it may have been intentionally transported to North America. The pollen of *P. lanceolata* is also a common allergen.

Kingdom: Plantae
Division: Magnoliophyta

POACEAE

Beckmannia syzigachne (Steud.) Fern. American sloughgrass
Syn: *Beckmannia eruciformis* auct. non; *Beckmannia eruciformis* ssp. *baicalensis*;
Beckmannia eruciformis var. *uniflora*; *Beckmannia syzigachne* ssp. *baicalensis*;
Beckmannia syzigachne var. *uniflora*
LIT
Origin: Cryptogenic

Found in wet meadows, swamps, marshes and shallow water. Range Eastern Europe to central Asia and North America. Most sources consider this to be a native, new-world grass and it is considered threatened and endangered in two midwestern states (Hickman 1993, USDA - NRCS 2004) but the Global Compendium of Weeds (HEAR 2004) lists its origins as China and Asia.

Deschampsia caespitosa (L.) Beauv. tufted
hairgrass
LIT
Origin: Cryptogenic

Distributed throughout the western and northern U.S. Most sources consider this to be a native, new-world grass (Hickman 1993, USDA - NRCS 2004) but the Global Compendium of Weeds (HEAR 2004) lists its origins as Eurasia, Africa, Australia (HEAR 2004).

Distichlis spicata (L.) Greene inland saltgrass
LIT
Origin: Native

Saltgrass is native to North America and is widely distributed (USDA - NRCS 2004). *Distichlis spicata* is the only saltgrass (*Distichlis*) native to the U.S.

Elymus glaucus Buckl. blue wildrye
LIT
Origin: Native

Origin: Nativegrass distributed throughout western North America (USDA - NRCS 2004). Hybridizes readily with other members of the genus *Elymus*.

Glyceria striata (Lam.) A.S. Hitchc. fowl mannagrass
LIT
Origin: Native

Kingdom: Plantae
Division: Magnoliophyta

Origin: Nativegrass, widely distributed throughout North America (USDA - NRCS 2004). Considered invasive in the Czech Republic (Dancak 2002).

Hordeum brachyantherum Nevski meadow barley
LIT
Origin: Native

Native to the western U.S., spotty distribution in the east may indicate that it is introduced to eastern North America (USDA - NRCS 2004).

Lolium arundinaceum (Schreb.) S.J. Darbyshire tall fescue
Syn: *Festuca arundinacea* Schreb. var. *arundinacea* Schreb.
LIT
Origin: Introduced

An agronomically important forage species native to Europe, *Lolium arundinaceum* is considered a pest species in the U.S. where it is widely distributed (USDA - NRCS 2004).

Phalaris arundinacea L. reed canarygrass
LCRANS, LIT
Origin: Introduced

Phalaris arundinacea is a rhizomatous perennial grass (Ecology 2002). Reed canarygrass forms dense, highly productive stands that grow so vigorously they are able to inhibit and eliminate competing species (Apfelbaum and Sams 1987). In addition, areas that have existed as reed canarygrass monocultures for extended periods of time may also be characterized by seed banks that lack any native species (Apfelbaum and Sams 1987, Ecology 2002).

Reed canarygrass is one of the most common species growing along the banks of the lower Columbia River system where it thrives in dense monocultures. Many recent habitat restoration projects along the system are investigating the efficacy of removing reed canarygrass stands.

Reed canarygrass is a circumboreal species (Larson 1993). While possibly native to North America, European cultivars have been widely introduced for use as hay and forage on the continent; there are no easy traits known for differentiating between the native plants and European cultivars (White et al. 1993, Ecology 2002) but it is thought that the invasive populations of reed canary grass are the result of these introduced cultivars. The species is common throughout most of southern Alaska and Canada, as well as all but the southeastern portion of the continental U.S. (Hitchcock et al. 1969).

Spartina spp. cordgrasses
LIT

Kingdom: Plantae
Division: Magnoliophyta

Origin: Introduced*

Several species of cordgrass (*Spartina alterniflora*, *S. anglica*, *S. densiflora*, and *S. patens*) are nonnative, invasive plants in several estuaries along the west coast of North America. As ecological engineers, spreading rapidly by both seeds and rhizomes and forming dense monocultures, they can severely alter the natural hydrology and ecology of invaded habitats (Pfauth et al. 2003). Dense mats of *Spartina* are very effective at trapping sediments and, because of this effect, *Spartina* has, in the past, been intentionally introduced into coastal areas for erosion control. *Spartina* also impacts resident and migratory shorebirds by converting their foraging habitat, the unvegetated, intertidal mudflats, to densely vegetated salt marsh (Pfauth et al. 2003). The growth of *Spartina* is also detrimental to eelgrass beds and the pelagic species that depend on them for food (Pfauth et al. 2003).

While *Spartina* has not been discovered growing in the lower Columbia River system, potentially viable seeds have been found associated with rafts of vegetation stranded along the interior mouth of the estuary (David Jay pers. com).

POLYGONACEAE

Polygonum hydropiperoides Michx.

swamp smartweed

LIT

Origin: Native

Origin: Native range: Western California, from the dry regions of the Great Basin and the Mojave desert to the Pacific coast north to Canada, eastern North America and Mexico. (Hickman 1993).

PONTEDERIACEAE

Eichhornia crassipes (Mart.) Solms

water hyacinth

Syn: *Eichhornia speciosa* Kunth, *Piaropus crassipes* (Mart.) Britton, *Piaropus mesomelas*, *Pontederia crassipes*, *Heteranthera formosa*

LIT

Origin: Introduced*

Origin: Introduced throughout the southern United States and California, *Eichhornia crassipes* is native to South America (Hickman 1993). It is not established in the lower Columbia River basin and, due to colder winter temperatures, probably can't overwinter in the Pacific Northwest. Nevertheless, this popular ornamental pond species has been found in a few Washington sloughs near Longview where it is now believed to have been successfully eradicated (Jennifer Parsons pers comm.). These were either escaped plants or unwanted plants from residential ponds.

Kingdom: Plantae
Division: Magnoliophyta

E. crassipes is an unwanted aquatic plant because its dense mats clog waterways, making boating, fishing and almost all other water activities, impossible while greatly reducing water flow and oxygen levels within the mats. Furthermore water hyacinth greatly reduces biological diversity: mats eliminate native submersed plants by blocking sunlight, alter emerged plant communities by pushing away and crushing them, and also alter animal communities by blocking access to the water and/or eliminating plants the animals depend on for shelter and nesting (IFAS 2004).

POTAMOGETONACEAE

***Potamogeton crispus* L.**
LCRANS, LIT
Origin: Introduced

Curly-leaf pondweed, curly pondweed

A native of Eurasia, *Potamogeton crispus* is now found worldwide. The earliest records of *Potamogeton crispus* in the United States that can be verified date its introduction as sometime in the 1860's. However, there are reports that date the species presence in this country to as early as 1807 (Cohen, Carlton 1995). The first documented appearance of *Potamogeton crispus* in Oregon was in the Rogue River, Curry County, 1947 (Stuckey 1979). The establishment of *Potamogeton crispus* is due to a combination of intentional introductions, careless disposal of aquaria and escapes from cultivation (Les and Mehrhoff 1999). Though, if the species were present as early as the 1807 reports state, this would point to yet another means of introduction. During the early 20th century *Potamogeton crispus* was deliberately planted in marshes for waterfowl forage and aquatic wildlife habitat. Migrating waterfowl may also have a role in dispersing *Potamogeton crispus*. Additionally, activities associated with fish hatcheries and stocking may have transported the species between water bodies. *Potamogeton crispus* also became a popular aquarium and water garden plant during the early 20th century (Les and Mehrhoff 1999). A cold-water species, it can survive the winter in most areas of the United States, which is likely one reason it became popular with water gardeners. The primary form of propagation in *Potamogeton crispus* is by turions, a form of vegetative reproduction. Turions are formed in late spring. Being a cold-water species, *Potamogeton crispus* dies back and goes dormant when water temperatures are high during the summer months. When fall arrives the turions germinate and develop into plants that remain viable throughout the winter. The plants are the most robust during the spring; this is usually when they become a nuisance (Les and Mehrhoff 1999). *Potamogeton crispus* is a highly invasive species with increasing populations.

Citations:

***Potamogeton epihydrus* Raf.**
LCRANS
Origin: Native

ribbonleaf pondweed

***Potamogeton foliosus* Raf.**
LIT

leafy pondweed

Kingdom: Plantae
Division: Magnoliophyta

Origin: Native

Potamogeton friesii Rupr.

Fries' pondweed

LIT

Origin: Native

Potamogeton natans L.

floating pondweed

LCRANS

Origin: Native

Potamogeton pectinatus (L.) Boerner

sago pondweed

LCRANS

Origin: Native

Potamogeton pusillus L.

small pondweed

LCRANS

Origin: Native

Potamogeton richardsonii (Benn.) Rydb.

Richardson's pondweed

LCRANS, LIT

Origin: Native

Potamogeton zosteriformis Fern.

flatstem pondweed

LCRANS

Origin: Native

There are about 80-90 species of *Potamogeton* in the world (IFAS) and perhaps 20 of them occur in the Pacific Northwest (Ecology 2001). Most of them are native species and several of them, such as *P. pectinatus*, are considered invasive species in other parts of the world. They occur in a variety of aquatic habitats. Some pondweeds are totally submersed, others have floating leaves. Although some may vary greatly in size and leaf shape, many *Potamogeton* species are notoriously difficult to tell apart. Pondweeds are very important as wildlife food and some are sold commercially as aquarium or pond plants.

RANUNCULACEAE

Caltha palustris L. var. *palustris*

yellow marsh marigold

LIT

Origin: Native

This marsh marigold is circumboreal in distribution and can be found along the edges of ponds and sloughs in moist soil (Rook 2002). The roots of *Caltha palustris* were commonly used by Native Americans for medicinal purposes.

ROSACEAE

Argentina anserina (L.) Rydb.

Pacific silverweed

Syn: *Argentina argentia*, *Potentilla anserina*

LIT

Kingdom: Plantae
Division: Magnoliophyta

Origin: Native

Native to the coastal dunes, marsh edges and sandy bluffs of the western U.S. from Alaska to Southern California coastal areas, *Argentina anserine* also is sometimes found inland at low elevations.

RUBIACEAE

Galium trifidum ssp. columbianum (Rydb.) Hultén threepetal bedstraw

LIT

Origin: Native

Galium trifidum ssp. columbianum is distributed throughout the western U.S. and parts of the northern states and Canada (USGS- NRCS 2004).

RUPPIACEAE

Ruppia maritima L.- widgeon-grass

LCRANS, LIT

Origin: Cryptogenic

Opportunistic and tolerant to a wide range of environmental conditions, *Ruppia maritima* L. is found worldwide. Typically an inhabitant of marginal seagrass habitats, *Ruppia maritima* L can also be present as a subdominant species, becoming dominate when environmental conditions change. There are indications that *Ruppia maritima* L. becomes dominant in environmentally degraded areas and under unfavorable climatic conditions (Johnson et al. 2003).

SALICACEAE

Salix hookeriana Barratt ex Hook. dune willow, coastal willow

LIT

Origin: Native

Native to the western coast of North America the coastal willow is found from Northern California to Alaska (Hickman 1993).

SCROPHULARIACEAE

Castilleja ambigua Hook. & Arn. ssp. ***ambigua*** johnny-nip, Indian paintbrush, owl
clover, purple owl's clover

Syn: *Orthocarpus exsertus*, *Orthocarpus purpurascens*

LIT

Kingdom: Plantae
Division: Magnoliophyta

waterways and interfere with recreation in shallow waters. The 6-foot-long, green reed has a small yellow flower that contains a bur-like fruit. The plant was not known to exist in the United States before the Dutch shipment arrived. State and federal agriculture inspectors scrambled to recover as many plants as possible but it is not known if attempts to collect all species were successful. Washington and Oregon are two states where shipments of the contaminated plants are believed to have been shipped.

Sparganium angustifolium Michx. narrowleaf bur-reed
LCRANS
Native

Narrow-leaf burr reed is native to the Western US, Alaska and can be found throughout the Great Lakes region. *Sparganium* is fodder for waterfowl, muskrats and deer. Stem base and tubers are edible (Ecology 2001).

TYPHACEAE

Typha angustifolia L.- Narrowleaf / Narrowed-leaved Cattail, Nail Rod
LIT, LCRANS
Origin: Introduced

Endemic to Eurasia, *Typha angustifolia* is now found in South America and throughout North America. The presence of *Typha angustifolia* on the eastern coast of the United States was reported in the 1820s (Cohen, Carlton 1995). The species was possibly introduced to the Atlantic coast by dry ship ballast. *Typha angustifolia* was apparently used for matting and pillow stuffing. Parts of the plant were also eaten. These uses may have facilitated the dispersal of the species (Mills et. al.1993). *Typha angustifolia* is a perennial plant that is invasive and capable of spreading rapidly.

Typha latifolia L. broadleaf cattail
LIT
Origin: Native

Origin: Nativespecies, distributed widely throughout North America and in temperate parts of Central America, Eurasia, and Africa (Hickman 1993).

ZANNICHELLIACEAE

Zannichellia palustris L. horned pondweed
LCRANS, LIT
Origin: Native

Kingdom: Plantae
Division: Magnoliophyta

A delicate underwater branching perennial this plant has a more or less worldwide distribution and is common throughout North America. In Washington, horned pondweed is common in hard water lakes of the Columbia Basin (Ecology 2001). It may be confused with *Ruppia maritime*.

ZOSTERACEAE

Zostera japonica Aschers. and Graebn Japanese eelgrass, dwarf eelgrass
LCRANS, LIT
Origin: Introduced

Native to Japan, *Zostera japonica* is now established on the coast of the Pacific Northwest. The first recorded collection of the species on the Pacific coast was from Washington State in 1957. *Zostera japonica* has been observed to be abundant in several areas of the Pacific Northwest coast that have been or are presently used for intensive oyster cultivation. It has been suggested that *Zostera japonica* was possibly used as packing material when oyster spat was shipped from Japan to oyster farms in the PNW. Being an annual plant, *Zostera japonica* is a prolific seed producer. Seeds may now be the primary mode of dispersal for this species (Harrison and Bigley 1982).

Zostera marina common eelgrass
LIT
Origin: Native

Zostera marina is widespread throughout the Atlantic and Pacific. In the eastern Atlantic it extends from the Arctic Circle to Gibraltar, including the Mediterranean. *Z. marina* forms large colonies on muddy substrates especially in estuaries, and also occurs on sandy substrates where there is weak wave action.

Kingdom: Animalia
Phylum: Porifera

Kingdom: Animalia

Phylum: Porifera

The sponges

The identification of freshwater sponges depends on characteristics of spicules and on features of intact gemmules. Species identifications depend absolutely on obtaining all types of the spicules (megascleres, gemmoscleres and, if present, microscleres) (Penny and Racek 1968; Thorp and Covich 2001). Gemmoscleres are particularly important but they may occur only during certain times of the year (Thorp and Covich 2001:115). Spicule preparations require digestion of the tissue in nitric acid in a tube immersed in boiling water for 1 hour, followed by centrifugation. The acid is then poured off and the spicules are washed in ethanol (Penny and Racek 1968, Thorp and Covich 2001).

The procedures necessary for preliminary identification were beyond the scope of this investigation although one sponge, a forest-green specimen was collected from a freshwater site at Sauvies Island, Oregon.

Phylum: Ciliophora

Class: Ciliatea

Ciliates

Protozoans are often overlooked but play a major role in nutrient cycling (Taylor and Sanders 2001).

Didiniidae		
<i>Mesodinium rubrum</i>	LCRANS	Native

DIDINIIDAE

Mesodinium rubrum (Lohmann, 1908)

Synonyms: *Cyclotrichium meunieri*, *Halteria rubra*, *Myrionecta rubra*

Origin: Native

LCRANS

Collected from Ilwaco Harbor and Young's Bay in October 2002 during a red tide, this is a solitary, bloom-forming, obligate autotroph (Lindholm 1985). This species contains a commensal photosynthetic alga (an endosymbiotic cryptophyte chloroplast) and is nontoxic. Identified by Dr. Rita Horner and Dr. Jin Wan Lee, it is probably a complex of

Kingdom: Animalia
Phylum: Cnidaria

closely related species. Dr. Horner relates that it is common in the northeast Pacific and considers it a native species. The unexplained global distribution of *Mesodinium rubrum* could result from it being a complex of closely similar geographically isolated species or from widespread introductions of one or more of its populations.

Phylum: Cnidaria

Class: Anthozoa

Edwardsiidae		
<i>Nematostella vectensis</i>	LCRANS, LIT	Introduced

EDWARDSIIDAE

Nematostella vectensis Stephenson, 1935

Syn: *Nematostella pellucida*

LCRANS, LIT

Origin: Introduced

Fifteen *Nematostella vectensis* were collected alive from muddy sand habitats and a shallow pool of a high *Carex* salt marsh in the lower Columbia River. J. T. Carlton (in correspondence) suggests that this species may have a trans-Arctic distribution i.e. ranging south from the Arctic on northern coastlines of the northern hemisphere to northern Japan, Puget Sound, Cape Cod, and the Bay of Biscay. Hand and Uhlinger (1991) demonstrated that the low latitude populations are a single species by interbreeding females from England, Maryland, Georgia, California, Oregon and Washington with males from Nova Scotia, Maryland, Georgia and Oregon in a total of 24 crosses which all produced healthy first and second generations. The global distribution of *N. vectensis* therefore appears unlikely to be of natural processes. The lack of large-scale genetic patterns among populations in different lagoons of Great Britain is consistent with occasional passive or anthropogenic dispersal of low number of individuals between lagoons (Pearson et al. 2002). Natural occurrences of the isolated British *Nematostella* populations therefore would be difficult to explain. More likely, the British populations are introduced.

Kozloff (1983) concludes that northeast Pacific *N. vectensis* are an Atlantic species for which “the exact date of introduction into our region is unknown” while Carlton (2000) lists *N. vectensis* as “cryptogenic” in Coos Bay. Confusion over the origins of the northeast Pacific *Nematostella* may partly result from poor information the likely expansion of its populations since the early 1900s and its occurrence only from San Francisco Bay north, a relatively narrow range if this were a native northeast Pacific species. Hand (1957) reported “This anemone probably is the ‘will-of-the-wisp’ species that I have hunted for more than 10 years in California. In 1946, the late Prof. S. F. Light described to me a very small anemone he had seen in small pools on the Salicornia marshes of Richardson’s Bay (a part of San Francisco Bay).” Since 1957, published reports of northeast Pacific, *Nematostella* are only from Puget Sound, Washington, Coos

Kingdom: Animalia
Phylum: Cnidaria

Bay, Oregon, Tomales Bay, California, and San Francisco Bay, California (Kozloff 1983, Hand & Uhlinger 1994). However, Jeff Cordell has found *N. vectensis* in almost every salt marsh of Oregon and Washington he has sampled in the last 20 years. In a 1994 survey of Trestle Bay in the lower Columbia River, prior to the breaching of the jetty, densities of *N. vectensis* were reported as 2,715/m² but two years after the breach no cnidaria were found (Hinton and Emmett 2000).

The geographical and climatic range of *Nematostella vectensis* on the eastern North American coast, from Nova Scotia to Georgia and western Florida to Louisiana (Hand & Uhlinger 1994) is much broader than the European or eastern Pacific ranges. Nova Scotia is colder and the Gulf of Mexico is warmer than temperatures of southern and eastern Britain and the eastern Pacific coast between Puget Sound and San Francisco. *N. vectensis* would therefore require pre-adapted thermal tolerances of occur western Atlantic and Gulf of Mexico range if it is native to the eastern Pacific or Europe. Therefore *N. vectensis* is more likely to be the native to the western Atlantic and Gulf of Mexico.

Hand and Uhlinger (1994) considered ballast water transport to be the most likely mechanism for dispersing *Nematostella* since it does not occur on hard substratums and is unlikely to be introduced with transplanted oysters, on ship hulls or in the fouling faunas associated with other hard substratums. The asexual reproduction of this species Hand and Uhlinger (1992) allows it to colonize new habitats with very few original propagules. Moreover, well-fed individuals can grow to 16 cm in length and individuals can survive 6 months of starvation (Hand and Uhlinger 1992). The extreme durability of this species and its close association with high intertidal sediments suggest that it could have been introduced to Britain and to the western United State in ballast sediments of early sailing ships. Many records of England indicate the regular use of ballast on board sailing vessels of the North Atlantic trade (Prowse 1895) and Dana (1840) reported sailing ship ballast dumped from the eastern United States directly into San Francisco Bay. Moreover Lindroth (1957) elegantly established the faunal connections between eastern and western North America and Great Britain via ballast sediments of sailing ships.

Class: Hydrozoa

Clavidae		
<i>Cordylophora lacustris</i>	LCRANS, LIT	Introduced
Hydridae		
<i>Hydra spp.</i>	LCRANS	

CLAVIDAE

Cordylophora lacustris Agassiz, 1862

Syn: *Cordylophora caspia*

LCRANS, LIT

Origin: Introduced

Kingdom: Animalia
Phylum: Cnidaria

Cordylophora lacustris is probably native to the Caspian Sea and the Black Sea. The first report of eastern Pacific *C. lacustris* is based on specimens collected in the lower Columbia River near Astoria, Oregon from pilings and posts in low salinity or fresh water in 1965 (Haertel and Osterberg 1967). However, Carlton (1979) found specimens collected from Lake Union, Washington in 1920 and (Cohen and Carlton 1995) found specimens from San Francisco Bay, California collected around 1930. *Cordylophora lacustris* was likely spread world wide prior to the 20th century in association with ship fouling and ballast water (Carlton 1979, Cohen and Carlton 1995).

Phylum: Ectoprocta
Class: Phylactolaemata

Ectoprocts were commonly lumped together with the entoprocts and referred to generally as “Bryozoa” (Thorp & Covich 2001). The class Phylactolaemata is an exclusively freshwater colonial group of ectoprocts. Adult stages attach to submerged surfaces such as branches, rocks and logs. The phylactolaemates form statoblasts dormant seed-like buds that are resistant to dessication and can remain dormant for long periods. The statoblasts are a likely life history stage for natural or anthropogenic transport between water bodies. The distributions of ectoprocts across North America are poorly known. Few large area surveys of bryozoa have been conducted in northwestern North America (see Wood 2001). Marsh and Wood (2002) were the first to survey freshwater bryozoans of the Pacific Northwest and records from outside of northeastern North America are few (Marsh and Wood 2002).

Fredericellidae		
<i>Fredericella browni</i>	LIT	Cryptogenic
<i>Fredericella indica</i>	LCRANS, LIT	Introduced
Pectinatellidae		
<i>Pectinatella magnifica</i>	LCRANS, LIT	Introduced
<i>Plumatella emarginata</i>	LIT	Cryptogenic
<i>Plumatella vaihirieae</i>	LIT	Cryptogenic

FREDERICELLIDAE

Fredericella browni Rogick, 1945

LIT

Origin: Cryptogenic

Collected from the Willamette River below the Oregon City Falls (Marsh and Woods 2002) and at three other Pacific Northwest sites. This is not a common species in Northeastern and Central United States where most bryozoan surveys have taken place (Marsh and Woods 2002). Specimen have also been reported in India (Pachut 1998).

Kingdom: Animalia
Phylum: Ectoprocta

Fredericella indica (Annandale, 1909)

LCRANS, LIT

Origin: Introduced

This species is common throughout North America especially in eastern states, at scattered sites in Europe, Africa, and Asia, and probably includes several species not yet distinguished (Thorp & Covich 2001). Distribution data for both U.S. states and Canadian provinces is likely incomplete. A month-long collection trip of bryozoans and sponges in the Pacific Northwest encountered this species at only four widely dispersed localities (Marsh and Wood 2002). While the origin of this species remains uncertain (likely eastern North America where it is very common) we consider *F. indica*, which is widespread in the lower bays of the basin (in brackish as well as freshwater), to be introduced into the lower Columbia River. Further surveys may reveal less disjoint distributions, however.

PECTINATELLIDAE

Pectinatella magnifica (Leidy, 1851)

Syn: *Fredericella magnifica*

LCRANS, LIT

Origin: Introduced

The gelatinous masses of *Pectinatella magnifica* form gelatinous colonies on submerged wood of any kind including docks Smith (2003). Massive colonies may exceed 60 cm in diameter, however colony sizes of less than 10 cm may go unnoticed for long periods until residents are "shocked" by its sudden appearance when ecological conditions favor massive "alien-like" colonies (Smith 2001). *P. magnifica*, is widely distributed east of the Mississippi River and is likely to be endemic to eastern and central North America (Smith 2001). Marsh and Wood (2002) found *P. magnifica* throughout Oregon including the Columbia River. The first records of *P. magnifica* in the lower Columbia River are from the late 1990s (see EMAP 2001 and Marsh and Wood 2002). Previously, *P. magnifica* had only been recorded from as far west as eastern Texas. *Pectinatella magnifica* has been introduced to Japan, Korea, India, and Europe (Smith 2001). The first records of *P. magnifica* in the lower Columbia River are from the late 1990s (see EMAP 2001 and Marsh and Wood 2002). The anchor-spiked statoblasts of *P. magnifica* are highly adapted for hooking onto fur and feathers for dispersal on birds and mammals between isolated water bodies in regions where it occurs.

Plumatella emarginata Allman 1844

LIT

Origin: Cryptogenic

Occurring in North America, Great Britain, India, Australia and Japan; it is cosmopolitan in northern hemisphere and may be endemic to Europe (Wood 2001).

Kingdom: Animalia
Phylum: Ectoprocta

Plumatella vaihiria (Hastings 1929)

Syn: *Hyalinella vaihiria*

LIT

Origin: Cryptogenic – probably introduced

Previously known only from four sites in North America three of which are wastewater treatment plants, Marsh and Wood (2002) collected *Plumatella vaihiria* from Oaks Bottom Slough (off of the Willamette River) in 1998. *Plumatella vaihiria* is a nuisance fouling organism (Wood and Marsh 1999). The type locality of *P. vaihiria* is a high mountain pond in Tahiti and it is known also from Hawaii and Argentina (Wood and Marsh 1999). An unconfirmed report of *P. vaihiria* is from Australia (Wood and Marsh 1999). Unlike *Pectinatella magnifica* *P. vaihiria* is characterized by rapid growth and massive colonies (Wood and Marsh 1999, Marsh and Wood 2002). Given its wide geographic range and limited literature citations this species is likely an invader but not enough information exists to confirm this.

Phylum: Entoprocta

The Entoprocta are a small group of species (~ 60 in all) that are distinct from the Ectoprocta but often lumped with them and referred to together as “Bryozoa.” *Urnatella* is the only freshwater genus in the phylum. Little is known about the distribution of entoprocta in North America as only a few large area surveys of bryozoa have been conducted (see Wood 2001) and most records from outside of northeastern North America only report relatively few species from a limited number of localities (Marsh and Wood 2002).

Urnatellidae			
<i>Urnatella gracilis</i>	LIT	Cryptogenic	

URNATELLIDAE

Urnatella gracilis Leidy 1851

LIT

Origin: Cryptogenic

Considered by Thorp and Covich (1991) to be the most common and widely distributed of the *Urnatella*, *Urnatella gracilis* is the only species of the genus reported from North America where its distribution ranges from the east to west coast and from Texas to Michigan. *U. gracilis* has a true cosmopolitan distribution as it is found on every continent but Antarctica and Australia (Thorp and Covich 1991).

Kingdom: Animalia
Phylum: Nemertea

Phylum: Nemertea

Class: Enopla

Nemertean identifications were not conducted by LCRANS.

Emplectonematidae			
<i>Paranemertes californica</i>	LIT		Native

EMPLECTONEMATIDAE

Paranemertes californica Coe 1904

LIT

Origin: Native

Native to littoral and benthic sites in the Pacific. Reported by EMAP 1999 and EMAP 2000 collections.

Phylum: Annelida

SubClass: Oligochaeta

Very few macroinvertebrates are more poorly studied in the lower Columbia River than the oligochaetes. Few prior studies on the lower Columbia conducted oligochaete identifications, only noting the presence of oligochaetes when encountered. There are several reasons for this. Oligochaete taxonomy is widely regarded as a difficult field and expert identifications may be beyond the scope of many projects. In addition, traditional sorting and preserving techniques used for benthic samples often damage worms beyond identification. Very little is known about native origins and transport of many species, the majority of species are simply labeled as having cosmopolitan or near cosmopolitan distributions

In the lower Columbia River special interest was paid to proper oligochaetes collection and preservation. In 2003 oligochaete samples were identified by Dr. Steve Fend. *Teneridrilus columbiensis* (a species named after its collection location – the Columbia River) was not found in the course of our sampling. Furthermore, some species limited to specific habitats (like banks or sandy weed beds) may not have been found at multiple stations because few such habitats were sampled overall. Of the seven native species collected, only three were found at nine or more stations (out of 45 possible stations) further indicating that collection efficiency was low and more comprehensive collection efforts should be undertaken.

Introduction mechanisms for oligochaetes are varied. Ballast water is a likely vector for many species, others may arrive in new habitats associated with sediments of nonnative ornamental aquatic plants or semi-aquatic plants.

Kingdom: Animalia
 Phylum: Annelida

While oligochaetes are considered freshwater organisms but species such as *Tubifex tubifex* and *Limnodrilus hoffmeisteri* can withstand exposures of up to 10 ppt (Brinkhurst and Gelder 2001). Most others can only survive exposures of 5 ppt or less. However, recent studies have shown that low salinity water may improve the ability of oligochaetes to withstand stress (Brinkhurst and Gelder 2001).

Enchytraeidae		
<i>Enchytraeus spp.</i>	LIT	
Lumbriculidae		
<i>Eclipidrilus n. sp.</i>	LCRANS, LIT	Native
<i>Kincaidiana hexatheca</i>	LCRANS	Native
<i>Rhynchelmis sp.</i>	LCRANS	
<i>Stylodrilus heringianus</i>	LCRANS	Introduced
Naididae		
<i>Amphichaeta sannio</i>	LCRANS	Cryptogenic
<i>Arcteonais lomondi</i>	LCRANS	Cryptogenic
<i>Chaetogaster diaphanous</i>	LCRANS	Introduced
<i>Chaetogaster nr. diastrophus</i>	LCRANS	Cryptogenic
<i>Dero digitata</i>	LCRANS	Cryptogenic
<i>Nais cf. elinguis</i>	LCRANS	Cryptogenic
<i>Nais cf. simplex</i>	LCRANS	Cryptogenic
<i>Nais communis</i>	LCRANS	Cryptogenic
<i>Nais pardalis</i>	LCRANS	Cryptogenic
<i>Nais variabilis</i>	LCRANS	Cryptogenic
<i>Ophidonais serpentina</i>	LCRANS	Cryptogenic
<i>Paranais frici</i>	LCRANS	Introduced
<i>Paranais litoralis</i>	LCRANS	Cryptogenic
<i>Pristina aequiseta</i>	LCRANS	Cryptogenic
<i>Pristina osborni</i>	LCRANS, LIT	Cryptogenic
<i>Slavina appendiculata</i>	LCRANS	Cryptogenic
<i>Stylaria lacustris</i>	LCRANS	Cryptogenic
Ocerodrilidae		
<i>Eukerria saltensis</i>	LCRANS	Introduced
Tubificidae		
<i>Aulodrilus pluriseta</i>	LCRANS	Cryptogenic
<i>Bothrioneurum vej dovskyanum</i>	LCRANS, LIT	Cryptogenic
<i>Branchiura sowerbyi</i>	LCRANS, LIT	Introduced
<i>Ilyodrilus frantzi</i>	LCRANS, LIT	Native
<i>Ilyodrilus templetoni</i>	LCRANS, LIT	Cryptogenic
<i>Limnodrilus hoffmeisteri</i>	LCRANS, LIT	Cryptogenic
<i>Limnodrilus silvani</i>	LIT	Cryptogenic
<i>?Limnodrilus udekemianus</i>	LCRANS, LIT	Cryptogenic
<i>Rhyacodrilus coccineus</i>	LCRANS, LIT	Cryptogenic
<i>Rhyacodrilus spp.</i>	LIT	

Kingdom: Animalia
Phylum: Annelida

<i>Spirosperma nikolskyi</i>	LCRANS, LIT	Cryptogenic
<i>Spirosperma spp.</i>	LIT	
<i>Tasserkidrilus harmani</i>	LCRANS, LIT	Cryptogenic
<i>Telmatodrilus vejovsky</i>	LIT	Cryptogenic
<i>Teneridrilus columbiensis</i>	LIT	Native
<i>Teneridrilus mastix</i>	LCRANS, LIT	Native
<i>Teneridrilus cf. calvus</i>	LCRANS, LIT	Native
<i>Tubifex tubifex</i>	LCRANS, LIT	Cryptogenic
<i>Tubificidae sp 1</i>	LIT	
<i>Tubificidae sp 2</i>	LIT	
<i>Varichaetadrilus pacificus</i>	LCRANS, LIT	Native

LUMBRICULIDAE

Eclipidrilus n. sp.

LCRANS, LIT

Origin: Native

First collected from the lower Columbia River during Miller Sands examination (Date) not enough specimens were collected to make ID. LCRANS collected this species from Miller Sands as well as other sites in Cathlamet Bay. Further collections need to be conducted to gather more type specimens.

Kincaidiana hexatheca Altman, 1936

LCRANS

Origin: Native

Identified and labeled as native to northwestern North America by Steve Fend.

Stylo-drilus heringianus Claparede, 1862

LCRANS

Origin: Introduced

This is a holarctic freshwater species whose status as an invasive species has been debated for many years (See Brinkhurst 1968, 1976). Likely native to Europe, this species has also been collected from places as diverse as Japan, Malaysia, and Egypt indicating that transport and introduction of *Stylo-drilus heringianus* is certainly possible.

NAIDIDAE

Amphichaeta sannio Kallstenius 1892

LCRANS

Origin: Cryptogenic – probably introduced

Kingdom: Animalia
Phylum: Annelida

Considered a European estuarine species by some (not reported from North America prior to the publication of Kathman and Brinkhurst 1998). Possibly synonymous with *A. raptisae*. Steve Fend considers this of unknown origin with a holarctic distribution.

Arcteonais lomondi (Martin 1907)

LCRANS

Origin: Cryptogenic

Widespread (Kathman and Brinkhurst 1998), unknown origin, holarctic distribution.

Chaetogaster diaphanus (Gruithuisen, 1828)

Syn: *Nais diaphana*, *Chaetogaster diaphanus cyclops*

LCRANS

Origin: Introduced

A freshwater species with a holartic distribution (S. Fend personal communication), *Chaetogaster diaphanous* is considered by the California Department of Fish and Game to be an introduced species (CDFG 2002). During their survey of the California coastal and estuarine waters this species was found only in the Sacramento San Joaquin Delta region of the San Francisco Bay (CDFG 2002). In the lower Columbia River a single specimen of *C. diaphanous* was found at a station located at the mouth of the Columbia Slough in Portland. Although its native range is unknown, the scattered and rare distribution of this species along the West Coast likely indicates that it is indeed nonnative to this region.

Chaetogaster nr. diastrophus (Gruithuisen 1828)

syn: *Pseudochaetogaster longmeri*, *C. langi*

LCRANS

Origin: Cryptogenic

Widespread (Kathman and Brinkhurst 1998), near cosmopolitan, possibly holartic in origin.

Dero digitata (Muller 1773)

Syn: *Nais digitata*

LCRANS

Origin: Cryptogenic – probably introduced

Widespread (Kathman and Brinkhurst 1998), near cosmopolitan, probably tropical in origin.

Nais cf. elinguis Muller 1773

LCRANS

Origin: Cryptogenic

Kingdom: Animalia
Phylum: Annelida

Widespread (Kathman and Brinkhurst 1998), near cosmopolitan, possibly holartic in origin.

Nais cf. simplex Piguet 1906

LCRANS

Origin: Cryptogenic

Widespread east of the Mississippi, also known from British Columbia (Kathman and Brinkhurst 1998), near cosmopolitan, possibly holartic in origin.

Nais communis Piguet 1906

LCRANS

Origin: Cryptogenic

Widespread (Kathman and Brinkhurst 1998). *Nais communis* and *N. variabilis* features often overlap, complex needs revision overall.

Nais pardalis Piguet 1906

LCRANS

Origin: Cryptogenic

Widespread, previously known as a variant of *N. bretscheri*, often confused with *N. variabilis* (Kathman and Brinkhurst 1998). Near cosmopolitan distribution, possibly holartic in origin.

Nais variabilis Piguet 1906

LCRANS

Origin: Cryptogenic

Widespread (Kathman and Brinkhurst 1998). *Nais communis* and *N. variabilis* features often overlap, complex needs revision overall.

Ophidonais serpentina (Muller 1773)

Syn: *Nais serpentina*

LCRANS

Origin: Cryptogenic

Widespread (Kathman and Brinkhurst 1998), unknown origin, near cosmopolitan distribution.

Paranais frici Hrabe, 1941

Syn: *Wapsa mobilis*?

LCRANS

Origin: Introduced

Kingdom: Animalia
Phylum: Annelida

Near cosmopolitan distribution. It is most often found in coastal waters, but usually in freshwater. This is a brackish water genus (likely originated in the Tethys) (Timm 1980). Considered introduced in San Francisco Bay and parts of Southern California by Brinkhurst and Cook (1980) and Cohen and Carlton (1995). Two specimens were collected in the lower Columbia River in a grab sample taken at the Sportsmen's Club boat launch in Kalama, WA. Species is present in Kozloff (1987) and probably established in the Pacific Northwest, but requires further confirmation. Considered introduced in the Baltic and in the Great Lakes. Timm (1980) considers it recently introduced to North America.

Paranais litoralis (Muller 1784)

Syn: *Nais litoralis*

LCRANS

Origin: Cryptogenic

Widespread, mostly coastal in tidal fresh or brackish water (Kathman and Brinkhurst 1998) but of unknown origin.

Pristina aequiseta Bourne 1891

Syn: *P. foreli* and *P. evelinae*

LCRANS

Origin: Cryptogenic

Widespread (Kathman and Brinkhurst 1998).

Pristina osborni (Walton 1906)

Syn: *Naidium minutum*, *Naidium osborni* *Pristina minutum*

LCRANS, LIT

Origin: Cryptogenic

Member of a "group of taxonomically problematic species" (Collado and Schmelz 2002). Kathman and Brinkhurst (1998) report it from Illinois, the east coast of North America and Argentina.

Slavina appendiculata (d'Udekem 1855)

Syn: *Nais appendiculata*, *Nais gracilis*

LCRANS

Origin: Cryptogenic

Widespread (Kathman and Brinkhurst 1998), near cosmopolitan, unknown origin.

Stylaria lacustris (Linnaeus 1767)

Syn: *Nereis lacustris*, *Nereis proboscidea*

LCRANS

Origin: Cryptogenic

Kingdom: Animalia
Phylum: Annelida

Widespread (Kathman and Brinkhurst 1998), holarctic and African distribution, possibly holartic in origin.

OCNERODRILIDAE

Eukerria saltensis (Beddard, 1895)

LCRANS

Origin: Introduced

Native to South America, this worm is considered an invasive pest species in Australia where severe infestations can damage rice crops (see http://www.ricecrc.org/reader/Oligochaeta_aquatic_earthworms.htm). It is not an obligate aquatic species, and can survive in irrigated pastures. It is considered a tropical species with a near cosmopolitan distribution.

TUBIFICIDAE

*Aulodrilus pluriset*a (Piguet 1906)

Syn: *Naidium pluriset*a

LCRANS

Origin: Cryptogenic

The genus *Aulodrilus* is currently being rewritten to clear up misidentifications especially *A. pluriset*a and *A. japonica* (Kathman and Brinkhurst 1998). A widespread species, most North American *A. pluriset*a may actually be *A. japonica*.

Bothrioneurum vej dovskyanum Stolc 1886

LCRANS, LIT

Origin: Cryptogenic

Widespread in North America, especially in sandy situations, may be synonymous with *B. americanum*.

Branchiura sowerbyi Beddard, 1892

LCRANS, LIT

Origin: Introduced

Native to tropical and sub-tropical Asia, *Branchiura sowerbyi*, is a widely introduced oligochaete. This tubificid worm may have originally been spread around the world in the water and sediments associated with ornamental aquatic plants such as water-lilies (Cohen and Carlton 1995). Often only conspicuous in artificially warm water (where it grows to a large size) *B. sowerbyi* can be found at locations scattered throughout North America (Brinkhurst 1986). The first record of this species in North America came from the Ohio River in 1930 (Spencer 1932). *B. sowerbyi* was discovered in San Francisco

Kingdom: Animalia
Phylum: Annelida

Bay in 1950 and the Bay had the only recorded west coast population until now (Cohen and Carlton 1995, NAS 2003). However, as only three specimens were found as a single sampling station on the lower Columbia River (in Crane Lake on Sauvie Island – note a shallow warm lake), we are uncertain as to how widespread or established this population is. In addition, fragments of *B. sowerbyi* may be erroneously identified as *Aulodrilus plurisetus* (Brinkhurst 1986)

Ilyodrilus frantzi Brinkhurst 1965
LCRANS, LIT
Origin: Native

Distributed throughout western North America.

Ilyodrilus templetoni (Southern 1904)
Syn: *Tubifex templetoni*
LCRANS, LIT
Origin: Cryptogenic

Widespread and common. Similar to *Tubifex tubifex* (Kathman and Brinkhurst 1998).

Limnodrilus hoffmeisteri Claparede 1862
LCRANS, LIT
Origin: Cryptogenic

Native to North America *Limnodrilus hoffmeisteri* is considered a pollution indicator species. *L. hoffmeisteri* can also inhabit brackish waters to 10 ppt (Brinkhurst and Gelder 2001).

Limnodrilus silvani Eisen, 1879
LIT
Origin: Cryptogenic

Limnodrilus udekemianus Claparede 1862
LCRANS, LIT
Origin: Cryptogenic – possibly native

May be native to North America but has a cosmopolitan distribution

Rhyacodrilus coccineus (Vejdovsky 1875)
Syn: *Tubifex coccineus*
LCRANS, LIT
Origin: Cryptogenic

Widespread North American distribution (Kathman and Brinkhurst 1998), the origin of this species is unclear and complicated by its cosmopolitan distribution.

Kingdom: Animalia
Phylum: Annelida

Spirosperma nikolskyi (Lastochkin and Sokolskaya 1935)
syn: *S. variegatus*, *S. oregonensis*, *Pelsoscolex oregonensi*,
LCRANS, LIT
Origin: Cryptogenic

Widespread, this genus may need more taxonomic work (Kathman and Brinkhurst 1998). Of unknown origin this species is distributed throughout Asia and North America.

Tasserkidrilus harmani (Loden 1979)
Syn: *Tubifex harmani*
LCRANS, LIT
Origin: Cryptogenic

This species is reported as widely distributed species throughout the North America but this is based on prior observations that were not made using all the accepted characteristics (Kathman and Brinkhurst 1998). It is a nearctic species with widely scattered records. It is probably native to North America.

Telmatodrilus vejdosky Eisen 1879
LIT
Origin: Cryptogenic

Teneridrilus columbiensis (Brinkhurst and Diaz 1985)
Syn: *Isochaetides columbiensis*
LIT
Origin: Native

Type specimen collected in the lower Columbia River at Miller Sands. Not known from any other locations (Brinkhurst and Diaz 1985, Erseus et al. 1990).

Teneridrilus mastix (Brinkhurst 1978)
Syn: *Ilyodrilus mastix*
LCRANS, LIT
Origin: Native

Collected from the Fraser River, British Columbia; Columbia River, Oregon; San Francisco Bay, California; and Pearl River, China (Brinkhurst 1986, Erseus et al 1990). Carlton and Geller (1993) list *T. mastix* as a nonnative species introduced via ballast water from China. The California Department of Fish and Game (2002) lists the same species as cryptogenic but identify its origin as Asia. Although some controversy exists as to the origin of this species we do not believe that enough information exists to contradict the original description of the species as native to western North America.

Teneridrilus cf. calvus Erseus and Brinkhurst 1990
LCRANS, LIT
Origin: Native

Kingdom: Animalia
Phylum: Annelida

Type specimen from Sacramento-San Joaquin Delta, California collected in freshwater muddy sediments (Erseus et al. 1990).

Tubifex tubifex (Muller 1774)

Syn: *Lumbricus tubifex*

LCRANS, LIT

Origin: Cryptogenic

Widespread but not as common as general texts suggest, this species occurs in marginal habitats (oligotrophic or hyereutrophic) and cold climates perhaps because it can avoid competition at such extremes (Kathman and Brinkhurst 1998). This is likely a complex with multiple variants. Susceptible to parasite infections such as whirling disease (*Myxobolus cerebralis*). *Tubifex tubifex*, like *Limnodrilus hoffmeisteri*, can withstand prolonged exposure to salinities up to 10ppt (Brinkhurst and Gelder 2001).

Varichaetadrilus pacificus (Brinkhurst 1981)

LCRANS, LIT

Origin: Native

Unlike many oligochaetes of the family Tubificidae, *Varichaetadrilus pacificus* is contaminant intolerant (Canfield et al., 1994). This species is native to North America.

Phylum: Annelida

Subclass: Polychaeta

Older polychaete keys specific to the Pacific Northwest are considered to be full of errors and thus the taxonomic certainty of polychaetes found during the literature review is uncertain. Polychaete taxonomy on a world-wide basis is in a state of flux and disagreements between experts on identifications, origins and distribution complicate the process of identifying introduced polychaetes in the lower Columbia River.

Errant Polychaetes

Glyceridae		
<i>Glycera americana</i>	LIT	Cryptogenic
<i>Glycera macrobranchia</i>	LIT	Native
<i>Glycera nana</i>	LIT	Cryptogenic
<i>Glycera tenuis</i>	LIT	Native
<i>Hemipodus borealis</i>	LIT	Native
Goniadidae		
<i>Glycinde armigera</i>	LIT	Native
<i>Glycinde picta</i>	LIT	Native
<i>Glycinde polygnatha</i>	LIT	Cryptogenic
Hesionidae		

Kingdom: Animalia
 Phylum: Annelida

<i>Hesionella mccullochae</i>	LIT	Native
<i>Podarkeopsis brevipalpa</i>	LIT	Cryptogenic
Nephtyidae		
<i>Nephtys caecoides</i>	LIT	Native
<i>Nephtys californiensis</i>	LIT	Native
<i>Nephtys cornuta</i>	LIT	Native
<i>Nephtys ferruginea</i>	LIT	Native
<i>Nephtys parva</i>	LIT	Native
Nereididae		
<i>Hediste limnicola</i>	LCRANS, LIT	Native
Phyllodocidae		
<i>Eteone columbiensis</i>	LIT	Cryptogenic
<i>Eteone dilatata</i>	LIT	Native
<i>Eteone lighti</i>	LIT	Cryptogenic
<i>Eteone longa</i>	LIT	Cryptogenic
<i>Eteone spilotus</i>	LCRANS, LIT	Native
<i>Eteone</i> sp.	LCRANS	
<i>Podarkeopsis brevipalpa</i>	LIT	Cryptogenic
<i>Podarkeopsis glabrus</i>	LIT	Cryptogenic#
Syllidae		
<i>Syllis</i> spp.	LIT	

GLYCERIDAE

The family family Glyceridae has been reevaluated by Markus Böggemann (2002). He concluded that of the 172 published species only 42 taxa remain valid. However the polychaete experts and members of the Southern California Association of Marine Invertebrate Taxonomists (SCAMIT 2002) disagreed with many of Böggemann's conclusions regarding Pacific taxa. In light of this lack of agreement on Glyceridae taxonomy the introduction status of many of these species remains unclear.

Glycera americana Leidy 1855

tufted gilled bloodworm

LIT

Origin: Cryptogenic

Glycera macrobranchia Moore 1911

Synonyms *Glycera convoluta*

LIT

Origin: Native

Glycera nana Johnson, 1901

LIT

Origin: Cryptogenic

Kingdom: Animalia
Phylum: Annelida

Glycera tenuis Hartmann 1944

LIT

Origin: Native

Many *Glycera* spp are reported from areas around the world. The actual origin of most of these species and their pattern of introduction is unknown. The genus *Glycera*, commonly known as blood worms, contains species typically found on the bottom of shallow marine waters, living on the sandy or silty bottoms of the intertidal or subtidal regions. Species such as *Glycera dibranchiata*, are extensively harvested for use as bait in fishing. While planktonic larval forms exist they may be demersal.

Hemipodus borealis Johnson, 1901

Syn: *Hemipodus roseus*

LIT

Origin: Native

Found in mudflats and gravelly or sandy beaches, *Hemipodus borealis*, is common along the shore from British Columbia to Southern California.

GONIADIDAE

Glycinde armigera Moore 1911

LIT

Origin: Native

Common along the Southern California coastal shelf, also recorded in the Fraser River this is a species with a marine to brackish salinity tolerance.

Glycinde picta (Berkeley, 1927)

LIT

Origin: Native

There is some debate over the validity of both *G. picta* and *G. polygnatha* as they are very similar morphologically. Genetic or developmental studies might be needed to resolve this question. *G. picta* was described from British Columbia.

Glycinde polygnatha Hartman, 1950

LIT

Origin: Cryptogenic

See *G. picta*

HESIONIDAE

Hesionella mccullochae Hartman, 1939

LIT

Kingdom: Animalia
Phylum: Annelida

Origin: Native

Specimens need to be examined to check the identification. The genera *Hesionella* and *Microphthalmus* are very close morphologically. Species belonging to *Microphthalmus* have been reported from many more localities in the Northeast Pacific than *Hesionella mccullochae*.

Podarkeopsis brevipalpa (Hartmann-Schroeder, 1959)

Synonyms: *Gyptis brevipalpa*

LIT

Origin: Cryptogenic

Probably also includes species mis-identified in the literature as *Podarkeopsis glabrus*.

Podarkeopsis glabrus Hartman 1961

LIT

See *Podarkeopsis brevipalpa*.

NEPHTYIDAE

Nephtys caecoides Hartman 1938

LIT

Origin: Native

Nephtys californiensis Hartman 1938

LIT

Origin: Native

Nephtys cornuta Berkeley and Berkeley, 1945

LIT

Origin: Native

Nephtys ferruginea Hartman 1940

LIT

Origin: Native

Nephtys parva Clark and Jones, 1955

LIT

Nephtys parva is a junior synonym of *N. cornuta*, however the specimens keyed out to this using local references probably belong to an undescribed species.

NEREIDIDAE

Kingdom: Animalia
Phylum: Annelida

Hediste limnicola (Johnson 1903)

Synonyms: *Neanthes limnicola*

LCRANS, LIT

Origin: Native

PHYLLODOCIDAE

Eteone columbiensis Kravitz & Jones, 1979

LIT

Origin: Cryptogenic

Recently described from the Columbia River mouth, this species could be either native or introduced.

Eteone dilatatae Hartman 1936

LIT

Origin: Native

Specimens mentioned in the literature should be examined to check the identification as there are several undescribed species in the Northeast Pacific.

Eteone lighti Hartman 1936

LIT

Origin: Cryptogenic

Described from San Francisco Bay and possibly introduced.

Eteone longa (Fabricius, 1780)

LIT

Origin: Cryptogenic

Specimens mentioned in the literature should be examined to check the identifications as it is unlikely that these are true *E. longa*. There are several undescribed species in the Northeast Pacific.

Eteone spilotus Kravitz & Jones, 1979

LCRANS, LIT

Origin: Native

Probably native, having been found in shelf sediments from California to Washington.

Sedentary Polychaetes

Ampharetidae		
<i>Hobsonia floridana</i>	LCRANS, LIT	Introduced

Kingdom: Animalia
 Phylum: Annelida

Capitellidae		
<i>Barantolla nr americana</i>	LIT	Native
<i>Capitella capitata</i>	LIT	Cryptogenic
<i>Heteromastus filiformis</i>	LIT	Cryptogenic
<i>Heteromastus filobranchus</i>	LIT	Native
<i>Mediomastus acutus</i>	LIT	Native
<i>Mediomastus californiensis</i>	LIT	Cryptogenic
<i>Mediomastus sp.</i>	LCRANS	
Cirratulidae		
<i>Chaetozone spinosa</i>	LIT	Cryptogenic#
<i>Cirratulus cirratus</i>	LIT	Cryptogenic#
Magelonidae		
<i>Magelona hobsonae</i>	LIT	Native
<i>Magelona pitelkai</i>	LIT	Native
<i>Magelona sacculata</i>	LIT	Native
Opheliidae		
<i>Armandia brevis</i>	LIT	Native
<i>Euzonus mucronata</i>	LCRANS	Native
<i>Euzonus williamsi</i>	LIT	Native
<i>Ophelia limacina</i>	LIT	Cryptogenic
<i>Ophelina acuminata</i>	LIT	Cryptogenic
<i>Ophelina breviata</i>	LIT	Cryptogenic#
Orbiniidae		
<i>Leitoscoloplos pugettensis</i>	LIT	Native
Oweniidae		
<i>Owenia fusiformis</i>	LIT	Native
Paraonidae		
<i>Paraonella platybranchia</i>	LIT	Native
Phyllodoceidae		
<i>Phyllodoce spp.</i>	LIT	
Polygordiidae		
<i>Polygordius spp.</i>	LIT	
Spionidae		
<i>Malacoceros fuliginosus</i>	LIT	Cryptogenic
<i>Polydora brachycephala</i>	LIT	Cryptogenic
<i>Polydora cornuta</i>	LCRANS, LIT	Introduced
<i>Polydora sp.</i>	LCRANS	
<i>Prionospio lighti</i>	LIT	Native
<i>Pseudopolydora kempfi</i>	LIT	Introduced
<i>Pseudopolydora sp.</i>	LCRANS	
<i>Pygospio californica</i>	LIT	Native
<i>Pygospio elegans</i>	LCRANS, LIT	Cryptogenic
<i>Scolecopsis foliosa</i>	LIT	Cryptogenic
<i>Scolecopsis squamata</i>	LIT	Cryptogenic
<i>Scolecopsis n. sp. ?</i>	LCRANS	Native
<i>Scoloplos armiger</i>	LIT	Cryptogenic

Kingdom: Animalia
Phylum: Annelida

<i>Spio butleri</i>	LIT	Native
<i>Spio filicornis</i>	LIT	Cryptogenic
<i>Spiophanes berkeleyorum</i>	LIT	Native
<i>Spiophanes bombyx</i>	LIT	Cryptogenic
<i>Streblospio benedicti</i>	LCRANS, LIT	Introduced
Sabellidae		
<i>Manayunkia aestuarina</i>	LCRANS, LIT	Introduced
<i>Manayunkia speciosa</i>	LCRANS, LIT	Introduced
<i>Manayunkia</i> sp.	LCRANS	

AMPHARETIDAE

Hobsonia floridana (Hartman 1951)
Syn: *Hobsonia florida*, *Amphicteis floridus*
LCRANS, LIT
Origin: Introduced

CAPITELLIDAE

Barantolla nr americana Hartman, 1963
LIT
Origin: Native

Specimens need to be examined to check the identification. *Barantolla americana* is found in shelf & slope depths off California. A related form, known as *B. nr. americana*, has been found in shallower water in Puget Sound and Alaska.

Capitella capitata (Fabricius, 1780)
Syn: *Lumbricus capitatus*
LIT
Origin: Cryptogenic

Should be referred to as "*Capitella capitata* complex". Formerly considered a cosmopolitan species but now recognized as a complex of sibling species that vary morphologically, genetically, and developmentally. Extensive laboratory work would be required.

Heteromastus filiformis (Claparde, 1864)
LIT
Origin: Cryptogenic

Considered to be cosmopolitan but records from around the world are likely to contain several species (pers. com. Leslie Harris). Believed to be native to the

Kingdom: Animalia
Phylum: Annelida

Atlantic Ocean from the Gulf of Mexico to the Arctic, it can also be found in South Africa, New Zealand and Australia. The first West Coast record of this worm is from San Francisco Bay in 1936, and it is now well established in California, Oregon, Washington and British Columbia. It is likely transported in sediment and ballast water.

Heteromastus filobranchus Berkeley and Berkeley, 1932

LIT

Origin: Native

Mediomastus acutus Hartman, 1969

LIT

Origin: Native

Mediomastus californiensis Hartman, 1944

LIT

Origin: Cryptogenic

Reported from several areas of the world but validity of all records is unknown, as is the origin and pattern of introduction.

CIRRATULIDAE

Chaetozone spinosa Moore, 1903

LIT

Origin: Cryptogenic – likely mis-identified

Local records are unlikely to be correctly identified. This is a deep-water species and there are no verified shallow water records for the Northeast Pacific.

Cirratulus cirratus (Müller, 1776)

Syn: *Lumbricus cirratus*

LIT

Origin: Cryptogenic – likely mis-identified

Unlikely to be correctly identified. Many previous Northeast Pacific records of this species have been assigned to local species.

MAGELONIDAE

Magelona hobsonae Jones 1978

LIT

Origin: Native

Magelona pitelkai Hartman, 1944

LIT

Kingdom: Animalia
Phylum: Annelida

Origin: Native

Magelona sacculata Hartman, 1961

LIT

Origin: Native

OPHELIIDAE

Armandia brevis (Moore, 1906)

Syn:

LIT

Origin: Native

Euzonus mucronata (Treadwell, 1914)

bloodworms

Syn:

LCRANS

Origin: Native

Euzonus mucronata is common in the upper intertidal of sandy/silty beaches along the West Coast. *Euzonus* have high hemoglobin content turning them a distinctive red color. These worms were found by LCRANS in the high salinity tidal pools along Clatsop spit.

Euzonus williamsi (Hartman, 1938)

Syn:

LIT

Origin: Native

See above.

Ophelia limacina (Rathke, 1843)

Syn:

LIT

Origin: Cryptogenic

A boreal species. Local specimens need comparison to type or topotype material in order to confirm the id.

Ophelina acuminata Oersted, 1843

LIT

Origin: Cryptogenic

Considered cryptogenic here due to the paucity of characters used to distinguish species. Genetic and development studies may be required for speciation.

Ophelina breviata (Ehlers, 1913)

Kingdom: Animalia
Phylum: Annelida

Syn: *Ammotrypane breviata*
LIT
Origin: Cryptogenic#

Known from Arctic and Subantarctic waters. Local specimens are likely to belong to another species.

ORBINIIDAE

Leitoscoloplos pugettensis (Pettibone, 1957)
Syn: *Leitoscoloplos elongatus*
LIT
Origin: Native

A marine species, probably only recorded from sampling at the mouth of the Columbia.

OWENIIDAE

Owenia fusiformis delle Chiaje, 1841 single-tube worm
LIT
Origin: Native

A widely distributed marine species probably only recorded from sampling at the mouth or outside of the Columbia.

PARAONIDAE

Paraonella platybranchia (Hartman, 1961)
Syn:
LIT
Origin: Native

A marine species, probably only recorded from sampling at the mouth or outside of the Columbia.

SPIONIDAE

Malacoceros fuliginosus (Claparede, 1868)
LIT
Origin: Cryptogenic

Specimens need to be compared to type or topotype material to confirm the identification. Found in the Eastern Atlantic in high salinity bays and lagoons.

Dipolydora caulleryi Hartman, 1936

Kingdom: Animalia
Phylum: Annelida

Syn: *Polydora brachycephala*
LIT
Origin: Cryptogenic

Reported from the Columbia River as *Polydora brachycephala* this species has been synonymized with *Dipolydora caulleryi*. *D. caulleryi* is reported from both sides of the US, Europe, and Surinam. Its origin & pattern of introduction is unknown. It is considered an introduced marine polychaete by the California Department of Fish and Game.

Polydora cornuta Bosc, 1802
Syn: *Polydora ligni*
LCRANS, LIT
Origin: Introduced

Verified records are found worldwide. The origin of the species and its pattern of distribution is unknown. Considered by Cohen and Carlton (1995) to be native to the North Atlantic and introduced to San Francisco Bay by the 1930s via ballast water or in association with oyster planting.

Prionospio lighti Maciolek, 1985
LIT
Origin: Native

Pseudopolydora kemp (Southern, 1921)
Syn: *Pseudopolydora kemp japonica*, *P. kemp kemp*
LIT
Origin: Introduced

Native to Japan, there remains some doubt as to whether the local specimens actually belong to this species. Specimens collected by LCRANS were only identified as *Pseudopolydora* sp. The subspecies *Pseudopolydora kemp japonica* has been considered both valid species and a junior synonym of *P. kemp*. We consider it a junior synonym as prior species identifications could not be verified. Also reported as introduced on the West Coast but not from the Columbia River is the closely related species *Pseudopolydora paucibranchiata*. Both species have planktonic larvae and could be readily transported via ballast water.

Pygospio californica Hartman 1936
LIT
Origin: Native

Found in marine intertidal sandflats (Blake 1975)

Pygospio elegans (Claparede, 1863)
Syn: *Spio rathbuni*

Kingdom: Animalia
Phylum: Annelida

LCRANS, LIT
Origin: Cryptogenic

Unknown if this is a species complex or a single widely distributed species; also its origin and pattern of introduction is unknown.

Scoelepis foliosa (Audouin and Milne Edwards, 1833)
Syn: *Nerine foliosa*, *Scoelepis foliosa occidentalis*
LIT
Origin: Cryptogenic

Local specimens need to be compared to type or topotype material to confirm the identification.

Scoelepis squamata (Mueller, 1806)
Syn: *Lumbricus squamatus*
LIT
Origin: Cryptogenic

Local specimens need to be compared to type or topotype material to confirm the identification.

Scoloplos armiger (Müller, 1776)
Syn: *Scoloplos elongata*
LIT
Origin: Cryptogenic

Local specimens may not be the same as the true *S. armiger* from Norway.

Spio butleri Berkeley & Berkeley, 1954
LIT
Origin: Native

Spio filicornis (Müller, 1776)
Syn: *Nereis filicornis*
LIT
Origin: Cryptogenic

Spiophanes berkeleyorum Pettibone, 1962
LIT
Origin: Native

Spiophanes bombyx (Claparede, 1870)
LIT
Origin: Cryptogenic

Kingdom: Animalia
Phylum: Annelida

Another cosmopolitan species that may consist of sibling species.

Streblospio benedicti Webster, 1879

LCRANS, LIT

Origin: Introduced

Origin and pattern of introduction of *Streblospio benedicti* are unknown. This variable species may prove to be another species complex.

SABELLIDAE

Manayunkia aestuarina (Bourne, 1883)

LCRANS, LIT

Origin: Introduced

Local references only used one character to speciate *Manayunkia*. Local records in the literature review must be compared to type or topotype material of *M. speciosa* for confirmation of identification. *Manayunkia aestuarina* is native to eastern North America and may have been introduced via ballast water or in association with stocked fish from eastern North America. EMAP specimens were confirmed as *M. aestuarina*.

Manayunkia speciosa Leidy, 1859

LCRANS, LIT

Origin: Introduced

Local references only used one character to speciate *Manayunkia*. This is inadequate and Local records in the literature review must be compared to type or topotype material of *M. speciosa* for confirmation of identification. *Manayunkia speciosa* is native to eastern North America and may have been introduced via ballast water or in association with stocked fish from eastern North America.

Phylum: Mollusca

SOME FRESHWATER MOLLUSKS OF THE LOWER COLUMBIA RIVER, OREGON AND WASHINGTON

Terrence J. Frest and Edward Johannes

Relatively little is known currently of the freshwater mollusk fauna of the mainstem Columbia River, particularly of its lower reaches, despite frequent visits by malacologists dating to before 1838. Historic data is considerable but mostly unpublished museum records. Much of the more recent information is in the rather

Kingdom: Animalia
Phylum: Mollusca

voluminous gray literature and needs to be reviewed and reidentified. A short survey of 12 sites in late June, 2002 from Portland, Oregon to the estuary provides some useful data as to historic vs. modern freshwater mollusk faunas. One emphasis was to search for so-called exotic (non-indigenous, non-native species). At least one such, the bivalve *Corbicula fluminea*, has been known to be present since perhaps 1937 (Burch, 1944; Counts, 1985).

Though the site coverage is limited, our results indicate that more detailed study would be rewarding. Exotics are more widespread than expected from the literature and native taxa have declined considerably. Still, more than one undescribed taxon was encountered. All of these considerations suggest that detailed survey should be undertaken.

We briefly review below necessary background information on the Columbia River freshwater malacofauna. We then systematically review species found. Finally, we discuss their significance within a historic context and within the wider context of other molluscan introductions.

MOLLUSK FAUNA OF THE COLUMBIA SYSTEM

There has been relatively little published on the malacofauna of the mainstem Columbia River, despite the fact that some of the earliest western U. S. mollusk records are from this stream. There are no particular titles devoted solely to it, in fact. However, numerous references are scattered through the literature and there are large numbers of largely untapped museum records. We have collected the system extensively since 1988. A fair number of recent records are contained in Neitzel & Frest (1989, 1993). Quite a few collections were made from the lower Columbia by NMFS teams during the last 20 years. Unfortunately, the quality of identifications in these latter publications is quite low (note numerous allusions to amnicolids, for example, which are not present). Also, recent revisions have made many of the older literature identifications clearly mistaken. For example, Hershler & Frest (1996) revised the described species of the lithoglyphid *Fluminicola*, one of the two most common Oregon-Washington freshwater snail genera. On their evidence, probably 90-95% of literature records and most museum records are wrong. Recent work by Frest & Johannes (unpublished) indicates a similar error ratio in identifications of the other very common genus, *Juga*. Another very widespread western U. S. genus, *Pyrgulopsis*, has been expanded from about 20 species to about 170 in the last fifteen years (Frest, 1995; Hershler & Sada, 2002). The majority of these new taxa are Western. Taylor (1975) opined that at least half of museum lots of Western freshwater mollusks were wrongly identified; Frest et al. (2002) reiterate this figure for Idaho lots and note that gray literature reports are proportionately even less likely to be correct. Hence, caution should be used in making mollusk identifications from Washington and Oregon freshwater sites, as elsewhere in the West, and dependence on older records is unwise.

Western freshwater habitats differ considerably in taxonomic composition from those elsewhere in the U. S. Large freshwater mussels (unionoids) are relatively non-diverse (about 10 vs. about 300 taxa) and hydrobiids are much more diverse. Only sphaeriids (fingernail clams) are about equal in diversity in both areas. Per site (a) diversity seems lower in the West; but overall (g) diversity is more or less comparable,

with hydrobiids and lithoglyphids making up for the low unionoid diversity. This faunal makeup may be universal for Western stream mollusks (Frest & Johannes, 2002). Western and Eastern malacofaunas differ considerably at the generic level, with the usual pattern being different genera or at least subgenera in those families held in common. Hence, *Fluminicola* (West) vs. *Somatogyrus* (East); *Juga* (West) vs. *Elimia* and 6 other pleurocerid genera (East), etc. It now appears likely that the western hydrobiid swarm differs at the generic level from the eastern also (Hershler, 1994; pers. comm., 2003), instead of *Pyrgulopsis* being common to both.

Leaving aside taxonomic composition, there are significant differences in Western and Eastern-Central U. S. preferred freshwater mollusk habitats as well. Spring and cold, clear, low nutrient, flowing habitats with few macrophytes are more typically Western stream habitats (Frest, 2002a,b), often relatively warm, turbid, with abundant macrophytes and comparatively high dissolved nutrient and lower dissolved oxygen levels, are more significant in the East. Large permanent streams are relatively uncommon in the West. Western drainages are relatively young for the most part but have been considerably modified by geologic factors. Endemism and short-range species are the norm; and perhaps only 40-50% of total diversity has been formally described to date (Frest & Roth, 1995). It is thus not surprising that several new taxa were noted in this brief survey (Table 2). Over the last 15 years, some 100+ newly described species have been added to the Western freshwater mollusk fauna (Frest, 1995). Moreover, Western mollusk biogeographic provinces are small (Frest & Johannes, 2001). There is nothing at all comparable in size in the western U. S. to the Mississippi freshwater Province. In effect, all western freshwater mussels occur in one Province, the Pacific, equal to the Eastern Division; while several very areally limited terrestrial provinces are needed for land forms. Based upon snail genera, terrestrial and freshwater provinces are surprisingly congruent (Frest & Johannes, 2001; in press). Hence, even large streams like the Columbia, Klamath, or Sacramento may range across provinces and not have a uniform fauna in the mainstem, let alone the tributaries. This situation is not limited to mollusks but characterizes the fish fauna as well (McPhail & Lindsey, 1986; Minkley et al., 1986).

In dealing with Western freshwater mollusks, it is important to keep in mind such biogeographic considerations. Despite their relative youthfulness, most Western streams are composite systems geologically, recently assembled and with segmented and composite biotas. This much complicates distributional scenarios both for fish and for mollusks (Minkley et al., 1986; Smith et al., 2000, 2002; Taylor, 1985, 1988a,b; Taylor & Bright, 1987; Hershler & Sada, 2002).

The lower Columbia, not surprisingly after the foregoing, thus had several taxa endemic to it historically and before damming was a cold-water, rocky bottom stream with little in the way of stable soft substrate habitats and macrophyte beds. Dams and dredging have much modified most of the original exposed bedrock (e.g., The Dalles) and hard substrate habitat (Magnuson, 1996). Lower Columbia endemics are believed to have included such taxa as *Fluminicola nuttalliana* and perhaps one other extinct species (see Hershler & Frest, 1996 for discussion); *Vorticifex neritoides*; *Physella columbiana sensu* Taylor (1985) and probably several other taxa. Most of these are either much reduced or perhaps even extinct currently (see Table 2 for most historic species and their habitats). Similarly, reduction in salmon (the glochidial host) runs and in suitable habitat

Kingdom: Animalia
 Phylum: Mollusca

seems to have nearly extirpated the formerly very widespread freshwater mussel *Margaritinopsis falcata* from the lower Columbia and habitat changes alone much reduced others, such as *Gonidea angulata*. Native pleurocerids, hydrobiids, and lithoglyphids have likely also declined considerably. The native lancid *Fisherola nuttalli*, a member of a subfamily or family restricted to the West, is also now quite rare (not found in this survey but living at a few of our lower Columbia sites). We believe that the aberrant planorbid genus *Vorticifex*, another Western endemic, was historically one of the more common lower Columbia snails; it is now one of the more rare.

Perhaps because of habitat changes, introduced taxa such as *Corbicula* are among the most commonly encountered forms. However, as yet relatively few taxa have been introduced. Recent finds of the New Zealand mudsnail are very disturbing, however, and the non-native *Radix auricularia* is a snail community dominant higher in the system (Frest & Johannes, pers. obs.). We expect that, in its current condition, the lower Columbia would provide excellent habitat for the zebra mussel and predict that it could readily become a major pest species, as well as further degrading the native mollusk fauna. The New Zealand mudsnail, *Potamopyrgus antipodarum*, is a serious pest snail in parts of the middle Snake River in Idaho and is rapidly spreading both up and down stream. We have considerably expanded its known range in the lower Columbia River from Astoria and areas ca. 20 miles upstream (Tongue Point) some 60 miles closer to Portland.

Taxonomy herein is based upon the names utilized in Burch (1972-1989), modified where necessary by Taylor (1981) and Turgeon et al. (1998). The latter is the source for common names. We have also used the periodical literature extensively to update all sources and to reflect more recent nomenclatorial changes.

Phylum: Mollusca

Class: Gastropoda

** - considered probably extinct in the lower Columbia River

Ancylidae		
<i>Ferrissia californica</i>	LCRANS, LIT	Native
<i>Ferrissia parallelus</i>	LIT	Native
<i>Ferrissia rivularis</i>	LCRANS, LIT	Native
<i>Ferrissia rowelli</i>	LIT	Native
Hydrobiidae		
<i>Flumicola</i> n. sp. 1	LCRANS	Native
<i>Flumicola</i> n. sp. 2	LCRANS	Native
<i>Flumicola</i> n. sp. 3	LCRANS	Native
<i>Flumicola fuscus</i>	LIT	Native**
<i>Flumicola nuttallianus</i>	LIT	Native**
<i>Flumicola virens</i>	LCRANS, LIT	Native
<i>Potamopyrgus antipodarum</i>	LCRANS, LIT	Introduced
Lymnaeidae		
<i>Fisherola nuttalli</i>	LIT	Native
<i>Fossaria</i> (B.) <i>bulimoides cockerelli</i>	LCRANS	Native
<i>Radix auricularia</i>	LIT	Introduced

Kingdom: Animalia
Phylum: Mollusca

<i>Stagnicola (Stagnicola) apicina</i>	LCRANS, LIT	Native
<i>Stagnicola caperata</i>	LCRANS	Native
<i>Stagnicola (Stagnicola) elodes</i>	LCRANS, LIT	Native
<i>Stagnicola</i> sp. (juv)	LCRANS	
Margaritiferidae		
<i>Margaritopsis falcate</i>	LIT	Native
Olividae		
<i>Olivella biplicata</i>	LIT	Native
Physidae		
<i>Physella (Physella) gyrina</i>	LCRANS, LIT	Native
<i>Physella (Physella) columbiana</i>	LIT	Native
<i>Physella (Physella) hordacea</i>	LIT	Native
<i>Physella (Physella) lordi</i>	LIT	Native
<i>Physella (Physella) propinqua</i>	LCRANS, LIT	Native
<i>Physella (Physella) traski</i>	LIT	Native
<i>Physella (Physella) virginea</i>	LIT	Native
<i>Physella</i> sp.	LCRANS	
Planorbidae		
<i>Gyraulus parvus</i>	LCRANS, LIT	Native
<i>Menetus (menetus) callioglyptus</i>	LCRANS	Native
<i>Menetus dilatatus</i>	LCRANS	Native
<i>Menetus opercularis</i>	LIT	Native
<i>Planorbella subcrenatum</i>	LIT	Native
<i>Planorbella columbiense</i>	LIT	Native
<i>Promenetus umbilicatellus</i>	LIT	Native
<i>Pyrgulopsis</i> n. sp. 1 cf. <i>robusta</i>	LCRANS	Native
<i>Pyrgulopsis</i> n. sp. 6	LIT	Native
<i>Vorticifex effusus effusus</i>	LCRANS	Native
<i>Vorticifex effusus costata</i>	LCRANS, LIT	Native
<i>Vorticifex neritoides</i>	LIT	Native
Pleuroceridae		
<i>Juga (J.)</i> n. sp.	LCRANS	Native
<i>Juga hemphilli</i>	LIT	Native
<i>Juga (J.) plicifera bulimoides</i>	LCRANS	Native
<i>Juga (J.) plicifera plicifera</i>	LCRANS, LIT	Native
<i>Juga silicula</i>	LIT	Native
Polygyridae		
<i>Vespericola</i> sp.	LCRANS	
Viviparidae		
<i>Cipangopaludina chinesis malleatus</i>	LCRANS	Introduced

ANCYLIDAE

Ferrissia californica (Rowell, 1863) fragile ancyloid
LCRANS, LIT

Kingdom: Animalia
Phylum: Mollusca

Origin: Native

Taylor (1981) believes that this name precedes *Ferrissia fragilis* for the common North American river limpet. This taxon is uncommon in the West and seems to prefer low-elevation, rather warm and eutrophic habitats, often with low flow (lotic) or is found in similar lentic habitats, such as ponds and lakes.

Ferrissia parallelus

LIT

Origin: Native

Ferrissia rivularis

LCRANS, LIT

Origin: Native

Ferrissia rowelli

LIT

Origin: Native

HYDROBIIDAE

***Fluminicola* n. sp. 1**

LCRANS

Origin: Native

There appear to be at least three *Fluminicola* in the lower Columbia and two in the lower Willamette. Aside from *virens*, or *virens*-like forms, at least one undescribed taxon occurs in both rivers. Formerly, both likely had the probably extinct *Fluminicola nuttalliana*; and there are historic records for *F. fuscus* (under the name *columbiana*) for the lower Columbia, and possibly the lower Willamette, as well (Neitzel & Frest, 1989, 1993). Hershler & Frest (1996) report another likely extinct taxon from the lower Willamette and possibly from the Columbia below Portland. There are only two remaining lower Columbia taxa found in some numbers; *virens* and this form. Both are probable cold-water stenotopes and often co-occur with *Juga (J.) plicifera plicifera*. Like most larger pebblesnails, this taxon seems to prefer cold and relatively pristine hard-substrate habitats, with little disturbance. Note that this taxon and the foregoing occurred historically in the Columbia upstream only as far as the Hanford Reach, while *fuscus* ranged into the Snake River (Frest, unpub.) and several other interior Washington tributaries (Neitzel & Frest, 1989, 1993; Hershler & Frest, 1996). This taxon has been cited as *Fluminicola* n. sp. 1 in Frest & Johannes (1993, 1995, 1996)

***Fluminicola* n. sp. 2**

LCRANS

Kingdom: Animalia
Phylum: Mollusca

Origin: Native

This *virens*-group taxon seems to be restricted to relatively small and more or less pristine oligotrophic stream habitats. So far, it appears that this undescribed taxon may be restricted to small tributaries in Oregon and Washington below Portland.

Fluminicola fuscus (Haldeman, 1841) Columbia pebblesnail

LIT

Origin: Native

Possibly locally extinct. This species until very recently was confused with several other taxa, and most commonly is cited as *Fluminicola columbiana* Hemphill. Original distribution: Lower Columbia River and a few of its major tributaries in WA, OR, ID, and BC (and probably MT as well). Possibly extinct in the lower Columbia River, WA-OR, and definitely extinct in most of the middle and upper Columbia River, WA, MT, and British Columbia.

Fluminicola nuttallianus

Fluminicola nuttalliana

LIT

Origin: Native

Probably extinct (See Frest on *Fluminicola* n. sp. 1)

Fluminicola virens (Lea, 1838) Olympia pebblesnail

LCRANS, LIT

Origin: Native

This pebblesnail taxon seems characteristic of the lower Columbia and middle to lower Willamette, although similar undescribed taxa occur widely in western Washington and Oregon. There is some possibility that the Columbia form is a distinct species: we are currently exploring that possibility using molecular genetic methods. The group including *virens*, recently redescribed by Hershler & Frest (1996), likely represents a monophyletic clade at a higher taxonomic level than species, as yet unnamed. Note that the common name is completely inappropriate. Pebblesnails are for the most part cold-water stenotopes and historically had very wide distribution in Oregon and Washington clear oligotrophic streams and springs. The common name is mysterious in origin, as the type locality is in Oregon and there is no reason to think Olympia, Washington *Fluminicola* are conspecific.

Potamopyrgus antipodarum (Grey 1853) New Zealand mudsnail

LCRANS, LIT

Origin: Introduced

The New Zealand mudsnail was first noticed in the Columbia River in 1995, at Youngs Bay near Astoria, Oregon (Wonham and Carlton in press). Since then, it

Kingdom: Animalia
Phylum: Mollusca

has been reported as far east as Cathlamet Bay, Oregon. We herein extend the species considerably eastward, to St. Helens, Oregon. Specimens at our two non-estuary sites are as yet quite rare; but massive increases are likely, to judge by the species' history in the middle Snake River. We expect that the Columbia will provide sufficient degraded habitat as to allow this taxon to become a true nuisance species. While Mackie (1999b) does not seem to regard this taxon as a nuisance, except possibly to native mollusks, experiences in the middle Snake River (Bowler & Frest, 1992; Frest & Johannes, 1992) suggest that it not only negatively impacts native mollusks but also can be both an aesthetic irritant and impediment to hydroelectric, trout rearing, and irrigation facilities. Aside from impacts on native species (USFWS, 1995; Richards et al., 2001: see also earlier references in Frest et al., 2002), the species is a biofouler. At one Idaho Power hydroelectric facility, for example, it has proved necessary to operations to remove some 30 tons of organic detritus per day. Half of that by weight is *P. antipodarum*. Impact is further discussed below.

This taxon may have been introduced independently several times into the U. S. Gangloff (1998) regards the Lake Ontario (1991-1994), Idaho (1987), Lower Columbia (1997 sic) and Yellowstone National Park (1995) occurrences as separate. We regard at least the Montana (Yellowstone)) as derived from Idaho sources. There is also another introduction, possibly independent, in the Colorado River system in Arizona (pre-1998). Since 1998, other introductions have turned up in Owens Valley, CA, Polecat Creek, Wyoming likely derived from Yellowstone populations, and in two other areas in coastal and interior Oregon (Frest & Johannes, unpub.). Ballast water is suggested as the venue in Lake Ontario (Zaranko et al., 1997) and generalized in Mackie (1999b) but this hypothesis is untenable for most introductions, the lower Columbia being a possible exception. Several reported introductions have proven incorrect and due to confusion with native hydrobiids. This is a problem in the lower Columbia as well, as native *Pyrgulopsis* occurs here also (see below). Supposed *P. antipodarum* finds should always be confirmed by a specialist.

LYMNAEIDAE

Fisherola nuttalli (Haldeman, 1841) shortfaced lanx
LIT
Origin: Native

The native lanciaid *Fisherola nuttalli*, a member of a subfamily or family restricted to the West, is also now quite rare (not found in this survey but living at a few sites along the lower Columbia). Type locality: "Lower Columbia River" near the old mouth of the Willamette River near Portland, Multnomah Co., OR (could have been from the Willamette River itself). Formerly widespread in the lower Columbia River, Snake River, and a few major tributaries, WA-OR-ID-MT-BC. The lower Columbia River populations are largely extinct due to habitat

Kingdom: Animalia
Phylum: Mollusca

modification caused by Bonneville Power Administration dams and impoundments (Frest and Johannes 1995).

Fossaria (B.) bulimoides cockerelli

LCRANS
Origin: Native

Radix auricularia (Linnaeus, 1758) Big-ear Radix

Syn: *Lymnaea auricularia*
LIT
Origin: Introduced

The non-native *Radix auricularia* is a snail community dominant higher in the Columbia River system (Frest & Johannes, pers. obs.), also introduced in the Great Lakes (<http://nas.er.usgs.gov>) prefers still or standing water, Eurasian aquarium species, first collected from Great Lakes in 1901 (Mills et al. 1993)

Stagnicola (Stagnicola) apicina

LCRANS, LIT
Origin: Native

Stagnicola caperata

LCRANS
Origin: Native

Stagnicola (Stagnicola) elodes (Say, 1821) marsh pondsnail

LCRANS, LIT
Origin: Native

We are more familiar with this taxon as a swamp and wetland taxon in the Midwest. It is relatively rare in the Western U.S. Large stream sites are more common in the West, while the typical eastern site is more likely to be a warm pond or ditch or very small stream. In much of the lower Columbia, including more or less undisturbed habitats, this taxon seems to be replaced by *Stagnicola apicina*, not noted at our sites during this survey.

MARGARITIFERIDAE

Margaritinopsis falcata (Gould 1850) western pearlshell

Syn: *Margaritifera falcata*
LIT
possibly locally extinct

Reduction in salmon (the glochidial host) runs and in suitable habitat seems to have nearly extirpated the formerly very widespread freshwater mussel

Kingdom: Animalia
Phylum: Mollusca

Margaritopsis falcata from the lower Columbia. Original distribution: Southern Alaska to central California, eastward to western Montana, western Wyoming, and northern Utah (Frest and Johannes 1995). Threats such as extensive diversion of rivers for irrigation, hydroelectric, and water supply projects has much reduced the WA, OR, ID, and CA range of this species. In the lower Columbia River region threats include impoundments: continued siltation and other impacts on the few remaining sites with habitat characteristics approximating pre-impoundment conditions on the lower Columbia. Harbor and channel “improvements” in the vicinity of The Dalles and John Day Dam; nutrient enrichment of the lower Columbia due to agricultural run-off. This taxon is declining, in terms of area occupied and number of sites and individuals.

OLIVIDAE

Olivella biplicata

LIT

Origin: Native

PHYSIDAE

Physella (Physella) gyrina (Say, 1821) tadpole physa

LCRANS, LIT

Origin: Native

Physids are among the common river snails in the Western U. S., as they are in the East as well. Taxonomy is badly in need of revision; and we follow Taylor (1981) and Burch (1982) here, recognizing a small number of taxa in the West. Forms of *gyrina* are widespread in a variety of habitats in Western North America. Many literature reports are more likely ascribable to *Physella (Physella) propinqua*. This taxon seems to prefer small stream, pond, and lake habitats locally.

Physella (Physella) columbiana

LIT

Origin: Native

Physella (Physella) hordacea

LIT

Origin: Native

Physella (Physella) lordi

LIT

Origin: Native

Physella (Physella) propinqua (Tryon, 1865) Rocky Mountain physa

Kingdom: Animalia
Phylum: Mollusca

LCRANS, LIT
Origin: Native

In contrast to *Physella (Physella) gyrina*, some forms of this taxon appear to prefer large river habitats, while others are more restricted (Frest & Johannes, 2001). Precise relationships of lower Columbia specimens remain to be determined. In relatively natural Columbia habitats, this taxon is rather rare. It seems to have benefited from siltation and eutrophication.

Physella (Physella) traski
LIT
Origin: Native

Physella (Physella) virginica
LIT
Origin: Native

PLANORBIDAE

Gyraulus parvus
LIT
Origin: Native

Menetus (menetus) callioglyptus (Vanatta, 1895) button sprite
LCRANS
Origin: Native

Note that most sources regard this taxon as *Menetus opercularis* (Gould, 1847); but Taylor (1981) argues that that name applies to snails from Mountain Lake, California and now extinct. This is a widespread taxon in western Washington, northern Oregon, and northwestern California in a variety of habitats. It is usually uncommon in larger streams.

Menetus dilatatus
LCRANS
Origin: Native

Menetus opercularis
LIT

INVALID NAME – See above section on *Menetes callioglyptus*

Planorbella subcrenatum
LIT

Kingdom: Animalia
Phylum: Mollusca

Origin: Native

Planorbella columbiense

LIT

Origin: Native

Promenetus umbilicatellus

LIT

Origin: Native

Pyrgulopsis n. sp. 1 cf. robusta

LCRANS

Origin: Native

This taxon was first noticed in the lower Columbia in the John Day and Bonneville pools by FWS personnel in 1988. Immature specimens possibly belonging to this taxon were noted far downstream during this survey. The taxonomic status of this taxon is currently under investigation using molecular genetic methods. We will need adults to obtain a full suite of morphological characters. The lower Columbia juveniles as live photographed differ in coloration from equivalent life stages of *Pyrgulopsis n. sp. 6* collected upstream. Relationships seem to be with other native U. S. *Pyrgulopsis*, notably *P. idahoensis*, *P. hendersoni*, and *P. robusta* (R. Hershler, pers. comm., 2003; pers. obs.).

Pyrgulopsis n. sp. 6

LIT

Origin: Native

Vorticifex effusus effusus

LCRANS

Origin: Native

Vorticifex effusus costata (Hemphill, 1890)artemisian ranshorn

LCRANS, LIT

Origin: Native

This appears to be the sole surviving species in the genus in the Columbia. We have not seen live *V. neritoides*, limited to the River historically below Portland, in the last few years. For distribution maps of these taxa, see Taylor (1985). *V. effusa costata* seems not to have occurred historically in the River above Grand Coulee. It is also absent from most tributaries, especially on the east side of the Washington and Oregon Cascade Mountains.

Vorticifex neritoides

LIT

Origin: Native

Possibly extinct (see above description)

Kingdom: Animalia
Phylum: Mollusca

PLEUROCERIDAE

Juga (J.) n. sp.

LCRANS

Origin: Native

This undescribed *Juga* taxon may be characteristic of immediate lower Columbia tributaries. It has been noted at several other sites in the first 100 river miles of the Columbia system (Frest & Johanes, unpub.). Sites are typically cold and oligotrophic, with clear water, moderate to high velocity currents, and rocky substrate.

Juga hemphilli

LIT

Origin: Native

Juga (J.) plicifera plicifera (Lea, 1838) pleated juga

LCRANS, LIT

Origin: Native

This is basically a big-river *Juga* species, characteristic of the Lower Columbia and middle-lower Willamette. While formerly much more widespread, it still appears frequently in lower Columbia habitats. Most reports from other streams appear to refer to other subspecies or other *Juga* taxa. Note that historically *Juga* may have reached no farther upstream than just below the Hanford Reach or the mouth of the Yakima River (Frest, unpub.).

Juga silicula

LIT

Origin: Native

POLYGYRIDAE

Vespericola sp.

LCRANS

The common Columbia River taxon is *Vespericola columbianus*; another taxon found in the lower Columbia region is *V. columbianus latilabris*. Other taxa are found by the mouth of the Columbia Gorge. These specimens appear to differ in morphology from any yet described.

VIVIPARIDAE

Cipangopaludina chinesis malleatus (Reeve, 1863) Chinese mystery snail

LCRANS

Origin: Introduced

This non-indigenous taxon has been reported widely in North America (Burch, 1989) but this is the first finding in the Columbia River system. It does not appear likely to become

Kingdom: Animalia
 Phylum: Mollusca

a pest species or to have major negative impact (Mackie, 1999c). However, its occurrence is symptomatic of many others likely to have been so far unnoticed. Hanna (1966) and Mackie (1999c) emphasizes food usage as the rationale for introductions. However, the aquarium trade route is much more likely for most (this species is not mentioned in Mackie, 1999a, nor are apple snails Pomacea). The species is raised specifically for this purpose in the middle Snake River region (Bowler & Frest, 1992) and has commonly seen in pet stores throughout the U. S., as are apple snails, for at least 30 years. Note that all of the non-native taxa mentioned in Bowler & Frest (1992) could quite easily be introduced into the Columbia: many may have already been.

Phylum: Mollusca

Class: Bivalvia

Cardiidae			
	<i>Clinocardium nuttallii</i>	LIT	Native
Corbiculidae			
	<i>Corbicula fluminea</i>	LCRANS, LIT	Introduced
Mactridae			
	<i>Tresus capax</i>	LIT	Native
Margaritiferidae			
	<i>Margaritifera (Margaritifera) falcata</i>	LIT	Native
Myidae			
	<i>Cryptomya californica</i>	LCRANS, LIT	Native
	<i>Mya arenaria</i>	LCRANS, LIT	Introduced
Mytilidae			
	<i>Mytilus edulis</i>	LIT	Cryptogenic
	<i>Mytilus ?trossulus?</i>	LCRANS	Native
Pharidae			
	<i>Siliqua patula</i>	LIT	Native
Pisidiidae			
	<i>Musculium raymondi</i>	LCRANS	Native
	<i>Musculium securis</i>	LIT	Native
	<i>Pisidium casertanum</i>	LCRANS, LIT	Native
	<i>Pisidium compressum</i>	LCRANS, LIT	Native
	<i>Pisidium pauperculum</i>	LCRANS	Native
	<i>Pisidium variabile</i>	LCRANS, LIT	Native
	<i>Sphaerium patella</i>	LIT	Native
	<i>Sphaerium simile?</i> (juv.)	LCRANS	Native
	<i>Sphaerium striatinum</i>	LCRANS	Native
Psammobiidae			
	<i>Nuttallia obscurata</i>	LCRANS	Introduced
Tellinidae			
	<i>Macoma baltica</i>	LCRANS, LIT	Native
Thyasiridae			
	<i>Axinopsida serricata</i>	LIT	Native

Kingdom: Animalia
Phylum: Mollusca

Unionidae		
<i>Anodonta californiensis</i>	LCRANS, LIT	Native
<i>Anodonta kennerlyi</i>	LIT	Native
<i>Anodonta nuttalliana</i>	LIT	Native
<i>Anodonta oregonensis</i>	LCRANS, LIT	Native
<i>Anodonta wahlametensis</i>	LCRANS, LIT	Native
<i>Gonidea angulata</i>	LCRANS, LIT	Native

CARDIIDAE

Clinocardium nuttallii (Conrad, 1837) Nuttall cockle
Synonyms: *Clinocardium corbis*
LIT
Origin: Native

CORBICULIDAE

Corbicula fluminea (Müller, 1774) Asian clam
LCRANS, LIT
Origin: Introduced

Corbiculids were native residents of North America for a considerable time before becoming extinct on the continent relatively recently (Taylor, 1988a,b). The recent introductions from the Western Pacific seem to have begun in the Columbia in the last 75 years; and this corbiculid is now widely distributed across the continent. Taxonomic status of *Corbicula* in North America is still somewhat cloudy, with claims for at least two taxa. More recently, morphological differences within the introduced populations have been ascribed to origin as separate clones of uncertain number, distribution, and status. If more than one taxon is present, the morphological range seen in the Columbia is great enough to suggest that two taxa may be present, although most populations may be mixes of two clones belonging to one only. Despite the early introduction, *Corbicula* is only moderately successful as an invader in Washington and Oregon, especially as compared with, say, the Tennessee Valley. It is a pest species with considerable economic impact in the central and eastern states.

McMahon (1999, fig. 22.2; 2001, fig. 11) seems to restrict *Corbicula* to the lower Columbia in Washington; but the species also occurs commonly to the Idaho border and in the Snake River in Idaho, as well as in Utah (Counts, 1985, 1986). The Idaho records date to at least 1966 (Hanna, 1966; Frest & Bowler, 1993; Frest & Johannes, 2001). McMahon (1999, p. 317) states that *Corbicula* in North America likely derives from a single introduction in northeastern Washington. Presumably, he means southwestern Washington, i.e., the lower Columbia River, as Counts (1986) says.

MACTRIDAE

Tresus capax (Gould, 1850) fat gaper

Kingdom: Animalia
Phylum: Mollusca

LIT

Origin: Native

MARGARITIFERIDAE

Margaritifera (Margaritifera) falcata (Gould, 1850) western pearlshell

LIT

Origin: Native

Populations in the Columbia River greatly reduced due to human mediated erosion, reservoir construction etc. Once an important food item for tribal peoples.

MYIDAE

Cryptomya californica (Conrad, 1837)

false mya, California softshell clam

LCRANS, LIT

Origin: Native

Mya arenaria Linnaeus 1758

softshell clam

LCRANS, LIT

Origin: Introduced

Established from Monterey Bay, CA to Prince William Sound, AK *Mya arenaria* is most abundantly in intertidal and shallow subtidal areas. Probably introduced unintentionally to the West Coast of North America with oyster shipments from the Atlantic coast, *Mya* was later intentionally planted to establish a commercially harvestable population in many West Coast bays.

MYTILIDAE

Mytilus edulis Linnaeus, 1758

blue mussel

LIT

Origin: Cryptogenic

Mytilus edulis is native to the Atlantic Coast. Introduced *M. edulis* have been reported in Puget Sound. Readily confused with *M. trossulus*, it can also hybridize with other *Mytilus* species.

Mytilus trossulus Gould, 1850

bay mussel, foolish mussel

LCRANS

Origin: Native

The native mussel, *Mytilus trossulus*, is often difficult to distinguish from *M. edulis* and *M. galloprovincialis*, two introduced mussels with which it can readily hybridize. No records of the Mediterranean *M. galloprovincialis* exist for the Columbia River Estuary

Kingdom: Animalia
Phylum: Mollusca

but it can be found in other bays along the West Coast and was probably introduced via ballast water.

PHARIDAE

Siliqua patula

LIT

Origin: Native

PISIDIIDAE

Musculium raymondi (Cooper, 1890) lake fingernail clam

LCRANS

Origin: Native

As the common name would suggest, this taxon is most often found in lentic habitats, or at least in low flow situations. It has been found elsewhere in the lower Columbia proper; but not yet here. The most frequent name seen in the literature for this taxon or others resembling it is *Musculium lacustre*; but Taylor (1981) feels that western U.S. populations are best ascribed to a separate taxon. *Lacustre* is a frequently seen taxon in eastern and central North America in warm-water, soft-sediment situations but is rather uncommon in the West (Frest & Johannes, 2001).

Musculium securis (Prime, 1852) pond fingernailclam

LCRANS

Origin: Cryptogenic

Despite the common name, this taxon in the Northwest is most frequently (not often, but increasingly!) seen in larger, warmer rivers with slow flow and definite nutrient enhancement. It is quite uncommon here in pristine habitats but very frequently encountered in eastern North America.

Pisidium casertanum (Poli, 1791) ubiquitous peaclam

LCRANS, LIT

Origin: Native

As the common name implies, this is a very frequently encountered sphaeriid species, perhaps the most widespread native mollusk in the northern hemisphere. It is rapidly spreading currently south of the Equator a well. Very frequent in a wide variety of habitats in the West. For examples, see Frest & Johannes (2001).

Pisidium compressum Prime, 1852 ridgebeak peaclam

LCRANS, LIT

Origin: Native

Kingdom: Animalia
Phylum: Mollusca

TELLINIDAE

Macoma baltica (Linnaeus, 1758) altic macoma
Syn: *M. inconspicua*
LIT, LCRANS
Origin: Native

Common in mid to low intertidal and distributed from San Francisco Bay to the Bering Strait (Ricketts et al. 1985). May have been introduced by man southern most limit in San Francisco Bay (Cohen and Carlton 1995).

THYASIRIDAE

Axinopsida serricata
LIT
Origin: Native

UNIONIDAE

Anodonta californiensis Lea, 1852 California floater
LCRANS, LIT

This mussel is widely but sporadically distributed in eastern Washington but is much less common west of the Cascades in Washington. The species may well be composite (Taylor, 1981; pers. obs.). It is currently rare in the southwestern states and southern California, which area includes the type locality, and is understudy for possible listing there. The species appears to be declining seriously in Washington, including in the Columbia proper.

Anodonta kennerlyi
LIT
Origin: Native

Anodonta nuttalliana Nuttal's floater
LIT
Origin: Native

This native floater has been found, along with *A. oregonensis* and *A. wahlametenis* in the Columbia River Slough by Al Smith (pers com 2004).

Anodonta oregonensis Lea, 1838 Oregon floater
LCRANS, LIT
Origin: Native

Kingdom: Animalia
Phylum: Mollusca

The Oregon floater was first described from the lower Columbia River but appears currently uncommon to rare in it. It is found over much of Washington and Oregon, although seldom in large numbers. Along the Cascade axis, it seems to be replaced by *Anodonta kennerlyi*, and is more often found in streams than that largely lentic taxon.

Anodonta wahlametensis Lea, 1838
LCRANS, LIT
Origin: Native

Willamette floater

Also first described from the lower Columbia and Multnomah Channel, this species has a disjunct range, with only a small portion in the lower Columbia River. Most of the range is in extreme southern Oregon and northern and central interior California. Much of the original range is no longer inhabited by the taxon (Taylor, 1981; 1985; pers obs.)

Gonidea angulata (I. Lea, 1838)
LCRANS, LIT
Origin: Native

Western ridged mussel

Should be a common species in the Columbia River but habitat changes may have significantly reduced its numbers.

Phylum: Arthropoda
Subphylum: Crustacea
Infraclass: Cirripedia

Acorn barnacles, Cirripedia, are conspicuous sessile crustaceans that form volcano like shells of their plates in massive numbers on solid substratums such as rocks, pilings boats and floats. Barnacles are very special crustaceans because they undergo two metamorphic changes (rather than one or none) during development. The acorn barnacles use their feet (cirripedia) to feed on plankton and are economically significant due to the problems the cause when attached to marine structures.

Balanidae		
<i>Balanus crenatus</i>	LIT	Native
<i>Balanus improvisus</i>	LCRANS, LIT	Introduced
<i>Balanus glandula</i>	LCRANS, LIT	Native
<i>Balanus</i> sp. unk	LCRANS	

BALANIDAE

Balanus crenatus Bruguière, 1789
LIT
Origin: Native

Kingdom: Animalia
Phylum: Arthropoda
Subphylum: Crustacea

Bering Sea to Santa Barbara, California. Pleistocene: Alaska, British Columbia, Washington, Oregon, California (Pitumbo & Ross 2002:100). Not expected in the low salinity areas of the Lower Columbia River where reports of it are probably misidentifications of *Balanus improvisus* or *B. glandula*.

Balanus improvisus Darwin, 1854

LCRANS, LIT

Origin: Introduced

Balanus improvisus is tolerant of long exposures to freshwater and full seawater and can reproduce in salinities as low as 10 PSU. *Balanus improvisus* is native to the north Atlantic and has been introduced all over the world on the hulls of sailing ships and with transplanted oysters. The east Pacific distribution of *B. improvisus* is from Vancouver Island, Canada to Monterey, California, and Ecuador (Pitumbo & Ross 2002:101, Carlton 1979:592-597, Zullo 1979, Cohen & Carlton 1995:79-80). The first record of *B. improvisus* in the lower Columbia River specimens occurring on the shells of the native crayfish, *Pacifasticus trowbidgii* collected in brackish waters of Young's Bay in 1957 (Miller 1965, Carlton 1979, Zullo 1979). *Balanus improvisus* is readily distinguished from all other northeast Pacific barnacles by the combination of its calcareous base, extended spur of the tergum, large adductor ridge of the scutum, wall plates with internal tubes and its occurrence in very low salinities.

Balanus glandula Darwin, 1854

LCRANS, LIT

Origin: Native

The most common balanoid of the northeastern Pacific, *B. glandula* occurs in bays and polyhaline waters and on the open rocky coast in the intertidal from the Unalaska Island, Aleutian Islands, Alaska to Bahia de San Quintin, Baja California, Mexico (Henry 1942) and in Pleistocene deposits (Ross 1976). This species was probably introduced to Puerto del Mar del Plata, Argentina (Newman & Abbott, 1980) from the Northeast Pacific. This is the most common barnacle in the lower Columbia River.

***Balanus* sp. unk** (Chapman)

LCRANS

Indeterminate

With carina, rostrum, lateral plates and deep spur of tergum similar to *B. improvisus*. The spur is up to 1/3 width of tergum and is far wider than expected for *B. improvisus*. The sharply quadrate articular ridge of the scutum is aligned with the angular adductor ridge but separated by a deep incision that is partially formed by a hatchet like extension of the articular ridge. The depressor muscle crests of the tergum are wide relative to illustrations of *B. improvisus*. (Specimens from 6.1332x, Port of Ilwaco, Washington, Baker Bay, Columbia River, April 17, 2002.)

Kingdom: Animalia
Phylum: Chordata
Class: Amphibia

Phylum: Arthropoda
Subphylum: Crustacea
Class: Ostracoda

Ostracods were neither targeted nor sent to experts for taxonomic identification by this survey. Further work is needed to determine both the native and introduced species present in the lower Columbia River.

Cyprididae		
<i>Cypria</i> spp.	LIT	
<i>Eucypris</i> spp.	LIT	
Candonidae		
<i>Candona</i> spp.	LIT	
Darwinulidae		
<i>Darwinula stevensoni</i>	LIT	Cryptogenic
Limnocytheridae		
<i>Limnocythere</i> spp.	LIT	

DARWINULIDAE

Darwinula stevensoni (Brady and Robertson, 1870)
Syn: *Polycheles improvisa*, *Polycheles stevensoni*

May be a cosmopolitan ostracod, asexual reproduction, common in European waters.

Phylum: Arthropoda
Subphylum: Crustacea
Subclass: Copepoda

Species descriptions by Jeff Cordell

The following copepods collected in the lower Columbia River consist of those taxa for which a strong case can be made for their status as introduced species. Several of these taxa (*Leimia vaga*, *Tachidius triangularis*) are regarded as cryptogenic because they are small and easily overlooked in typical sampling programs, and their distributions are poorly known. However, they are included in the list because they occur in widely disjunct populations, and/or previous authors have regarded them as introduced to the northeastern Pacific. A number of other harpacticoid copepods were collected in this survey that were described from elsewhere and may have been introduced to the northeastern Pacific. These were not included in this list because they are very widely distributed, (e.g., on both coasts of the United States and in Europe) and/or their taxonomy is poorly known, and therefore their status as introduced or cryptogenic is less

Kingdom: Animalia
 Phylum: Arthropoda
 Subphylum: Crustacea

clear. These species include *Coullana canadensis*, *Huntemmania jadensis*, *Limnocletodes behningi*, *Microarthridion littorale*, *Nannopus palustris*, *Onychocamptus mohammed*, *Paronychocamptus cf huntsmanni*, and *Tachidius (Tachidius) discipes*. Also, several specimens of an unidentified species of Thermocyclops were found in this survey. Thermocyclops has not been previously recorded from western North America, but is widespread, occurring in southeastern North America, Central and South America, Europe, Asia, the Indian subcontinent, and Africa (Ueda and Reid 2003). The disposition of this species as introduced is unknown, and will become clearer if enough specimens can be examined to make a specific identification.

CALANOID COPEPODS		
Calanoid	LIT	
Acartiidae		
<i>Acartia tonsa</i>	LIT	
<i>Acartia</i> sp.	LIT	
<i>Acartia clausi</i>	LIT	
<i>Acartia longiremis</i>	LIT	
<i>Acartiella sinensis</i>	LIT	Introduced
Calanidae		
<i>Calanus</i> sp.	LIT	
<i>Calanus finmarchicus</i>	LIT	
Centropagidae		
<i>Centropages</i> sp.	LIT	
<i>Centropages abdominalis</i>	LIT	
<i>Centropages mcmurrichi</i>	LIT	
<i>Osphranticum labronectum</i>	LCRANS	Cryptogenic
<i>Sinocalanus doerri</i>	LCRANS, LIT	Introduced
Diaptomidae		
Diaptomidae	LCRANS	
<i>Diaptomus ashlandi</i>	LIT	
<i>Diaptomus novamexicanus</i>	LIT	
<i>Diaptomus franciscanus</i>	LIT	
<i>Diaptomus</i> sp.	LIT	
<i>Hesperodiaptomus kenai</i>	LCRANS	Native
<i>Leptodiaptomus novamexicanus</i>	LCRANS, LIT	Cryptogenic
<i>Leptodiaptomus</i> sp.	LCRANS	
<i>Skistodiaptomus pallidus</i>	LCRANS	Cryptogenic
<i>Skistodiaptomus</i> sp., undescribed	LCRANS	
Eucalanidae		
<i>Eucalanus</i> sp.	LIT	
<i>Eucalanus bungii</i>	LIT	
Metridiidae		
<i>Metridia lucens</i>	LIT	
Paracalanidae		
<i>Paracalanus parvus</i>	LIT	
<i>Paracalanus</i> sp.	LIT	

Kingdom: Animalia
 Phylum: Arthropoda
 Subphylum: Crustacea

Pontellidae		
<i>Epilabidocera longipedata</i>	LIT	
<i>Epilabidocera amphitrites</i>	LIT	
Pseudocalanidae		
<i>Clausocalanus arcuicornis</i>	LIT	
<i>Clausocalanus parapergens</i>	LIT	
<i>Ctenocalanus vanus</i>	LIT	
<i>Microcalanus</i> sp.	LIT	
<i>Pseudocalanus</i> sp.	LIT	
<i>Pseudocalanus minutus</i>	LIT	
Pseudodiaptomidae		
<i>Pseudodiaptomus forbesi</i>	LCRANS, LIT	Introduced
<i>Pseudodiaptomus inopinus</i>	LCRANS, LIT	Introduced
Scolecithricidae		
<i>Scolecithricella</i> sp.	LIT	
Temoridae		
<i>Epischura nevadensis</i>	LIT	
<i>Eurytemora affinis</i>	LCRANS, LIT	Native
<i>Eurytemora americana</i>	LIT	
<i>Eurytemora hirundoides</i>	LIT	
<i>Eurytemora</i> sp.	LIT	
Tortanidae		
<i>Tortanus discaudatus</i>	LIT	
CYCLOPOID		
Cyclopoida	LCRANS, LIT	
Corycaeidae		
<i>Corycaeus affinis</i>	LIT	
<i>Corycaeus anglicus</i>	LIT	
<i>Corycaeus</i> sp.	LIT	
Cyclopidae		
<i>Acanthocyclops robustus s.l.</i>	LCRANS	Native
<i>Acanthocyclops vernali</i>	LIT	
<i>Cyclops bicuspidatus thomasi</i>	LIT	
<i>Cyclops vernalis</i>	LIT	
<i>Cyclops</i> sp.	LIT	
<i>Diacyclops thomasi</i>	LCRANS, LIT	Native
<i>Eucyclops cf. elegans</i>	LCRANS	Cryptogenic
<i>Eucyclops conrowae</i>	LCRANS	Cryptogenic
<i>Eucyclops elegans</i>	LCRANS	Cryptogenic
<i>Halicyclops spp.</i>	LCRANS	
<i>Macrocyclus albidus</i>	LCRANS, LIT	Cryptogenic
<i>Mesocyclops edax</i>	LCRANS, LIT	Cryptogenic
<i>Orthocyclops modestus</i>	LCRANS	Cryptogenic
<i>Paracyclops chiltoni</i>	LCRANS	Cryptogenic
<i>Paracyclops fimbriatus</i>	LIT	
<i>Paracyclops poppei</i>	LCRANS	Cryptogenic

Kingdom: Animalia
 Phylum: Arthropoda
 Subphylum: Crustacea

<i>Thermocyclops sp.</i>	LCRANS	Cryptogenic
<i>Limnoithona sinensis</i>	LIT	Introduced
<i>Limnoithona tetraspina</i>	LCRANS	Introduced
Oithonidae		
<i>Oithona similis</i>	LIT	
<i>Oithona spinirostris</i>	LIT	
<i>Oithona sp.</i>	LIT	
HARPACTICOID		
Harpacticoida		
Ameiridae		
<i>Nitocra sp.</i>	LIT	
Canthocamptidae		
<i>Attheyella illinoisensis</i>	LCRANS	Cryptogenic
<i>Attheyella sp.</i>	LIT	
<i>Bryocamptus hiemalis</i>	LIT	
<i>Bryocamptus sp.</i>	LIT	
<i>Canthocamptus robertcokeri</i>	LCRANS	Cryptogenic
<i>Elaphoidella bidens</i>	LCRANS	Cryptogenic
<i>Mesochra alaskana</i>	LCRANS, LIT	Cryptogenic
<i>Mesochra lillijeborgi</i>	LIT	
<i>Mesochra pygmaea</i>	LIT	
<i>Mesochra rapiens</i>	LCRANS	Cryptogenic
<i>Mesochra sp.</i>	LIT	
Canuellidae		
<i>Coullana canadensis</i>	LCRANS, LIT	Cryptogenic
Cletodidae		
<i>Huntemannia jadensis</i>	LCRANS, LIT	Cryptogenic
<i>Leimia vaga</i>	LCRANS, LIT	Cryptogenic
<i>Limnocletodes behningi</i>	LCRANS, LIT	Cryptogenic
<i>Nannopus palustris</i>	LCRANS	Cryptogenic
Cylindropsyllidae		
<i>Paraleptastacus sp.</i>	LIT	
Diosaccidae		
<i>Schizopera knabeni</i>	LIT	
<i>Schizopera sp.</i>	LCRANS, LIT	
Ectinosomidae		
<i>Ectinosoma sp.</i>	LIT	
<i>Microsetella sp.</i>	LIT	
<i>Pseudobradya sp.</i>	LCRANS, LIT	
Harpacticidae		
<i>Harpacticus sp.</i>	LIT	
Laophontidae		
<i>Onychocamptus mohammed</i>	LCRANS, LIT	Cryptogenic
<i>Paronychocamptus cf. huntsmanni</i>	LCRANS, LIT	Cryptogenic
Tachidiidae		
<i>Microarthridion littorale</i>	LCRANS, LIT	Cryptogenic

Kingdom: Animalia
 Phylum: Arthropoda
 Subphylum: Crustacea

<i>Microarthridion</i> sp.	LIT	
<i>Tachidius discipes</i>	LCRANS, LIT	Cryptogenic
<i>Tachidius triangularis</i>	LCRANS, LIT	Introduced
<i>Tachidius</i> sp.	LIT	
Thalestridae		
<i>Diarthrodes</i> sp.	LIT	
OTHER COPEPODS		
Nicothoidae		
<i>Hansenulus trebax</i>	LIT	Native

Pseudodiaptomus inopinus (Burkardt, 1913)

Pseudodiaptomus inopinus is native to the Indo-Pacific, and occurs in a variety of fresh and brackish water habitats from Siberia to the South China Sea, and on both coasts of Japan. The first record of this species on the west coast of the North America was in 1990, from the Columbia River estuary (Cordell et al. 1992). It was subsequently found to be established in many smaller estuaries in the Pacific Northwest, probably via introduction by ballast water (Cordell and Morrison 1996). *P. inopinus* appeared to be a stable and dominant component of the zooplankton in the tidal tributaries of the Columbia River estuary until 2002, when it was found to have been replaced by two other Asian calanoid copepods, *Pseudodiaptomus forbesi* and *Sinocalanus doerri*. Studies on the Chehalis River, which is north of the Columbia River, have found that when *P. inopinus* dominates the plankton in tidal brackish areas it can be important prey of the native shrimps *Neomysis mercedis* and *Crangon franciscorum* (J. Cordell, unpublished data). However, in this survey its abundance peak in the late summer-early fall did not correspond to times when juvenile salmon and other planktivorous fish are present, and it did not occur in their diets. In addition, *P. inopinus* may have ecological effects on other zooplankton. For example, another estuarine copepod, *Eurytemora affinis*, appears to be restricted temporally and spatially with regard to its expected distribution when *P. inopinus* is present (J. Cordell, unpublished data).

Pseudodiaptomus forbesi (Poppe & Richard, 1890)

In its native range, *P. forbesi* has been reported from the Yangtze River in China and from Japan (<http://www.obs-banyuls.fr/Razouls/Webcd/Pseudodiaptomidae.htm>). It was first collected in the northeastern Pacific from the upper San Francisco Bay estuary in 1987 (Orsi and Walter 1991), where it now appears to be a permanent part of the brackish-oligohaline plankton assemblage. This species was first found in the Columbia River estuary in benthic samples taken by the WEMAP survey. Along with *Sinocalanus doerri*, it appears to have replaced *P. inopinus* in this estuary. In 2003 samples from this survey, *P. forbesi* was one of the most abundant mesozooplankton species in tidal tributaries of the main estuary, comprising up to 52% of the plankton numbers in the Grays River. It occurred in the furthest upstream samples taken in this survey, and in

Kingdom: Animalia
Phylum: Arthropoda
Subphylum: Crustacea

summer 2003 comprised 31% of the plankton numbers in Crane Lake, which is located near the city of Portland, Oregon.

Sinocalanus doerri (Brehm, 1909)

This species was introduced to San Francisco Bay from its native range in mainland China (Orsi et al. 1983). In the early 1980s it was the most abundant copepod in the oligohaline-tidal fresh region of the Sacramento-San Joaquin delta, but by the mid 1990s it had declined greatly (Orsi 1999). It first was first reported from the Columbia River estuary in 2002, by this survey. It occurred upstream to Crane Lake near Portland, Oregon, and was very abundant in tidal tributaries of the estuary, where it comprised up to 47% of the plankton numbers in summer 2003 samples.

Limnoithona sinensis (Burkhardt, 1912)

This cyclopoid copepod was first collected in San Francisco Bay estuary in 1979, from the San Joaquin River. It is a fresh water species native to the Yangtze River. It was also collected from the Columbia River from 1979 to 1980 during the CREDDP surveys. This species was not found in the present survey. It has been reported to have disappeared from the San Francisco Bay estuary, having been replaced by its congener *L. tetraspina*, another introduced species (Orsi and Ohtsuka 1999). However, recent analyses of ballast water taken from upper San Francisco Bay in 1999 show that *L. sinensis* was still present at that time (J. Cordell and G. Ruiz, unpublished data). Therefore, this species may still exist in a restricted range in upper San Francisco Bay.

Limnoithona tetraspina (Zhang & Li, 1976)

Limnoithona tetraspina, which is native to the Yangtze River, first occurred on the North American west coast in 1993, in the upper part of San Francisco Bay. Since its introduction there, it has been the most abundant copepod in the bay, with mean abundances of $>10,000\text{ m}^{-3}$. Three specimens of this species were found in 2003 samples from this survey, from both lower (Grays River) and upper (Trojan Power Plant) sites.

Leimia vaga (Willey, 1923)

This harpacticoid copepod can be regarded as a cryptogenic species. Described from Nova Scotia, it is also abundant in many estuaries in Oregon and Washington, where it is restricted to brackish water (J. Cordell, unpublished data), and has also been reported from Prince William Sound, Alaska (Hines and Ruiz, 2000). It was not reported from brackish water habitats in the Nanaimo River estuary, British Columbia in Kask's (1982) checklist of harpacticoids from there. The fact that *L. vaga* has restricted habitat requirements and apparently disjunct populations on the Pacific coast may indicate that it has been introduced.

Tachidius (Neotachidius) triangularis Shen and Tai, 1963

Kingdom: Animalia
 Phylum: Arthropoda
 Subphylum: Crustacea

This species is one of the most abundant harpacticoids in marine-influenced tidal channels of coastal Pacific Northwest estuaries; and also occurs in eelgrass beds. Described from the Pearl River delta, South China, Kask et al. (1982) regarded it as a probable introduction to the Nanaimo River estuary, British Columbia. In this survey *T. (N.) triangularis* occurred in Baker Bay and in the early 1990s it was recorded in Trestle Bay in an unpublished USFWS study.

Phylum: Arthropoda
 Subphylum: Crustacea
 Suborder: Cladocera

Cladocera	LCRANS, LIT	
Bosminidae		
<i>Bosmina</i> sp.	LIT	
<i>Bosmina longirostris</i>	LCRANS	Cryptogenic
Chydoridae		
<i>Alona rustica</i>	LIT	
<i>Alona costata</i>	LIT	
<i>Alona</i> sp.	LIT	
<i>Alona quadrangularis</i>	LIT	
<i>Alona affinis</i>	LIT	
<i>Alona guttata</i>	LIT	
<i>Alonella</i> sp.	LIT	
<i>Camptocercus reticrostris</i>	LIT	
<i>Chydorus sphaericus</i>	LIT	
<i>Chydorus</i> spp.	LCRANS, LIT	
<i>Eurycercus lamellatus</i>	LIT	
<i>Eurycercus</i> sp.	LCRANS, LIT	
<i>Leydigia quadrangularis</i>	LIT	
<i>Leydigia acanthocercoides</i>	LIT	
<i>Leydigia</i> sp.	LCRANS, LIT	
Other Chydoridae	LCRANS	
<i>Monospilus dispar</i>	LIT	
<i>Pleuroxus striatus</i>	LIT	
<i>Pleuroxus denticulatus</i>	LIT	
<i>Pseudochydorus globosus</i>	LIT	
Daphnidae		
<i>Ceriodaphnia pulchella</i>	LIT	
<i>Ceriodaphnia quadrangula</i>	LIT	
<i>Ceriodaphnia reticulata</i>	LIT	
<i>Ceriodaphnia</i> spp.	LCRANS, LIT	
<i>Daphnia parvula</i>	LIT	

Kingdom: Animalia
 Phylum: Arthropoda
 Subphylum: Crustacea

<i>Daphnia retrocurva</i>	LIT	
<i>Daphnia galeata</i>	LIT	
<i>Daphnia rosea</i>	LIT	
<i>Daphnia pulex</i>	LIT	
<i>Daphnia longispina</i>	LIT	
<i>Daphnia</i> spp.	LCRANS, LIT	
<i>Scapholeberis mucronata</i>	LIT	
<i>Scapholeberis</i> sp.	LCRANS	
Holopediidae		
<i>Holopedium gibberum</i>	LCRANS	Cryptogenic
Leptodoridae		
<i>Leptodora kindtii</i>	LCRANS, LIT	Cryptogenic
Macrothricidae		
<i>Ilyocryptus sordidus</i>	LIT	
<i>Ilyocryptus</i> sp.	LIT	
<i>Macrothrix</i> spp.	LCRANS, LIT	
Moinidae		
<i>Moina</i> spp.	LIT	
Polyphemidae		
<i>Evadne nordmanni</i>	LIT	
<i>Pleopsis polyphaemoides</i>	LIT	
<i>Podon leuckartii</i>	LIT	
<i>Podon polyphemoides</i>	LIT	
<i>Podon</i> sp.	LIT	
Sididae		
<i>Diaphanosoma brachyurum</i>	LIT	
<i>Diaphanosoma</i> sp.	LCRANS	
<i>Sida crystallina</i>	LIT	
<i>Sida</i> sp.	LCRANS	

Phylum: Arthropoda
 Subphylum: Crustacea
 Class: Malacostraca
 Peracarida – Cumacea

Section write ups by John Chapman

Cumaceans small motile animals that brood their young in a pouch. Few species produce more than one or two brood in their life but they can reach great abundances in some areas nevertheless. Only *Cumella vulgaris* and *Nippoleucon hinumensis* were collected in the lower Columbia Riversurvey. Both species are tolerant of reduced salinities and are likely to be the only species that permanently reside in the estuary. All other

Kingdom: Animalia
 Phylum: Arthropoda
 Subphylum: Crustacea

Cumacean species reported in the lower Columbia River are either obligate marine species that are perhaps were incidental or are likely misidentifications.

Diastylidae		
<i>Colurostylis occidentalis</i>	LIT	Native^
<i>Colurostylis</i> spp.	LIT	
<i>Diastylopsis dawsoni</i>	LIT	Native^
<i>Diastylopsis</i> spp.	LIT	
Lampropidae		
<i>Lamprops</i> sp. A	LIT	
Leuconidae		
<i>Eudorellopsis</i> sp.	LIT	
<i>Hemileucon comes</i>	LIT	Introduced#
<i>Hemileucon</i> spp.	LIT	
<i>Leucon</i> sp.	LIT	
<i>Nippoleucon hinumensis</i>	LCRANS, LIT	Introduced
Nannastacidae		
<i>Cumella vulgaris</i>	LCRANS, LIT	Native

= probable misidentification, ^ = marine species

DIASTYLIDAE

Anchicolurus occidentalis (Calman, 1912)

LIT

Origin: Native

(Calman, 1912); *Colurostylis* (?) *occidentalis* - Calman, 1912:605,670, figs.100-112; *Colurostylis occidentalis* - Zimmer, 1936:439; Zimmer, 1940:61; Zimmer, 1941:35, fig.44; Lie, 1969:23; *Anchicolurus occidentalis* - Stebbing, 1912:176; Stebbing, 1913:130-131, figs.85-86; Gladfelter, 1975:242, tab.2; Gladfelter, 1975b:275; Bacescu, M., 1992:267,

An offshore marine species not encountered in the present survey and of doubtful occurrence in the non-marine LCR.

Diastylopsis dawsoni (Smith, 1880)

LIT

Origin: Native

Diastylopsis Dawsoni - Smith, 1880:(app B), 215B; Sars 1900:3(5-6):64; *Diastylopsis dawsoni* - Zimmer 1908:8(3):190; Calman 1912:41, 605, 662-666, fig. 81-90; Stebbing 1913:39:110, 111, fig. 66-67; Zimmer 1930:16(4)653; Zimmer 1941:5(1)(4):22, ffigs. 21-22; Zimmer 1943:12(1):169; Gamo 1963:79, pl 12 fig. 1; Lie 1969:23; Gladfelter, 1975b:275.

A probable native species not encountered in the present survey and of doubtful occurrence in the non-marine LCR. A complication with *Diastylopsis dawsoni*, however, is that it has been reported from the North Atlantic, and the western Pacific in addition to

Kingdom: Animalia
Phylum: Arthropoda
Subphylum: Crustacea

the eastern Pacific. Possibly eastern Pacific records of *D. dawsoni* are in fact the extremely similar native *Diastylopsis tenuis* and the western populations are a separate species. On the other hand fully marine species have been introduced to the eastern Pacific (Gosliner, T. 1995. The introduction and spread of *Philine auriformis* (Gastropoda: Opisthobranchia) from New Zealand to San Francisco Bay and Bodega Harbor. *Marine Biology*, 122: 249-255).

LEUCONIDAE

Hemileucon comes Calman, 1907

LIT

Origin: Introduced – probable misidentification

Calman 1907:38-39, pl. 9, figs. 26-32; Bacescu 1988:149.

Hemileucon comes is native to New Zealand and its occurrence in the northeastern Pacific is unconfirmed. However it resembles and thus is a probable misidentified record of *Nippoleucon hinumensis* in the LCR.

Nippoleucon hinumensis Gamo, 1967

LCRANS, LIT

Origin: Introduced

Hemileucon hinumensis - Gamo 1967:151-156, fig 5-7; Cohen & Carlton, 1995:146; *Nippoleucon hinumensis* - Watling 1991:576; Hancock et al., 1997:524,574; Fields, W. & C. Messer, 1999:40; Ruiz, et al. 2000:503; Carlton, J.T., 2001:20.

The type locality of *N. hinumensis* is the brackish water Lake Hinuma, Honshu Japan. In the northeast Pacific, *Nippoleucon hinumensis* has been variously misidentified in collections from the northeast Pacific as *Leucon* or *Hemileucon*. Carlton (1979) did not find it in his comprehensive survey of San Francisco Bay NAS but then it became abundant and widespread in San Francisco Bay since at least 1986 (Cohen & Carlton 1995). *Nippoleucon hinumensis* could be a ballast water introduction (Cohen and Carlton 1995) but it occurs even in estuaries of the NEP that do not receive ballast water traffic, and its spread along the northeastern Pacific coast prior to 1986 is unknown. Other mechanisms of introduction, including transplanted oysters from Japan have not been examined closely. *Nippoleucon hinumensis* is one of the many likely introductions of the NE Pacific that have not yet been published in the peer-reviewed sources. *N. hinumensis* ranges between Elliot Bay, Puget Sound Washington to San Francisco Bay in the NE Pacific (Cohen et al. 2001). Surprisingly Wasson et al. (2001) do not report *N. hinumensis* from Elkhorn Slough, California, which is only 150 km south of San Francisco Bay. *Nippoleucon hinumensis* can readily be confused with *Leucon* or *Hemileucon*.

NANNASTACIDAE

Cumella vulgaris Hart, 1930

LCRANS, LIT

Origin: Native

Kingdom: Animalia
Phylum: Arthropoda
Subphylum: Crustacea

Hart, J.F.L. 1930:37-38, fig.5A-D; Zimmer 1943:154-158, figs.38-47; Lomakina 1958:255-257, fig.171; Lomakina 1968:69, fig.9(7-9); Shih, Figueira & Grainger, 1971:161; Gladfelter 1975:242,244, tab.2; Gladfelter 1975b:275; Valentin 1978:3; Bacescu, M., 1992:227-228.

Cumella vulgaris is tiny and common to abundant on shallow subtidal muddy/sand bottoms, of marine intertidal and rocky intertidal pools. It ranges from Alaska to central California (Gladfelter 1975, Basecu 1992:227) and tolerates extremely broad temperature and salinity ranges. Few species have such broad geographical and physiological ranges. *Cumella vulgaris* may consists of more than one species.

Phylum: Arthropoda
Subphylum: Crustacea
Class: Malacostraca
Peracarida – Tanaidacea

Section write ups by John Chapman

Tanaidacea are distant relatives of Isopoda with long bodies and chelate first walking legs. Tanaidaceans undergo complex sequential sex and morphology changes in response to local population and environmental conditions. The enormous morphological changes greatly complicate taxonomic analyses. The taxonomy of northeastern Pacific tanaidaceans is poorly resolved. As in all peracaridans, juvenile development is direct, requires significant parental care and occurs without a pelagic larval dispersal stage.

Leptocheliidae		
<i>Leptochelia dubia</i>	LCRANS	Cryptogenic
Tanaidae		
<i>Sinelobus stanfordi</i>	LCRANS	Introduced

LEPTOCHELIIDAE

Leptochelia savignyi (Kroyer, 1842)

LCRANS

Origin: Cryptogenic

(Kroyer, 1842); Ishimaru 1985(with citations); Dojiri & Sieg, 1997:213-214,217, figs.3.9, 3.10; Carlton, J.T., 2001:20.

Leptochelia savignyi has also been referred to as *Leptochelia dubia* in the northeast Pacific. However *L. dubia* is one of many synonyms of *L. savignyi*. The *Leptochelia savignyi* complex occurs on all temperate and boreal marine coasts of the northern hemisphere but not in the Arctic Ocean (Ishimaru 1985). The biogeography and taxonomy of the species are too poorly resolved to decipher the native or introduced origins of this species. *Leptochelia savignyi* is a dominant benthic organism in many high salinity areas and its tube building can effect significant alterations of sediment

Kingdom: Animalia
Phylum: Arthropoda
Subphylum: Crustacea

stability in northeast Pacific estuaries. Few *L. dubia* were encountered in the Columbia River.

TANAIDAE

Sinelobus stanfordi (Richardson, 1901)

Syn: *Leptocheilia philetaerus*, *Tanais estuaries*, *Tanais herminiae*, *Tanais philetaerus*, *Tanais stanfordi*, *Tanais sylviae*

LCRANS

Origin: Introduced

Richardson 1901b; Nunomura 1979; Sieg 1976; Lang 1956; Gardiner 1975; Gutu & Ramos 1995; Menzies & Miller 1970 & Miller 1968 (as "*Tanais sp.*"); Sieg, J. & R.N. Winn, 1981:315-343; Sieg, J., 1983:31-39; Heard 2002:376.

Sinelobus stanfordi is a cosmopolitan, tropical and temperate latitude freshwater and marine, shallow water species with a complex taxonomy and massive list of synonymies (Sieg 1980:60-68, Sieg & Winn 1981:329, fig. 6). Very likely transported around the world since 1500 in association with solid ballast, in fouling communities associated the hulls of sailing ships and then again with ballast water and aquaculture transplants. Its origins in the LOWER COLUMBIA RIVER could be due to many mechanisms. The specific name is by consensus among local taxonomists and the species epithet is very unlikely to remain after its taxonomy is more clearly resolved.

Phylum: Arthropoda

Subphylum: Crustacea

Class: Malacostraca

Peracarida – Isopoda

Section write ups by John Chapman

Isopoda occur in fresh and marine waters and in most terrestrial environments. Most isopods are dorsoventally flattened and have 7 pairs of walking legs of similar form. *Argaia*, and *Liriopsis* are parasites of marine fish, encountered only incidentally within the Columbia River. The all native Idoteidae species are marine and also are encountered only incidentally within the lower Columbaia River.

A notable missing species in the lower Columbaia River is the Asian idoteid *Synidotea laevidorsalis* Miers, 1881 introduced to San Francisco Bay over 100 years ago. *Synidotea laevidorsalis* can reproduce in salinities as low as 10 PSU and occurs in Willapa Bay, Washington, immediately north of the lower Columbia River but has not been reported from the lower Columbaia River. Possibly, the record of *Synidotea angulata* (below) was actually *S. laevidorsalis*.

Epicaridea	LIT
------------	-----

Kingdom: Animalia
 Phylum: Arthropoda
 Subphylum: Crustacea

Asellidae		
<i>Caecidotea occidentalis</i>	LCRANS, LIT	Native
<i>Caecidotea racovitzai racovitzai</i>	LCRANS, LIT	Introduced
<i>Caecidotea</i> sp cf <i>racovitzai</i> (fem)	LCRANS	Introduced
<i>Caecidotea tomalensis</i>	LIT	Native
<i>Caecidotea</i> sp.	LCRANS	
Bopyridae		
<i>Argeia pugettensis</i>	LIT	Native
Chaetiliidae		
<i>Mesidotea entomon</i>	LCRANS, LIT	Cryptogenic
Cirolanidae		
<i>Excirrolana chiltoni</i>	LCRANS	Native
Cryptoniscidae		
<i>Liriopsis pygmaea</i>	LIT	Native#
Idoteidae		
<i>Idotea fewkesi</i>	LIT	Native
<i>Synidotea angulata</i>	LIT	Native#
<i>Synidotea</i> spp.	LIT	
Ligiidae		
<i>Ligia pallasii</i>	LCRANS	Native
Limnoriidae		
<i>Limnoria lignorum</i>	LIT	Native
Oniscidae		
<i>Porcellio scaber</i>	LIT	
Sphaeromatidae		
<i>Bathycopaea daltonae</i>	LIT	Native
<i>Gnorimosphaeroma insulare</i>	LCRANS	Native
<i>Gnorimosphaeroma oregonense</i>	LCRANS, LIT	Native
<i>Gnorimosphaeroma</i> spp.	LIT	
<i>Tecticeps convexus</i>	LIT	Native#

= probably misidentification

ASELLIDAE

The epigeal *Asellus* of the northeast Pacific consist of the native *A. alaskensis* Bowman & Holmquist, 1975, *A. occidentalis* Williams, 1972 and *A. tomalensis* Bowman 1974 and the introduced nonindigenous *A. hilgendorfi* Bovallis, 1886 and *A. racovitzai racovitzai* Williams, 1970. The incomplete taxonomy and geographical information on these species greatly complicates efforts to resolve their origins.

Caecidotea occidentalis (Williams 1970)

LCRANS, LIT
 Origin: Native

Kingdom: Animalia
Phylum: Arthropoda
Subphylum: Crustacea

See discussions of *A. tomalensis* and *A. racovitzai racovitzai* below.

Caecidotea racovitzai racovitzai (Williams, 1970)

LCRANS, LIT

Origin: Introduced

Asellus racovitzai racovitzai - Williams 1970:16, 17, 43-47, figs. 29, 31, 32; *Asellus communis* – Racovitzai 1920:79-115; *Asellus tomalensis* – Winger et al. 1972:;
Caecidotea racovitzai – Toft et al. 1999:; Toft et al. 2002:190, 193, fig. 2.

The palm of the propodus of the first pereopod bears a triangular process near the midpoint and the first pleopod of the male is subequal to the second pleopod. The mesial process of the endopod of the second male pleopod is present and the cannula is relatively long and narrow with the caudal process acutely pointed. *Asellus communis* was the first species of North American *Asellus* to be described. Say's (1818) brief description provided no details or figures of the male sexual pleopods. It is uncertain whether any of the several subsequent redescriptions of this species (none of which referred to the type material) in fact apply to *A. communis* Say, 1818. Williams (1970) reports *Asellus racovitzai racovitzai* and *A. communis* from broad regions of the eastern Great Lakes and the northeastern U.S. Williams's (1970) western North American records of both species are from Echo Lake, Kings County, Washington.

The male triangular extension of the mid propodus of pereopod 1 and three tipped endopodite of the second pleopod clearly distinguish *Asellus racovitzai* from *A. communis*, *A. occidentalis* and *A. tomalensis*. Hatch (1947) reports *A. communis* from Ontario, Quebec and Nova Scotia, from the Arboretum, and the Plantation Pond, Lake Washington, Univ. Washington Campus. This species has been referred under *Caecidotea* sp. (Smith 2001, Thorp & Covich 2001). However, Birstein (1951:48-59) argues for the synonymy of *Caecidotea* under *Asellus*, which appears to have been accepted by Williams (1970) and Bowman (1974) and Miller (1975). Hatch's (1947) records and others assumed by Bowman (1975) to be *A. occidentalis* are not confirmed and could be in fact be *A. racovitzai* or *A. communis* "occurring in the side channels and on vegetated shores in areas of dense aquatic vegetation of the Columbia River".

Toft et al. (2002) review the criteria for nonindigenous species that apply to the possible introduction of *C. racovitzai* to the San Francisco Bay delta

Origins: Very likely, an introduction from the eastern U.S. but requiring more detailed taxonomic analyses.

Caecidotea* sp cf *racovitzai (Chapman)

LCRANS

These female specimens are possibly *Caecidotea racovitzai racovitzai* but cannot be identified with confidence. They should be counted as a record for the genus but not as additions to species lists.

Caecidotea tomalensis Harford, 1877

LIT

Origin: Native– possibly misidentified

Kingdom: Animalia
Phylum: Arthropoda
Subphylum: Crustacea

Asellus tomalensis Harford 1877:53-54; Richardson 1904a:224-226, figs. 110-112; Richardson 1904b:668-669, figs. 15-17; Richardson 1905:431-433, figs. 487-489; Johansen 1922:156; Fee 1926:20-21; Van Name 1936:459-461, fig. 288 (part); Van Name 1940:133; Carl 1937:451; Hatch 1947:170-171, figs. 31-32; Ellis 1971:passim; Bowman 1974:431-441, figs. 9-11, 18-20, 26-28, 29-39, 35-37; Miller 1975:298, 308.

The dactyl and propodus palm of pereopod 1, postmandibular lobes of head and the distal endopod of male pereopod 2 closely match *A. occidentalis* of Williams (1970) and Bowman (1974). However, the male pleotelson is more similar to Williams (1970, fig. 53G) than to Bowman (1974, fig. 18). The pleotelson shape is constant among males ranging from 3-8 mm in length in sample 8.501x.

Asellus occidentalis is distinguished from *A. communis* and *A. racovitzae* (the only other species known from Washington and Oregon) by the absence of an anterior tooth and mid triangular process on the palm of the propodus of male pereopod 1 and by the absence of a process on the lateral edge of the base of the endopod of the male second pleopod. Characters that distinguish these *Asellus occidentalis* from the *A. tomalensis* are the long, triangular apex of the endopod of male pleopod 2, which is rounded in *A. tomalensis*, and the acute postmandibular lobes of the head, which are evenly rounded in *A. tomalensis*. The steeply inclined posterior edges of the telson of these specimens more closely match *A. tomalensis* of Bowman (1974). Ellis (1971) found *A. occidentalis* (as *A. tomalensis*) in an intermittent pond adjacent to the south fork of the Klaskanin River in Clatsop County, Oregon, but not in apparently suitable habitats of the south fork above and below the pond. Williams (1970) in his revision of 14 epigeic species of North American *Asellus*, lists only *A. occidentalis* as restricted to the Pacific coast (Oregon, Washington, British Columbia). The only other Pacific coast epigeic species that Williams listed, (*A. communis* and *A. racovitzai*) were known then only from Echo Lake, Washington. Both *A. communis* and *A. racovitzai* were collected by E. L. Bousfield 20 August 1955. Both of these species appear to be introduced to the Pacific coast from the eastern United States (Bowman 1974, Toft et al. 2002).

Williams (1970:13) considered Bousfield's material from Echo Lake and personal notes to be "of considerable interest" and included them in his publication: Bousfield personal communication to Williams, (1 Sept. 1967) "Echo Lake is the type locality of *Crangonyx richmondensis occidentalis* H. & H., one of a species complex that is usually found together with *A. communis* in the east. . . . *Crangonyx pseudogracilis* Bousf., formerly thought to be endemic to eastern North America, has also turned up in material from Oregon and Washington cf. Bousfield, 1961, and indicates that freshwater peracaridans may have much wider distributions than formerly believed." Indeed, since Bousfield's 1967 note, the introductions of *C. pseudogracilis* (Costello 1993, Chapman 2000) and *A. communis* (Williams 1972, Chapman 2000) were discovered in Europe, and *A. racovitzai* has since appeared in San Francisco Bay (Toft et al. 1999, 2002).

Williams (1972) described his Pacific coast *Asellus* material as a new species (*Asellus occidentalis*) rather than *Asellus tomalensis* Harford, 1877 (as others had done e.g., Fee

Kingdom: Animalia
Phylum: Arthropoda
Subphylum: Crustacea

1926, Carl 1937, Hatch 1947, Ellis 1971). William's justification of this designation was that the published descriptions of *A. tomalensis* were inadequate (Bowman 1974). The single type specimen of *A. tomalensis*, collected by Lockington in "Tomales Bay, and vicinity", California, was in the California Academy of Sciences, collections that were destroyed in the 1906 San Francisco earthquake. Attempts by William's colleagues to collect more specimens from Tomales Bay were unsuccessful. Williams was therefore uncertain whether *A. occidentalis* was in fact, a distinct species from *A. tomalensis*. Bowman, concluding that *C. tomalensis* is a freshwater species, and accordingly searched adjacent creeks and ponds of Tomales Bay for it but without success. However, E. Iverson and J. T. Carlton later found specimens in a shallow pond adjacent to Bolinas Lagoon, less than 24 km south of Tomales Bay. Bowman's (1974) compared these topotypes with *A. occidentalis* and concluded that the two species are valid. However, the synonymies proposed by Bowman (1974) are for dates prior to Williams 1970 since the specimens were not examined.

The discovery of *A. racovitzai* in the Columbia River (see below) and its recent appearance in San Francisco Bay, since its discovery in Echo Lake in 1955, indicate that it is spreading on the Pacific coast. The inability of William's colleagues and of Bowman to find *A. tomalensis* around Tomales Bay suggests that this species has a restricted or limited distribution in the region. Toft et al. (2002) could not confirm previous records of *A. tomalensis* in San Francisco Bay its absence the bay delta prior to European settlement while occurring in surrounding drainages is unlikely. The exclusive occurrence of *A. racovitzai* and *A. hilgendorffii* and complete absence of *A. tomalensis* in thousands of samples from the San Francisco Bay delta may indicate the local extinction of *A. tomalensis* and perhaps its replacement by *A. racovitzai* and *A. hilgendorffii*. Native to eastern Pacific but perhaps confused in the Columbia River with native or nonindigenous species.

BOPYRIDAE

Argeia pugettensis Dana, 1853

Syn: *Argeia pauperata* Stimpson, 1857; *Argeia calmani* Bonnier, 1900; *Argeia pingi* Yu 1935.

LIT

Origin: Native

Ranging from the Bearing Sea to southern California, Japan and Korea, *Argeia pugettensis* is a branchial parasite of Crangonid shrimps.

CHAETILIIDAE

Mesidotea entomon (Linnaeus, 1767).

Mesidotea Richardson, 1905. = *Saduria* Adams in White, 1852; Kussakin 1982:73-77, figs. 49-50; *Saduria entomon* - Schultz 1969:59, fig. 63.

LCRANS, LIT

Kingdom: Animalia
Phylum: Arthropoda
Subphylum: Crustacea

Origin: Native

Mesidotea entomon was formerly placed under *Oniscus* and several other genera. However, the identity of this large isopod has remained clear in the literature. In the northeast Pacific this species has been commonly known as *Saduria entomon*. The species is rare in muddy sands and gravels of coastal rivers, bays and beaches of Washington and Oregon but attracts attention due to its large size, reaching 30 mm in length.

Distribution - Circumpolar, western coast of North America to Pacific Grove, CA; Stockholm, Germany, Labrador, Kara Sea.

CIROLANIDAE

Excirrolana chiltoni (Richardson, 1905)

LCRANS

Origin: Cryptogenic

(Formerly placed in *Cirolana*). British Columbia to CA; Japan, Taiwan, Hong Kong. Intertidal. = *E. kincaidi* (Hatch, 1947); = *E. vancouverensis* (Fee, 1926); = *E. japonica* Richardson, 1912 (See Brusca *et al.* 2004)

An open coastal and marine bay species of clean sand. Northeast Pacific species of these environments are commonly considered to be native due to the vast majority of other species in those habitats that are native. However, the spread of the introduced Asian clam *Nuttallia obscurata* in these same environments from along the coast (see below) and the probable introduction of the surf zone diatom *Chaetocerus armatum* (Lewin, J. and Norris, R.E. 1970, Lewin, J. and Rao, V.N.R. 1975, Lewin, J. and Schaefer, C.T. 1983) indicate that the origins of many of these species should be examined more carefully.

CRYPTONISCIDAE

Liriopsis pygmaea (Rathke, 1843)

LIT

Origin: Native– probably misidentified

An obscure nearly cosmopolitan hyperparasite of rhizocephalin barnacles that infect lithodid crabs and hermit crabs (Lovrich *et al.* 2004). The occurrence of this species in the LOWER COLUMBIA RIVER should be held in doubt since none of the hosts are reported from the LCR.

IDOTEIDAE

Idotea fewkesi Richardson, 1905

LIT

Origin: Native

Kingdom: Animalia
Phylum: Arthropoda
Subphylum: Crustacea

Richardson 1905:359-360, fig. 387-388; Fee 1926:17-18; Hatch 1947:218; Mezies 1950:161-164, pl. I, fig. A-I; Schultz 1969:76, fig. 93; Kussakin 1982:147-148, ffig. 108.

A common inhabitant of shallow water and rocky intertidal macrophytes from Alaska to southern California . *Idotea fewkesi* is a probable incidental species of the LOWER COLUMBIA RIVER and unlikely permanent resident.

Synidotea angulata Benedict, 1897

LIT

Origin: Native- possibly misidentified

Benedict 1897:395-396, fig. 6; Richardson 1899a:847-848, Richardson 1899b:268; Richardson 1905:376, figs. 418-419; Hatch 1947:220, fig. 97; Schultz 1969:68, fig. 77; Kussakin 1982: 245-247, figs. 181-182; Rafi & Laubitz 1990:2674, figs. 19-20;

The range of *Synidotea angulata* is British Columbia to Northern California and it occurs in full marine deep waters (57-69 m) that would not be expected in the LCR. However, *S. angulata* resembles and could be confused with the introduced *Synidotea laevidorsalis* Meirs, 1881. *Synidotea laevidorsalis* is a full estuarine low salinity species introduced over 100 years ago, (Chapman and Carlton 1991, 1994) but is known in the eastern Pacific only from San Francisco Bay, California and Willapa Bay, Washington.

Origin: Native if correctly identified. *Synidotea laevidorsalis* was reported for the first time in the northeast Pacific along with the original description of *S. angulata* (Benedict, 1897).

LIGIIDAE

Ligia pallasii Brandt, 1833

LCRANS

Origin: Native

Van Name 1936:46-44, fig. 7 (with synonymy); Hatch 1947:187-188.

Ligia pallasii is a cockroach-like isopod that scavenges decaying plant and animal material. It occurs in deep crevices of high intertidal rocky areas predominantly on open coasts and often near freshwater seeps. Females reach 2.5 cm in length.

Distribution - Alaska to Santa Cruz, California.

LIMNORIIDAE

Limnoria lignorum (Rathke, 1799)

LIT

Origin: Native

Hatch 1947:211-212, fig. 81; Kussakin 1979:315-316, figs. 181-182 (with synonymy).

Limnoira lignorum is conspicuous where it occurs because it bores into wood.

Kingdom: Animalia
Phylum: Arthropoda
Subphylum: Crustacea

Distribution – Kodiak, Island, Alaska to Pt. Arena, California, 0-20 m, tolerant of low salinities.

ONISCIDAE

Porcellio scaber Latreille, 1804

LIT

Origin: Introduced

Van Name 1936:226-227 (with synonyms)

The most common introduced terrestrial isopod of western North America. A cosmopolitan species of European origin. = *Porcellio scaber americanus* Arcangeli, 1932 (Brusca et al. 2004)

SPHAEROMATIDAE

Bathycopea daltonae (Menzies and Barnard, 1959)

LIT

Origin: Native

Ancinus daltonae - Menzies and Barnard, 1959:31, fig. 25; *Ancinus granulatus* - Holmes & Gay; Schultz 1969:115; *Bathycopea daltonae* – Lyola & Silva 1971:217-222, fig. 5-7. Subtidal marine species of medium coarse gray sands. Monterey Bay to San Miguel Islands, CA. 19-20 m. Occurrence of this species in the Columbia River would be a range extension and thus is more likely to be a misidentification.

Distribution – Monterey to Santa Cruz Island, California, unless this record stands.

Gnorimosphaeroma insulare (Van Name, 1940)

LCRANS

Origin: Native

Syn: *Gnorimosphaeroma lutea* (Van Name, 1940). The species was formerly placed also in *Exosphaeroma* (Brusca et al. 2004, Kussakin 1979:409-410, figs. 263-264).

Distinguished from *G. oregonense* by pointed rather than square hinge notches between telson and 3rd pleonite, by the projection of the 3rd pleonite short of the lateral edge of the pleon. Morphological differences between *Gnorimosphaeroma insulare* and *G. oregonense* are subtle (especially comparing 4.897x and 5.898x of 10 July 2002, Young's Bay Rip rap). The shape of hinge notches vary with angle of perspective and the only illustrations are at different angles. The extension of the third pleonite and the body length are also variable. There seems to be no salinity gradation associated with their distributions in the Columbia River and they seem doubtfully distinct species.

Distribution - Popof Island, Alaska to San Nicolas Island, California. Fresh and brackish water estuaries and lagoons along the northeast Pacific coast. = *G. oregonensis lutea* Menzies, 1954; = *G. lutea* Menzies, 1954.

Kingdom: Animalia
Phylum: Arthropoda
Subphylum: Crustacea

Hoestlandt, H. 1977. Description complementaire de l'isopode flabellifere
Gnorimosphaeroma insulare Van Name et synonymie de *G. luteum* Menzies avec cette
espece. *Crustaceana* **32**:45-54.

Gnorimosphaeroma oregonense (Dana, 1852)

LCRANS, LIT

Origin: Native

Syn: *Sphaeroma oregonensis*, *Sphareoma olivacea*, *Exosphaeroma oregonensis*,
Neosphaeroma oregonense, *Gnorimosphaeroma oregonensis oregonensis*
(Dana, 1852); *Sphaeroma oregonense* Dana, 1852:778; Atlas, 1855:pl.52; Stimpson,
1857:509; Richardson, 1899:836; Richardson, 1900a:223; Richardson, 1904b:214;
Richardson, 1904c:659; Richardson, 1905:216; *Sphaeroma olivacea* Lockington,
1877:45; *Exosphaeroma oregonensis* Richardson, 1905b:296-298, figs.315,316;
Richardson, 1909:92; Van Name, 1936:450-451, fig.282; Hatch, 1947:213, figs.82-83;
Neosphaeroma oregonense Monod, 1932:67-82, fig.74; Monod, 1936:123-
124(partim:fig.70); *Gnorimosphaeroma oregonensis oregonensis* Menzies, 1954:8-11,
fig.5,7A-E, 12; Riegel, 1959:272-284; *Gnorimosphaeroma oregonense* Hoestlandt,
1964:872-877; Miller, 1968:12-13; Schultz, 1969:129, fig.187a; Hoestlandt, 1973b:355-
369, figs.1-9; Kussakin, O., 1979:406-407,409, figs.260-262.

Distinguished from *G. insulare* by square rather than pointed hinge notch between telson
and 3rd pleonite, by the projection of the 3rd pleonite to the lateral edge of the pleon.
This is one of the most ubiquitous northeast Pacific coastal isopods.
Distribution - Central California to Alaska, intertidal to 22 m.

Tecticeps convexus Richardson, 1899

LIT

Origin: Native

Tecticeps convexus - Richardson, 1899:837; Richardson 1905b:278, figs. 290-291;
Kussakin 1979:347-350, figs. 210-211.

The previously known range of *T. convexus* is Oregon border to Point Conception,
California (Brusca et al. 2004). Thus, specimens from the Columbia River would be a
range extension or, the specimens could also be misidentified. *Tecticeps convexus* is a
full marine species that occurs at depths of 0- 9 m

Phylum: Arthropoda

Subphylum: Crustacea

Class: Malacostraca

Peracarida – *Amphipoda*

Section write up by John Chapman

Kingdom: Animalia
 Phylum: Arthropoda
 Subphylum: Crustacea

The large order Amphipoda is represented locally by the suborders, Gammaridea, Caprellidea and Hyperidea but only the Gammaridea permanently occupy the lower Columbia River. The Gammaridea however, are by far the most abundant and familiar suborder of benthic Crustacea in the fresh, brackish and marine waters of the lower Columbia River and occupy even the supralittoral fringe and in a few almost terrestrial habitats. Gammaridean amphipods, brood their eggs in a pericardial pouch from which the fully formed young emerge. The juveniles do not have a specialized larval dispersal stage. The native *Corophium salmonis* and *Corophium spinicorne* are critical food sources of juvenile salmon in the lower Columbia River.

Ampeliscidae			
	<i>Byblis</i> spp.	LIT	
Ansiogammaridae			
	<i>Anisiogammarus</i> sp.	LIT	
	<i>Eogammarus confervicolus</i>	LCRANS, LIT	Native
	<i>Eogammarus</i> sp. A	LCRANS, LIT	Native
	<i>Eogammarus</i> sp.	LIT	
	<i>Rammellogammarus oregonensis</i>	LIT	Native#
	<i>Rammellogammarus</i> sp. A	LCRANS, LIT	
Aoridae			
	<i>Grandidierella japonica</i>	LCRANS, LIT	Introduced
Atylidae			
	<i>Atylus tridens</i>	LIT	Native
Corophiidae			
	<i>Americorophium brevis</i>	LCRANS, LIT	Native
	<i>Americorophium salmonis</i>	LCRANS, LIT	Native
	<i>Americorophium spinicorne</i>	LCRANS, LIT	Native
	<i>Corophium acherusicum</i>	LIT	
Crangonyctidae			
	<i>Crangonyx floridanus</i> subgroup	LIT	Cryptogenic#
	<i>Crangonyx pseudogracilis</i>	LCRANS	Introduced
	<i>Crangonyx</i> spp.	LIT	
Haustoriidae			
	<i>Eohaustorius brevicuspis</i>	LCRANS	Native
	<i>Eohaustorius estuaries</i>	LCRANS, LIT	Native
	<i>Eohaustorius</i> sp.	LIT	
	<i>Monoporeia affinis</i>	LIT	Cryptogenic#
	<i>Monoporeia</i> sp.	LIT	
Hyalellidae			
	<i>Hyalella azteca</i>	LCRANS, LIT	Cryptogenic
Hyalidae			
	<i>Allorchestes angusta</i>	LIT	Native
Hyperiididae			
	<i>Hyperoche</i> spp.	LIT	

Kingdom: Animalia
 Phylum: Arthropoda
 Subphylum: Crustacea

Isaeidae		
<i>Photis macinerneyi</i>	LIT	Native
<i>Photis</i> spp.	LIT	
Lysianassidae		
<i>Hippomedon columbianus</i>	LIT	Native
Melitidae		
<i>Melita cf. nitida</i>	LCRANS	Introduced
Oedicerotidae		
<i>Americhelidium shoemakeri</i>	LIT	Native
<i>Americhelidium</i> spp.	LIT	
<i>Pacifoculodes spinipes</i>	LIT	Native
<i>Pacifoculodes</i> spp.	LIT	
Phoxocephalidae		
<i>Foxiphalus obtusidens</i>	LIT	Native
<i>Grandifoxus grandis</i>	LCRANS, LIT	Native
<i>Mandibulophoxus gilesi</i>	LIT	Native
<i>Paraphoxus</i> sp.	LIT	
<i>Rhepoxynius abronius</i>	LIT	Native
<i>Rhepoxynius daboius</i>	LIT	Native
<i>Rhepoxynius heterocuspadata</i>	LIT	Native
<i>Rhepoxynius tridentatus</i>	LIT	Native
<i>Rhepoxynius</i> spp.	LIT	
Talitridae		
<i>Megalorchestia pugettensis</i>	LCRANS	Native
<i>Traskorchestia traskiana</i>	LCRANS	Native

ANSIOGAMMARIDAE

Only two native species of Anisogammaridae appear to exist in the present lower Columbia River, *Eogammarus confervicolus* and *Ramellogammarus sp. A*. *Ramellogammarus sp. A* appears to be a new species, distinct from *Ramellogammarus oregonensis* and *R. vancouverensis*.

Eogammarus confervicolus (Stimpson, 1856)

LCRANS, LIT

Origin: Native

Mara confervicola - Stimpson, 1856:90; *Gammarus confervicolus* - Stimpson, 1857:520-521; Holmes 1904:239; Bate, 1862:218, pl.38, fig.9; *Melita confervicola* - Stebbing, 1906:428; *Anisogammarus (Eogammarus) confervicolus* - Barnard 1954a:9-12, pls.9-10; Bousfield, 1958:86, fig.10; Tzvetkova 1972; Tzvetkova 1975; *Anisogammarus confervicolus* - Saunders 1933:248; Carl 1937; Barnard, J.L. 1954; Filice, F.P., 1958:183; Shoemaker, 1964:423-427, figs.14-15; Bousfield, E.L. & J.D. Hubbard, 1968:3; Barnard 1975:351,358; *Eogammarus confervicolus* - Bousfield 1979:317-319, fig.4; Klink, R.W. 1980:242; Barnard & Barnard 1983:585; Austin, 1985:607; Carlton, J.T. & J. Hodder 1995:725; Staude, 1997:373, 383, fig. 18.75; Bousfield 2001:108.

Kingdom: Animalia
Phylum: Arthropoda
Subphylum: Crustacea

Among the most prevalent species of estuary samples sites in the LCR. Sample 14.1135x has a particularly large specimen.

Distribution - Southeastern Alaska to southern California, 0-30m .

Eogammarus oclairi Bousfield, 1979

LIT

Origin: Native

Eogammarus oclairi - Bousfield 1979:319-321; Barnard & Barnard 1983:585; Austin, 1985:608; Staude, 1997:373, 383; Bousfield 2001:108.

The presence of two spines rather than one on the distal ends of the telson lobes are the primary feature distinguishing *Eogammarus oclairi* from *E. confervicolus*. Whether differences between *E. confervicolus* and *E. oclairi* are due to speciation or intraspecific allometric variation is unclear. The largest specimens in the collections (samples, 28.725x, 17.1229x, 12.1249x, 40.1252x) are mixed in with *E. confervicolus* morphotypes. This largest specimen has two stout distal spines on one telson lobe and one on the other. *Eogammarus oclairi* is thus a doubtful species.

Rammellogammarus sp. A

LCRANS, LIT

Specimen 1.1164x (female, Ft. Canby interior, 25 June 2003) has tiny pleonal spines that might be considered spines. This species occurs only in completely fresh water and appeared to be replaced by *E. confervicolus* occurred where salinities exceeded about 5 PSU. The possibility that this "new" *Rammellogammarus* is the long lost *Rammellogammarus ramellus* seems remote. (Weckel, 1907) reports *Rammellogammarus* (*Gammarus*) *ramellus* from Portland, Oregon. But also that: "These specimens were larger and stouter than those from California." Possibly Weckel misidentified his material and had *Rammellogammarus* sp. A. of this study. Either we did not find *Rammellogammarus ramellus* or Weckel's illustrations are misleading.

Rammellogammarus sp. A also does not appear to be *Rammellogammarus oregonensis* (Shoemaker, 1944) or *Rammellogammarus vancouverensis* Bousfield, 1979. Dorsal pleon spines of *E. sp. A* are 6-12 and variable in number and positioned on the extreme posterior edge of the pleonites. The pleonal spines of *R. oregonensis* and *R. vancouverensis* are clearly more anterior to the posterior pleonal edge than the spines of *R. sp. A*. Moreover, *R. ramellus* is without pleonal spines and *R. vancouverensis* have only 1-2 spines. Previous reports of *Rammellogammaurs ramellus*, *R. oregonensis* and *R. vancouverensis* from the Columbia River are probably in fact, *Rammellogammarus* sp. A. Only a single species seems to be involved whether it is a new species cannot be addressed here.

The eyes lack pigment and spines occur on the absolute posterior edges of pleonites 3, 2 and sometimes 1 which do not occur on *E. confervicolus*. Specimens 4.1085x (Gray's

Kingdom: Animalia
Phylum: Arthropoda
Subphylum: Crustacea

River Log scrape, 6/26/2003); specimens 3.1329x (Sportsmens Club boat launch, Kalama, 6/27/2002, sample 9.3). A large male with typical *Eogammarus* peg-spines was found among among specimens 7.1013x (Creek below Lewis & Clark Falls, Gravel bucket swirl, 26 June 2003). An ovigerous female, the largest specimen is in sample 6.1153x (Gray's River, bryozoans etc. scraped from a log, 26 June 2003). The posterodorsal pleonites of this specimen are lined only with setae but the remaining 25 specimens have stouter spines.

Distribution – Known only known freshwater reaches of the Columbia River.

Origins – Presumed native, if it is indeed a good species, due to its extreme similarity to the native *Rammellogammarus* species and *Eogammarus confervicolous*.

Rammellogammarus oregonensis (Shoemaker, 1944)

LIT

Origin: Native

Anisogammarus (Eogammarus) oregonensis - Shoemaker, 1944:89-93, figs.1-2; Barnard 1954a:13; Bousfield 1961:5; *Rammellogammarus oregonensis* - Bousfield, 1979:340-341; Austin, 1985:608; Bousfield 2001:108.

Bousfield (1979) reports *R. oregonensis* from Creeks and lakes of Lincoln and Lane Counties in Oregon and Lake Oswego (Bousfield 1979). However, this species was also not observed in the survey.

Distribution - A freshwater species that ranges from Eureka, California north to Cape Flattery, Washington.

AORIDAE

Grandidierella japonica Stephensen, 1938

LCRANS, LIT

Origin: Introduced

Grandidierella japonica - Stephensen, 1938:179-184, figs. 1-2; Ueno 1938:156; Nagata 1960:179. Pl. 17, fig. 103; Barnard, J.L. 1975:333(key), 360; Chapman & Dorman 1975:105-108, figs.1-4; Page & Stenzel 1975; Stenzel et al. 1976; Nichols 1977; Carlton, J.T. 1979a:127,144,146-147,152,179,192,662-663,866-868,880; Carlton, J.T. 1979b:433; Hirayama 1984a:15, figs. 53, 55, 56; Austin, 1985:614; Barnard & Karaman 1991:196; Ishimaru 1994:33-34; Greenstein, D.J. & L.L. Tiefenthaler, 1997:101-105; Muir, D.G. 1997:51; Staude, 1997:386; Smith et al. 1999:8-9, figs. 1, 3; Carlton, J.T., 1999:9; Chapman 2000:tab. 2; Bousfield 2001:112; Lowry & Stoddart 2003:71.

Distribution - Japan: Eastern coast of Japan, from Nakaminata, Honshu to southern Point of Kyushu, and southern coast of Korea between Pusan and Wando, including islands of Korea Strait. North America: Frasier River estuary, British Columbia, south to Bahia de San Quintin, Baja California, Mexico. Australia: Sydney, from Port Macquarie south to Cape Howe at New South Wales on the Victoria border. Europe: southern

Kingdom: Animalia
Phylum: Arthropoda
Subphylum: Crustacea

England. The Fraser River and English populations are at the highest latitudes any other populations of *Grandidierella* and far exceed the maximum latitude of the native *Grandidierella* populations of Japan. *Grandidierella japonica* is an estuarine species transferred around the world most likely with transplanted oysters and ballast water.

ATYLIDAE

Atylus tridens (Alderman, 1936)

LIT

Origin: Native

Nototropis tridens - Alderman, 1936:58-59, figs 20-25; *Atylus tridens* - Mills, 1961:25, fig.3; Barnard, J.L. 1966a:61; Barnard, J.L., 1975:340(key),346,359, fig.216; Klink, R.W. 1980:240; Austin, 1985:604; Staude, 1987:382, figs. 18.54, 18.63; Barnard & Karaman, 1991:265; Bousfield & Kendall, 1994a:10,20,22, fig. 9; Staude, 1997:361, 382, fig. 18.63; Bousfield 2001:97.

Distribution - Queen Charlotte Islands south along the outer coasts of British Columbia to Oregon and central California (Bousfield & Kendella 1994:22), 0-135m. *Atylus tridens* is an entirely marine species that is only likely to occur in the lower Columbia River incidentally

COROPHIIDAE

Americorophium brevis (Shoemaker, 1949)

LCRANS, LIT

Origins: Native

Corophium brevis - Shoemaker, 1949:70-72, fig.4; Barnard, J.L., 1954a:36; Barnard, J.L., 1975:340(key),359, figs.67,116,148,149; Otte, G., 1975:9, figs.4i-k,5g-I; Coyle & Mueller, 1981:9; Austin, 1985:615; Staude, 1987:349(key),386; Barnard & Karman, 1991:185; Staude, 1997:349, 386; *Americorophium brevis* - Bousfield & Hoover, 1997:90,92,95,97-98, fig. 17; Bousfield 2001:115.

Americorophium brevis is a predominately shallow water marine and high salinity estuary species that usually occurs in fouling communities and open coasts and marine bays.

Distribution - Prince William Sound, Alaska to San Francisco Bay, California, subtidal to 35 m (Bousfield & Hoover 1997:98).

Americorophium salmonis (Stimpson, 1857)

LCRANS, LIT

Origin: Native

Corophium salmonis - Stimpson, 1857:514-515; Stimpson 1857:74-75; Stebbing, 1906:692; Bradley 1908:235-241, pl.11, figs.20-27, pl.12, figs.28-35, pl.13, figs.38-39; MacGinitie 1935:700; Crawford 1937:603; Shoemaker, C.R. 1949:66-68, fig.1; Barnard, J.L., 1954a:36; J. Exp. Mar. Biol. Ecol. 1964:50; Otte, 1975:9(key), figs.4d-h,5d-f; Eckman 1979:437-457; Albright & Armstrong 1981:63 pp.; Wilson, S.L., D.L. Higley, & R.L. Holton 1981:273; Taghon 1982:295-304; Eckman 1983:241-257; Austin, 1985:615;

Kingdom: Animalia
Phylum: Arthropoda
Subphylum: Crustacea

Staude, 1987:349(key),386, fig.18.26; Barnard & Karaman, 1991:186; Staude, 1997:349, 386, fig. 18.26; *Americorophium salmonis* - Bousfield & Hoover, 1997:90,92,94, figs.14-15; Bousfield 2001:116.

Distribution - Its northeast Pacific range extends from south Alaska to Humboldt Bay, California. *Americorophium salmonis* is an endemic estuary species that has been introduced above the tidal range of the Columbia River dams and into Putah Creek, California. *Americorophium salmonis* does not attach its tubes to solid substratums and occurs exclusively on muddy to sandy bottoms in of estuaries, and slow moving rivers. Native to coastal regions and introduced inland.

Americorophium spinicorne (Stimpson, 1856c)

LCRANS, LIT

Origin: Native

(Stimpson, 1856c); *Corophium spinicorne* - Stimpson, 1856c:89; Stimpson, 1857:514; Bradley, 1908:227, pls.9-10; Essig 1925:189-190; MacGinitie 1935:700; Carl 1937:450; Crawford 1937:604; Shoemaker, 1949:74-76, fig.6; Barnard, J.L. 1952b:33; Barnard, J.L., 1954a:36-37; Bousfield 1958b:111; Filice 1958:184; Aldrich, 1961:21, fig.2; Bousfield 1961:2; Reish & Barnard 1967:16; Bousfield & J.D. Hubbard 1968:6; Eriksen 1968:1-12; Barnard, 1975:340(key),359, fig.141; Otte, G., 1975:9, figs.4a-c,5a-c; Siegfried, Kopache & Knight 1980:296; Austin, 1985:615; Staude, 1987:349(key); Barnard & Karaman, 1991:186; Staude, 1997:349, 386; *Americorophium spinicorne* - Bousfield & Hoover, 1997:90-93, fig.13; Bousfield 2001:115; Thorp & Covich 2001:780,785.

An endemic polyhaline species endemic to tidal bays, estuaries and freshwater river mouths of the northeast Pacific that ranges between Amchitka Island, Alaska to Morro Bay, California. *Americorophium spinicorne* has been introduced above the tidal range of the Columbia River (e.g., Thorp & Covich 2000) and up other rivers by human activities. *Americorophium spinicorne* occurs on fouling surfaces and mud bottoms in association with the NZMS. Native to coastal areas.

Monocorophium acherusicum (Costa, 1851)

LIT

Origin: Introduced

Podocerus cylindricus - Say 1818:387-388; Lucas 1842:232; Stebbing 1914:372-373; *Corophium cylindricum* - Smith 1873:566; Holmes 1905:521-522; Paulmier 1905:167, fig.37(in part); Holmes 1905:521-522, fig.; Johansen 1930:93; Cowles 1930:351; Shoemaker 1930a:128-129; Kunkel 1981:171-173, fig.52; (*Corophium cylindricus* ?Stebbing 1914:372-373;); *Audouinia acherusica* - Costa 1851:24; *Corophium contractum* - Thomson 1881:220-221, fig.9; *Corophium crassicorne* - Walker 1895:318; *Corophium bonellii* - Barnard, K.H., 1932:244; *Corophium acherusicum* - Costa 1853:178; Costa 1857:232; Bate 1862:282; Heller 1867:51-52, pl.4, fig.14; Della Valle 1893:367, pl.1, fig.2, pl.8, figs.17-18,20-41; Sowinsky 1897:9; Sowinsky 1898:455;

Kingdom: Animalia
Phylum: Arthropoda
Subphylum: Crustacea

Chevreaux 1900a:109; Graeffe 1902:20; Stebbing, 1906:692-740; Chevreaux 1911:271; Barnard, K.H., 1916:272-274; Stebbing 1917a:448; Ussing & Stephensen 1924:78-79; Chevreaux 1925c:271; Chevreaux & Fage 1925:368, fig.376; Chevreaux 1926:392; Cecchini 1928b:309-312, fig.1; Cecchini 1928e:8, pl.1, fig.6a; Schellenberg 1928:672; Miloslavskaya 1931:61(footnote); Schijfsma 1931a:22-25; Monod 1931a:499; Fage 1933:224; Candeias 1934:3; Shoemaker 1934c:24-25; Cecchini-Parenzan 1935:227-229, fig.52; Shoemaker 1935c:250; Crawford 1936:104; Schellenberg 1936c:21; Schijfsma 1936:122-123; Crawford 1937:617-620, fig.2; Crawford 1937a:650; Monod 1937:13; Miloslavskaya 1939:148-149; Barnard, K.H. 1940:482; Bassindale 1941:174; Stephensen 1944a:134; Shoemaker 1947:53, figs.2,3; Mohr & LeVeque 1948a; Shoemaker 1949a:76; Soika 1949:210-211; Gurjanova 1951:977-978, fig.680; Reid 1951:269; Stock & Bloklader 1952:4-5; Barnard, J.L. 1954a:36; Hurley 1954e:442-445, figs.35-39; Reish & Winter 1954; Barnard, J.L. 1955a:37; Irie 1957:5-6, fig.6; Barnard, J.L. 1958; Barnard, J.L. 1959b:58; Barnard, J.L. 1959c:38 (with references); Nayar 1959:43-44, pl.15, figs.14-20; Reish 1959b:39; Nagata 1960:177; Reish 1960:100-101; Barnard, J.L. 1961:173,175,182; Barnard, J.L. 1961:169,176; Reish 1961a; Reish 1961c; Jones, M.L. 1961:288; Reish 1963a; Reish 1963b; Barnard, J.L. 1964a:111, chart 5; Reish 1964b; Reish 1964c; Johnson & Juskevics 1965; Nagata 1965c:317; Painter 1966; Reish & Barnard 1967:12-13,16; Ledoyer 1968:214; Fearn-Wannan 1968b:134-135; Reish 1968b:49; Keith 1969; Mordhukai-Boltovskoi 1969:485, pl.25, fig.2; Sivaprakasam 1969d:156, fig.14; Bellan-Santini 1971:260-261; Barnard, J.L., 1971a:59; Reish 1971a; Stout 1971:68; Barnard, J.L., 1972b:48; Reish 1972:78; Bousfield, 1973:201, pl.62.2; Griffiths 1974a:181-182; Griffiths 1974b:228; Griffiths 1974c:281; Barnard, J.L. 1975:338-340(key),359, figs.143,144, 147; Chapman & Dorman 1975; Griffiths 1975:109; Otte, G., 1975:10, figs.6i-k,7d-f; Page & Stenzel 1975; Reish et al. 1975; Standing et al. 1975; Armstrong et al. 1976; Otte, 1976:8(key), figs. 6,7; Chapman 1978; Carlton, J.T. 1979:144-145,152,156,172,192,202,629,653-656,658,859-860,863-875,879; Carlton, 1979:655 (distribution list, Alaska to California); Klink, R.W. 1980:240; Hong, 1983:143-147, figs. 6-8; Hirayama, 1984:13, fig.50; Austin, 1985: 615; Staude, 1987:386, fig.18,27; Barnard & Karaman, 1991:185; Kim, 1991:114, fig.26; Ishimaru, 1994:35; Staude 1997:351, 386, fig. 18.16, 18.27; Carlton, J.T., 1999:9; *Monocorophium acherusicum* - Bousfield & Hoover 1997:111,112, 117,118,119, fig.30; Bousfield 2001:116; Lowry & Stoddart 2003:90. . . . and on and on(note - John Chapman) Distribution – One of the most widely distributed and reported medium to high salinity estuary organisms, *Monocorophium acherusicum* occurs in all large estuaries at all latitudes less than 50° (north or south).

In the northeast Pacific, its presence in central Alaska is not confirmed. However, it occurs in nearly every estuary from the Strait of Georgia to the Panama Canal and has likely been in the northeast Pacific for 200+ years. Not finding it in the lower Columbia River survey was a surprise.

CRANGONYCTIDAE

Reports of *Crangonyx floridanus* subgroup and *Crangonyx spp.* in the lower Columbia River are likely to comprise a single species, *Crangonyx pseudogracilis*.

Kingdom: Animalia
Phylum: Arthropoda
Subphylum: Crustacea

Crangonyx floridanus

LIT

Origin: Introduced – probably misidentified

Bousfield, E.L., 1963:2-6, figs.1-2; Barnard & Barnard 1983:434; Toft, J., T. Cordell, & C. Simenstad, 1999:35-36, tab.1, fig.1A; Bousfield 2001:101.

This Gulf coast species differs from *C. pseudogracilis* only by subtle, mostly microscopic characters that are seldom examined in routine synoptic surveys. The Columbia River records of this species are doubtful. More likely it is *Crangonyx pseudogracilis*.

Distribution - Gulf coast, sloughs, swamps, caves, and ponds, San Francisco Bay, California. Introduced into San Francisco Bay but doubtful in the lower Columbia.

Crangonyx pseudogracilis Bousfield 1958

LCRANS

Origin: Introduced

Melita parvimana- Holmes 1905; ?*Crangonyx gracilis* - Forbes 1876:6; Hynes 1955; ?*Melita parvimana* Holmes 1904:506, fig.; *Eucrangonyx gracilis* - Kunkel 1918:94, fig.20; Johansen 1920:128; Hubricht & Mackin 1940:199, fig.7; *Eucrangonyx gracilis* - Tattersall 1937:593; *Crangonyx gracilis* - Hubricht 1943:691; *Crangonyx pseudogracilis* Bousfield 1958:102-105, fig.17; Mills 1964a:4-5; Bousfield 1973:68-69, pl 8.1; Holmes 1975; Gledhill et al. 1976; Thomas, J.D. 1976:90; Barnard & Barnard 1983:435; Austin, 1985:597; Pinkster et al. 1992; Costello 1993:292; Bousfield 2001:101.

Bousfield (1963) described *C. pseudogracilis* from the Napanee River, Ontario and from other material from Quebec, Vermont and Missouri. In the same paper, Bousfield reports the introduction of *C. pseudogracilis* to the British Isles based on specimens from Gloucestershire, England. Bousfield (1958:105) further reports that Holme's (1905:94, fig.) "*Melita parvimana*" from Connecticut is "unquestionably a *Crangonyx* and very probably a pseudocrangonyx". Bousfield (1958) distinguishes *C. pseudogracilis* from the superficially similar *C. gracilis* Smith 1871, "hence the specific name."

Crangonyx pseudogracilis "breeds in spring and throughout the summer" Bousfield (1958) and is frequently taken "along with *Gammarus fasciatus* and *Hyalella azteca*, though less often with *C. gracilis* and *G. pseudolimnaeus* (in northern areas)". The distribution and ecology of the species is "rivers, river mouths, lakes, sloughs, quarry ponds, dams, and other larger freshwaters that tend to be somewhat turbid and warm in summer".

The combination of bifid spines lining the palm, and singly inserted simple setae on lateral anterior edge of the propodus of female gnathopod 1 place specimens 31.503x, outside of Holsinger (1972) couplet 1. However, Bousfield (1958, fig. 17) clearly indicates the presence of these characters. Bousfield's (1973, pl. VIII) illustration of *P. pseudocrangonyx*. Figure 2A of *Crangonyx floridanus* from San Francisco Bay

Kingdom: Animalia
Phylum: Arthropoda
Subphylum: Crustacea

Toft et al. (2002) is an unpublished illustration of *Crangonyx forbesi* (Hubricht and Mackin 1940) from the Subterranean Amphipod Database (http://web.odu.edu/sci/biology/amphipod/cc_pictu.htm).

The combination of comb setae lining the dorsal lateral edge of the outer ramus of male uropod 2, and special ventral spines on the inner margin of the outer ramus of male uropod 2 (unique among species of *Crangonyx*) distinguish this species from all others (Zhang 1998). However, Zhang's illustrations of *C. floridanus* and *C. pseudogracilis* indicate that morphological differences are subtle if they are real.

Distribution - Introduced to Great Britain and Ireland (Costello 1993), NW and NE North America, Oregon. Inhabits aquatic vegetation in still an slow flowing waters, including organically polluted and saline waters (Holmes 1975, Gledhill et al. 1976, Pinkster *et al.* 1992, Costello 1993). It clings to plants when removed from water and is thus further distributed in Ireland (O'Connor et al. 1991).

HAUSTORIIDAE

Eohaustorius brevicuspis Bosworth, 1973

LCRANS

Origin: Native

Eohaustorius brevicuspis - Bosworth, 1973:255, 257, 259, fig. 1k-o, fig. 2b, f, n; Austin, 1985:605; Staude, 1987:383,372(key); Barnard & Karaman, 1991:363; Bousfield & Hoover 1995:50, fig.10; Staude, 1997:372, 383, fig. 18.11; Bousfield 2001:107.

Samples 11.1389x, (Baker Bay, Fort Columbia Tide flats, 11 June 2002), specimens do not have a cusp on the dorsal posterior of basis of pereopod 7 and pereopod 6 have only a single seta on the lateral faces of articles 5 and 6. These differences are consistent and suggest that these populations are a new species. However, a single individual of specimens 7.993x (Sand Island, Outer Beach, High Intertidal 25 June 2003) has the dorsal cusp and all specimens have two or more setae on the lateral faces of articles 5 and 6 of pereopod 6. Size, instar, age, seasonal differences in morphology should be examined in these species.

Distribution - Central California north to the Strait of Juan de Fuca (Bousfield & Hoover 1995:50) in high beach pools, river mouths, and estuaries in clean sand, 0-1 m.

Eohaustorius estuarius Bosworth 1973

LCRANS, LIT

Origin: Native

Eohaustorius estuarius - Bosworth, 1973:257-258, 259, figs. 2c, g, i-m; Austin, 1985:607; Staude, 1987:372(key),383, fig. 18.11; Barnard & Karaman, 1991:363; Bousfield & Hoover 1995:40,41,42, fig.4; Staude, 1997:372, 383; Bousfield 2001:107. Distribution - Occurring in clean sand areas of estuaries and freshwater seeps and is very abundant in sandy areas of the lower Columbia River. Does not occur in completely

Kingdom: Animalia
Phylum: Arthropoda
Subphylum: Crustacea

fresh water. Sample 120.991 (inside Coast Guard Jetty 25 June 2002). *Eohaustorius estuarius* was the most abundant *Eohaustorius* and the only species other than *E. brevicuspis* encountered in the LCR. Since *Eohaustorius* are difficult to distinguish, the other species identified previously from the LCR, *Eohaustorius sawyeri* and *Eohaustorius washingtonianus* are more likely to be *E. estuaries*. Distribution - Oregon, Eureka, California north to Cape Flattery, Washington, 0-7m.

Monoporeia sp.

Syn: Previously misidentified as *Pontoporeia affinis*.

LIT

Origin: Cryptogenic

The Columbia River population is the only population of the genus reported south of Alaska. This disjunct distribution has all appearances of a cold-water introduction, which would be unique among NE Pacific amphipods. However, the rapidly evolving state of the taxonomy of pontoporeiids prevents a definitive identification of this species presently. This species is reported only from lower Columbia River, however, Jeff Cordell has seen it in other adjacent estuaries.

HYALELLIDAE

Hyalella azteca (Saussure, 1858)

LCRANS, LIT

Origin: Cryptogenic

Amphithoe aztecus - Saussure 1858:474; *Allorchestes knickerbockeri* - Bate 1862:250; *Hyalella dentata* - Smith 1874:609, fig.1; *Lockingtonia fluvialis* - Harford 1877:54; *Hyalella knickerbockeri* - Weckel 1907:54, fig.15; *Hyalella Hyalella azteca* - Bousfield 1996:183; Bousfield 2001:104; *Hyalella azteca* - Stebbing 1906:575; Stout, V.R. 1913:635; Saunders 1933:245, fig.1; Shoemaker 1942b:80,82; Bulycheva 1957:181, figs.66a-b; Bousfield 1958b:109, fig.20; Bousfield 1961:5; Bousfield 1973:154, pl 43.2; Thomas, J.D. 1976:91-92; Barnard & Barnard 1983:708; Austin, 1985:595; Bousfield 1996:206, 207-209, figs. 3, 12, 17E; Hendrycks & Bousfield 2001:28, figs.4-5,6a,14; Bousfield 2001:104; Gonzalez & Watling 2002:173-183, figs. 1-5.

Specimen from 7.767x Carroll's Channel Log raft, 26 June 2002 has particularly prominent dorsal carina on pleonites. This distinctive species, or species complex, has eluded taxonomist for 150 years and I am unable to resolve it here. Its very broad geographic distribution and many associations with introduced species, including introduced aquatic plants, suggest the almost certain possibility that populations have been moved about. However, the existence of many species within this complex is also likely. Resolution of the evolutionary origins of these populations and the role of humans in their distributions is not yet possible. Figure 2A of *Hyalella* from San Francisco Bay Toft *et al.* (2002) is an illustration from (Cole & Watkins 1977) of a specimen from Montezuma Well, Yavapai Co., Arizona.

Kingdom: Animalia
Phylum: Arthropoda
Subphylum: Crustacea

Distributions - Fresh waters of north and central America and Caribbean islands north to the tree line of North America and in larger rivers seaward into tidal fresh waters, and fresh-water barrier beach lagoons (Bousfield 1973:154) and freshwater and slightly brackish waters of lakes, rivers, upper estuaries of Mexico and California north to Alaska (Hendrycks & Bousfield 2001:28).

Hyalidae

Allorchestes angusta Dana, 1856

LIT

Origin: Native

Allorchestes angustus - Dana 1856:177; Barnard 1952:20-23, pl. 5, figs. 2-6; *Allorchestes angusta* - Barnard 1974:42; Barnard, J.L. 1975:343(key),358; Barnard 1979:91, figs. 50-52 (part); Bousfield 1981:81, figs. 12, 13; Bousfield 1996:178, fig. 1; Barnard; Bousfield 1996:181; Hendrycks & Bousfield 2001:10, 24-25, 1-6h, 12; non *Allorchestes angustus* - Barnard 1954c:21-23, Pl. 21 (=A. *bellabella*); *Allorchestes oculatus* - Stout 1913:651?
Distribution - Japan northward through Kuriles, across Aleutian Chain to Alaska then southward to California, generally intertidal, phycophilous, rarely subtidal (Barnard, 1979), high rocky intertidal and among algae wrack in protected bays and high salinity estuaries, 0-4m

ISAEIDAE

Photis macinerneyi Conlan, 1983

LIT

Origin: Native

Conlan, 1983:54, fig.27; Austin, 1985:612; Barnard & Karaman, 1991:226; Staude, 1997:351, 385, fig. 18.33; Bousfield 2001:114; Cadien 2001:98.

Distribution - Lady Ellen Point, Broughton Strait, Vancouver Island south to Neah Bay, Clallam County, Washington (Conlan 1983) and southern California (Cadien, 2001), 0-45m. A probable incidental species in the LCR.

LYSIANASSIDAE

Hippomedon columbianus Jarrett and Bousfield, 1982

LIT

Origin: Native

Hippomedon denticulatus - Barnard 1954:4, pls. 2,3 (in part) not Bate 1857; Hurley 1963:137-140, fig.45; Barnard 1971:31-34, fig.21(form with gaped gnathopod 2);

Hippomedon columbianus - Jarrett & Bousfield 1982:109-111, fig.3; Barnard & Karaman, 1991:490; Bousfield 2001:76; Cadien 2001:94.

Distribution - Oregon, 100-150m (Barnard 1971:34), British Columbia (Jarrett & Bousfield 1982), Southern California (Cadien 2001), 4-320m, probably incidental marine species of the LCR.

Kingdom: Animalia
Phylum: Arthropoda
Subphylum: Crustacea

MELITIDAE

Melita cf. nitida (Smith, 1874)

LCRANS

Origin: Introduced

Melita nitida - Smith, 1874; Shoemaker, C.R., 1935b:70-71, fig.2; Light 1941:180-190; ?Barnard 1954f:161; Mills 1964a:5-7, fig. 1; Bousfield, 1973:65, pl.9.2; Barnard, J.L. 1975:361; Chapman & Dorman 1975; Levings & McDaniel, 1976:5?; Thomas, J.D. 1976:90-91; Chapman, C.J., 1977:101; Carlton, J.T. 1979a:120,146-147,192,672-673,859,868-869,877; Carlton 1979b:433; Sheridan 1980:61-62, Figs. 1-2; Barnard, J.L. & C.M. Barnard, 1983:665; Ortiz 1983:26; Austin, 1985:610(part); Chapman, J.W. 1988:372-374, fig.5F; Jarrett & Bousfield 1996:51,57,59, figs.35,36; Carlton, J.T., 1999:9; Bousfield 2001:110; Faasse & Moorsel 2003:16-18, figs. 1&2, tabs. 1&2; *Melita sp.* - Light 1941:180; *Melita sp.A* - Barnard 1975:361; ?*Melita setiflagella* - Yamato, 1988:80-86, figs. 2-6; ? Kim et al. 1992b:116, 119, fig. 3; Jarrett & Bousfield 1996:51,61, fig.38c; non *Melita nitida* - Shoemaker, 1935:70, fig.2.

This is the first likely record of *M. nitida* from the Columbia River. The two damaged specimens, one male and one female are similar to *M. nitida* in the nearly bare posterior urosome, quadrate epimeron and general shape of male gnathopod 1, but the female coxa 5 does not have the extended posterior that appears to be a stridulating organ on *M. nitida* s.s.

Distribution - Southern British Columbia and northern Washington, also in Columbia estuary, parts of San Francisco Bay and south of Point Conception, in summer-warm brackish localities (Jarrett & Bousfield 1996). North-western Atlantic distribution is from New England to at least the southern Gulf of Mexico, 0-20m. It may also occur in Japan if *Melita setiflagella* Yamato, 1988 proves to be a junior synonym.

OEDICEROTIDAE

Americhelidium shoemakeri (Mills, 1962)

LIT

Origin: Native

Synchelidium shoemakeri - Staude, 1997:362, 368; *Synchelidium shoemakeri* - Mills 1962:15-17, figs.4, 6A; Barnard, J.L. 1966a:79; Barnard, J.L. 1966b:27; Barnard, 1969a:195; Barnard, J.L. 1971b:51; Barnard, 1975:345(key), 363, fig.136; Klink, R.W. 1980:246; Austin, 1985:591; Staude, 1987:378; Barnard & Karaman, 1991:566; Thomas & McCann 1997:57, fig.2,36; *Americhelidium shoemakeri* - Bousfield & Chevrier 1996:132-134, fig.37; Bousfield 2001:91.

Americhelidium shoemakeri occurs in full marine sandy sediments from British Columbia to southern California and in the intertidal to 183m (Thomas & McCann 1997). Its occurrence within the LOWER COLUMBIA RIVER is likely to be incidental.

Kingdom: Animalia
Phylum: Arthropoda
Subphylum: Crustacea

Pacifoculodes spinipes (Mills, 1962)

LIT

Origin: Native

Monoculodes spinipes - Mills, 1962:12-14, fig.3,6C; Barnard, J.L., 1962e:368-369, fig.10; Barnard, J.L., 1966b:26; Barnard, J.L., 1971b:51; Klink, R.W., 1980:246; Austin, 1985:591; Staude, 1987:378; Barnard & Karaman, 1991:560; Staude, 1997:362, 378; Thomas & McCann, 1997:55-56, fig.2.34; *Pacifoculodes spinipes* - Bousfield & Chevrier, 1996:103-104, fig.16; Bousfield 2001:92; non *Monoculodes spinipes* - Mills, cf. Barnard, J.L., 1962:368, fig.10.

Distribution - British Columbia to southern California, intertidal to 98m (Thomas & McCann 1997); North-eastern Pacific boreal, 0-50m. Occurrences of thus fully marine species in the LOWER COLUMBIA RIVER are probably incidental.

PHOXOCEPHALIDAE

Grandifoxus grandis (Stimpson, 1856)

LCRANS, LIT

Origin: Native

Phoxus grandis - Stimpson, 1856:90; Stimpson, 1857: 81-82; Stimpson, 1857:521-522; *Pontharpinia grandis* - Stebbing, 1906: 147; *Pontharpinia milleri* - Thorsteinson, 1941:82, pl. 5, figs.52-62; *Paraphoxus milleri* - Barnard, J.L., 1958:147; Barnard, J.L., 1960:266, pl 40; Barnard, J.L., 1975:362; *Pontharpinia longirostris* - Gurjanova 1938:263-267,385, fig.7; Gurjanova 1951:385-387, fig.235; *Pontharpinia robusta* - Gurjanova 1938:262-263, fig.6a; Gurjanova 1951:384-385, figs.233-234; *Grandifoxus grandis* - Barnard, J.L., 1979:375; Barnard, J.L., 1980b:495-500, fig.1 upper right; Coyle, 1982:449, fig. 10 g, h; Austin, 1985:597; Barnard & Karaman, 1991:611; Jarrett & Bousfield, 1994a:63,64,67,68, fig. 1; Staude, 1997:363, 380, 503; Bousfield 2001:86. Distribution – The range of *Grandifoxus grandis* is Dixon Entrance, Alaska to Pacific Grove, California, often occurring in reduced or brackish salinities (Jarrett & Bousfield 1994a:67) at depths of 0-1m. *Grandifoxus grandis* may permanently reside in the LCR.

Mandibulophoxus gilesi Barnard, J.L., 1957

LIT

Origin: Native

Mandibulophoxus uncistrostratus - Barnard 1960a:359; Barnard 1969a:196;

Mandibulophoxus gilesi - Barnard 1957a:433-435, figs. 1-2; Gray & McCain, 1969:189, fig.1; Barnard 1975:348(key),361; Barnard, J.L. & Drummond, 1978:91(key); Barnard & Karaman, 1991:620; Jarrett & Bousfield 1994b:78,80, figs.3,4; Klink, R.W. 1980:247; Staude, 1997:363, 380, 503; Bousfield 2001:87.

Distribution - Central British Columbia to southern California, intertidal to shallow subtidal depths and subtidally in substrata exposed to tidal currents (Jarrett & Bousfield 1994b:80) boreal, 0-14m

Foxiphalus obtusidens (Alderman, 1936)

Kingdom: Animalia
Phylum: Arthropoda
Subphylum: Crustacea

LIT

Origin: Native

Pontharpinia obtusidens - Alderman 1936:54-56, figs.1-13,19; Hewatt 1946:199; Barnard, J.L. 1954a:4; *Parapinia sic pontarpioides* - Gurjaonva 1953:229; *Paraphoxus obtusidens* - Barnard, J.L., 1958a:147; Barnard, J.L., 1960:249-259, pl.33-37; Barnard, J.L. 1964a:105, chart 6; Barnard, J.L. 1964c:244; Barnard, J.L., 1966a:89; Barnard, J.L., 1966b:29; Barnard, J.L. 1969a:197; Barnard, J.L., 1970b:3; Barnard, J.L. 1971b:70; Barnard, J.L., 1975:362, pl. 72(22); *Foxiphalus obtusidens* - Barnard, J.L., 1979a:373; Klink, R.W., 1980:247; Barnard, J.L. & C.M. Barnard, 1982b:4-12, fig.1(part); Austin, 1985:597; Barnard & Karaman, 1991:610; Jarrett & Bousfield, 1994a:63,93,94; Staude, 1997:364, 380, 503; Thomas & McCann, 1997:78-79, figs.2.53, 2.59; Bousfield 2001:86; not - *Paraphoxus obtusidens major* - Barnard, J. L 1960:259-261, pl.32.
Distribution - Kuril Islands, Okhotsk Sea, Alaska, common from California to British Columbia in sandy tidepools, 0-210 m (Thomas & McCann 1997:79), 0-210, ?459m.

Rhepoxynius abronius (Barnard, J.L. 1960)

LIT

Origin: Native

Paraphoxus abronius - Barnard, 1960:203, pl.5; Barnard, J.L. 1966a:88; *Rhepoxynius abronius* - Barnard & Barnard, 1982a:26; Slattery, P.N., 1985:635--647; Robinson, A.M., et al. 1988:953-958, tables 1-4; Bousfield, 1990:13; Barnard & Karaman, 1991:629; Bousfield, 1991:84; Jarrett & Bousfield 1994a:63,108,109-110, fig. 21; Staude, 1997:363, 380, 506; Bousfield 2001:86.
Distribution - Queen Charlotte Islands southward to California, commonly and abundantly inshore and sub-tidally, mostly at surf-protected localities, in sand of protected bays and shorelines to below 50 m (Jarrett & Bousfield 1994a); San Diego: Point Conception, southern California south to Ensenada, Baja California, Bathyal, 9-274 m.

Rhepoxynius daboius (Barnard 1960)

LIT

Origin: Native

Paraphoxus daboius - Barnard 1960a:210-212, pls.10-11; Barnard 1966a:88; Barnard 1971b:70; *Rhepoxynius daboius* - Barnard 1979:372; Klink, R.W., 1980:248; Barnard, J.L. & C.M. Barnard, 1982a:30-32; Austin, 1985:599; Barnard & Karaman, 1991:629; Jarrett & Bousfield, 1994a:63,108,122, fig.26; Staude, 1997:363, 380; Thomas & McCann, 1997:95-96; Bousfield 2001:86.
Distribution - Alaska to southern California, 77-813m. An entirely marine species possibly incidental or misidentified in the LCR.

Rhepoxynius heterocuspидatus (Barnard, J.L., 1960)

LIT

Origin: Native

Kingdom: Animalia
Phylum: Arthropoda
Subphylum: Crustacea

Paraphoxus heterocuspoidatus - Barnard 1960:224-226, pls.19-20; Barnard 1964a:103,105; Barnard 1966a:89; Barnard 1969a:196-197; *Rhepoxynius heterocuspoidatus* - Barnard 1979:372; Barnard, J.L. & C.M. Barnard, 1982a:38-42, fig.4 (part); Austin, 1985:599; Barnard & Karaman, 1991:629; Jarrett & Bousfield 1994a:63; Staude, 1997:364, 380; Thomas & McCann, 1997:96, fig.2.75; Bousfield 2001:87. Distribution - Point Conception, California to Bahia de Los Angeles, Baja California, 0-146m (Thomas & McCann 1997). Occurrences of *R. heterocuspoidatus* in the LOWER COLUMBIA RIVER are doubtful.

Rhepoxynius tridentatus (Barnard 1954)

LIT

Origin: Native

Pontharpinia tridentata - Barnard, J.L. 1954a:4-6, pls.4-5; *Paraphoxus tridentatus* - Barnard, J.L. 1960a:261-265, pls.38-39; Barnard, J.L. 1966a:90; Barnard, J.L. 1969b:224; Barnard 1975:362; *Rhepoxynius tridentatus* - Barnard 1979:372; Klink, R.W. 1980:248; Barnard & C.M. Barnard, 1982a:42-44, fig. 6 bP7; Austin, 1985:599; Barnard & Karaman, 1991:629; Jarrett & Bousfield, 1994a:63,108,110; Staude, 1997:363, 380; Bousfield 2001:87; not: *Paraphoxus tridentatus pallidus* - Barnard 1960:261, pls. 38-39; not: *Paraphoxus heterocuspoidatus* - Barnard 1960:224, pls.19-20. Distribution - Puget Sound, Washington to vicinity of Point Conception, California, 0-89m (Barnard & Barnard 1982a). Occurrences of *R. tridentatus* in the LOWER COLUMBIA RIVER are probably incidental.

TALITRIDAE

Megalorchestia pugettensis (Dana, 1853)

LCRANS

Origin: Native

Orchestia (Talitrus) pugettensis - Dana, 1853&1855:859, t.57, fig.3a-d; Stimpson 1857:516; *Orchestoidea pugettensis* - Thorsteinson, 1941:pl.1, figs.1-9; Bousfield 1958:890, fig.2a,10i; Bousfield, 1961:7, fig.3; Bowers, 1963:317, figs.3e,4; Bousfield, 1975:355,364, fig.232; Bowers, 1975:357, fig.228; Staude et al., 1977:12, fig.20a; Klink, R.W., 1980:249; Bousfield, 1981:fig.18; *Orchestoidea corniculata* - Thorsteinson, 1941:55; *Talorchestia tridentata* - Stebbing, 1899:398, t.30b(male); *Megalorchestia pugettensis* - Bousfield, 1982b:37-38, fig.16; Austin, 1985:596; Staude, 1997:353, 355, 380; Bousfield 2001:106; Not - *O. Pugettensis* - Stebbing 1906a:528.

Distribution - Southern Alaska to central California, high intertidal coastal beaches and estuaries in high salinities and brackish water areas.

Traskorchestia traskiana (Stimpson, 1854)

LCRANS

Origin: Native

Kingdom: Animalia
Phylum: Arthropoda
Subphylum: Crustacea

Orchestia traskiana - Stimpson 1854:98; Stimpson 1856:90; Stimpson 1957:517-518; Bate, S. 1862:19, pl.3, fig.4; Stebbing 1906a:534; Stout 1912:134, figs.74-75; Stout, V.R. 1913:635; Thorsteinson, E.D. 1941:54-55, pl.1, figs.1-9; Shoemaker 1942:13; Barnard, J.L. 1952b:23; Barnard, J.L., 1954a:23; Bousfield 1958a:885-887, figs.2d,10d; Bousfield 1961:3, fig.1-2; Barnard 1964a:116; Bousfield 1975:363, fig.236; Klink, R.W. 1980:249; Bousfield 1981:83, fig.17; *Orchestia sp.* - O'Clair 1977:446; *Traskorchestia traskiana* - Bousfield 1982b:10-13, fig.5; Staude, 1997:355, 380; Bousfield 2001:105; not *Orchestia taskiana* - Bulychева 1957:166, fig.60.

Distribution - Amchitka Island, Alaska to Bahia de San Quintin, Baja California.

Phylum: Arthropoda

Subphylum: Crustacea

Class: Malacostraca

Peracarida – Mysida

Section write up by John Chapman

Mysids are integral components of nearshore, estuary and freshwater food-webs of western North America both as predators of and food for many commercially and recreationally important fishes. Mysids, being peracaridan crustaceans, brood their eggs in a brood pouch (thus the vernacular name "opossum shrimp"). The brood pouch is formed by inner lamellae extending from the walking legs and the hatched young emerge from the pouch after they are fully formed. And the young emerge fully formed. Although half of all mysid species in San Francisco Bay are introduced, no introduced mysids were found in this survey of the lower Columbia River.

Mysidae		
<i>Acanthomysis macropsis</i>	LIT	Native
<i>Archaeomysis grebnitzkii</i>	LIT	Native
<i>Exacanthomysis</i> spp.	LIT	
<i>Neomysis integer</i>	LIT	Native
<i>Neomysis kadiakensis</i>	LIT	Native
<i>Neomysis mercedis</i>	LCRANS, LIT	Native
<i>Neomysis rayii</i>	LIT	Native
<i>Neomysis</i> spp.	LIT	

Acanthomysis macropsis (Tattersall, 1932)

LIT

Alenacanthomysis macropsis Tattersall 1932; *Neomysis macropsis* Tattersall 1932; Li 1936; *Alienacanthomysis macropsis* Tattersall 1932; Holmquist 1981; Daly & Holmquist 1986:1208.

Distribution – California to Alaska in shallow water among eelgrass and algae, "not uncommon".

Origin: Native.

Kingdom: Animalia
Phylum: Arthropoda
Subphylum: Crustacea

Archaeomysis grebnitzkii Czerniasky, 1882

LIT

Callomysiss maculata - Holmes 1894; non *Archaeomysis maculata* - (Holmes 1894); Tattersall 1932 (see Holmquist 1975).

Distribution – California to western Alaska and Japan. Intertidal, common to abundant in open coastal, unprotected inland waters above sandy bottoms, uncommon in brackish waters.

Origins: native.

Neomysis integer (Leach, 1815)

LIT

Neomysis integer, *Praunus integer*, *Neomysis vulgaris*, *Mysis scoticus* - See Gordan 1957:367-368 for synonymy.

Neomysis integer is a dominant mysid shrimp in the upper reaches of estuaries in Europe where it occurs in non-tidal lagoons, isolated bodies of nearly freshwater, and in high shore hypersaline pools, but is rare in fully marine habitats. There are no other reports of this species in the lower Columbia. This record is either an unrecognized introduction into the lower Columbia River or a misidentification.

Origins: Introduced (not seen)

Neomysis awatchensis see *N. mercedis*

LIT

Neomysis kadiakensis Ortmann, 1908

LIT

Tattersall 1951:192-194; Gordan 1957:368; Daly & Holmquist 1986:1209; Kathman et al. 1986:202-203, fig.

Distribution – southern Alaska to southern California, neritic, to 200 m.

Origin: Native (not seen):

Neomysis mercedis Holmes, 1897

LCRANS, LIT

Neomysis awatchensis – Brandt 1851; Tattersall 1951:190-192; Banner 1954; Gordan 1957:366-367; *Neomysis mercedis* – Holmes 1897:199; Holmes 1900:222; Tattersall 1932b:318; Tattersall 1933:11; Scheffer & Robinson 1939:135; Banner 1948b:75; Tattersall 1951:187; Pennak 1953:323, 422; Smith et al. 1954:136; Gordan 1957:368; Daly & Holmquist 1986:1209; Kathman et al. 1986:204-205, fig.

Distribution - Southern Alaska to southern California, euryhaline, fresh to marine, littoral and shallow neritic marine waters.

A few 15 mm specimens (the greatest length observed), had a pointed antennal scale, quadrangular or rounded rostrum, widely separated spines of the telson of both sexes, and by the 4th male pleopod which has a short terminal article of the exopod (less than 1/3 length first article) and the thick, short lateral extension of the endopod.

Origin: Native

Kingdom: Animalia
Phylum: Arthropoda
Subphylum: Crustacea

Neomysis rayii

LIT

Tattersall 1951:181-186, figs. 68-71; Gordan 1957:368-369; Daly & Holmquist 1986:1209; Kathman et al. 1986:206-207, fig.

A probable misidentification or incidental species in the LCR.

Distribution – Kamchatka Peninsula, Russia and central California to northern Alaska, neritic to 300 m.

Origins: native.

Phylum: Arthropoda

Subphylum: Crustacea

Class: Malacostraca

Decapoda

Section write up by John Chapman

Decapoda have ten feet and a carapace that covers the united head and thorax. Decapoda are the most important crustacean food source for humans. *Cancer magister* is the most important commercially fished crustacean in the northeast Pacific. Decapod juveniles hatch out of the booded eggs at the nauplius stage (except in penaeid shrimps) and undergo extended larval dispersal before metamorphosis and settling back to the benthos.

* - unsuccessful introduction

Astacidae		
<i>Pacifastacus leniusculus klamathensis</i>	LIT	Native
<i>Pacifastacus leniusculus trowbridgii</i>	LIT	Native
<i>Pacifastacus leniusculus leniusculus</i>	LCRANS, LIT	Native
Callinassidae		
<i>Neotrypaea californiensis</i>	LCRANS, LIT	Native
Canceridae		
<i>Cancer magister</i>	LIT	Native
<i>Cancer oregonensis</i>	LIT	Native
<i>Cancer</i> spp.	LIT	
Crangonidae		
<i>Crangon franciscorum franciscorum</i>	LCRANS, LIT	Native
<i>Crangon nigromaculata</i>	LIT	Native
<i>Crangon</i> spp.	LIT	
<i>Lissocrangon stylirostris</i>	LIT	Native
<i>Neocrangon alaskensis</i>	LIT	Native
Grapsidae		
<i>Eriocheir japonica</i>	LIT	Introduced*
<i>Hemigrapsus oregonensis</i>	LCRANS, LIT	Native
Hippolytidae		

Kingdom: Animalia
 Phylum: Arthropoda
 Subphylum: Crustacea

<i>Heptacarpus brevirostris</i>	LIT	Native
Nephropidae		
<i>Homerus americanus</i>	LIT	Introduced*
Palaemonidae		
<i>Exopalaemon modestus</i>	LCRANS, LIT	Introduced
Porcellanidae	LIT	
Upogebiidae		
<i>Upogebia pugettensis</i>	LIT	Native

ASTACIDAE

Pacifastacus leniusculus klamathensis (Stimpson 1857a)

LIT

Hobbs 1989:7, fig. 6 (with synonymy)

Distribution – British Columbia and Idaho south to central California, in cold, swift streams.

Origins: Native.

Pacifastacus leniusculus trowbridgii (Dana, 1852)

LIT

Hobbs 1989:7-8, fig. 5 (with synonymy)

Distribution – In North America: British Columbia, California, Idaho, Nevada, Oregon, Utah, and Washington, in streams and lakes. Introduced to Sweden (Svardson 1965:92) and Japan (Kamita 1970:140).

Origins: Native.

Pacifastacus leniusculus leniusculus (Dana, 1852)

LCRANS, LIT

Hobbs 1989:7, fig. 6 (with synonymy)

Distribution – British Columbia and Idaho south to central California, in cold, swift streams.

Origins: Native.

NEPHROPIDAE

Homerus americanus Milne-Edwards, 1837

LIT

Many unsuccessful introductions have been attempted in the region since 1874 without success Carlton 1979:691-695.

Not established*

THALASSINIDEA

CALLIANASSIDAE

Callianassidae see *Neotrypaea*

Neotrypaea californiensis (Dana, 1854)

Kingdom: Animalia
Phylum: Arthropoda
Subphylum: Crustacea

LCRANS, LIT

Callianassa californiensis – Dana 1854; Hart 1982:58, 60, fig. 15; *Neotrypaea californiensis* - Manning & Felder 1991:771, fig. 10 (with synonymy); Jensen 1995:43, 78, fig. 158.

Distribution – Mutiny Bay, Alaska to Punta Banda, Baja California, 0 – 50 m.

Origins: Native.

Manning, R. B. and D. L. Felder 1991. Revision of the American Callianassidae (Crustacea: Decapoda: Thalassinidea), Proceedings of the Biological Society of Washington 104-764-792.

UPOGEBIIDAE

Upogebia pugettensis (Dana, 1852)

LIT

Upogebia pugettensis - Schmitt 1921:115-116, fig. 77 (with synonymy); Williams 1986 (with synonymy); Hart 1982:52-53, fig. 12; Jensen 1995:43, 78, fig. 160.

Distribution – Valdez Narrows, Alaska to Morrow Bay, California.

Origins: native.

Williams, A. B. 1986. Mud shrimps, *Upogebia*, from the eastern Pacific (Thalassinidea: Upogebiidae), San Diego Natural History, Memoir 14:1-60.

BRACHYURA

CANCRIDAE

Cancer magister Dana, 1852

LIT

Hart 1982:23, 33, 34, 212, fig. 87; Jensen 1995:14, 27, 28, fig. 31.

Distribution – Pribilof Islands, Alaska to Santa Barbara, California, 0 – 179 m.

Origin: Native

Cancer oregonensis (Dana, 1852)

LIT

Cancer oregonensis - Schmitt 1921:234-235, Pl. 36, figs. 3-4 (with synonymy); Hart 1982:23, 33, 34, 210, fig. 87; Jensen 1995:36, fig. 29.

Distribution – Bering Sea to Santa Barbara, California, 0- 436 m.

Origin: native

CARIDEA

CRANGONIDAE

Crangon franciscorum franciscorum Stimpson, 1856

LCRANS, LIT

Holthuis 1980:150 (with citations); Butler 1980:101-102, 107; Jensen 1995:40-41, fig. 57.

Specimen #1120x, Ft. Canby, Jetty exterior, 25 June 2003, has C.

nigricauda antenna scale and palm of leg 1 but short tooth on abdominal segment 5 rather than prominent long tooth.

Kingdom: Animalia
Phylum: Arthropoda
Subphylum: Crustacea

Distribution – Resurrection Bay, Alaska to San Diego, California, intertidal to 91 m.
Crangon franciscorum franciscorum is the estuarine form that occasionally occurs in nearly fresh water and was once the basis of a commercial fishery in San Francisco Bay. This was the only form found in the survey.

Origin: Native

Crangon nigromaculata Stimpson, 1856

LIT

Holthuis 1980:150-151 (with synonymy); Butler 1980:95, 102; Jensen 1995:41, fig. 60.
Crangon nigromaculata was once fished commercially along with *Crangon franciscorum franciscorum* in San Francisco Bay (Holthuis 1980).

Distribution – Northern California to Baja California, sand bottoms 5 – 174 m. Probably not correctly identified in the LCR.

Origins: Native.

Lissocrangon stylirostris (Holmes 1900)

LIT

Crangon stylorostris – Butler 1980:98-99, fig.; Jensen 1995:41, fig. 61; *Lissocrangon stylirostris* – Kuris & Carlton 1977:551-552.

Distribution – Chirikov Island, Alaska to San Louis Obispo Bay, California.

Origins: native.

Neocrangon alaskensis (Lockington, 1877)

LIT

Crangon alaskensis - Holthuis 1980:150-151 (with original citations); Butler 1980:108-109, fig.; Kuris & Carlton 1977:547; Jensen 1995:40, fig. 59; *Neocrangon alaskensis* - **Zarenkov 1965.**

Crangon alaskensis and *C. nigricauda* may be hybrids (Jensen 1995).

Distribution - Bering Sea to San Diego, 0 – 555 m.

Origins: Native.

Zarenkov, N.A. 1965. Revision of the genus *Crangon* Fabricius and *Sclerocrangon* G.O. Sars (Decapoda, Crustacea) Zool. Zhur., 44(12): 1761-1775 (InRussian).

HIPPOLYTIDAE

Heptacarpus brevirostris (Dana, 1852)

LIT

Holthuis 1980:126 (with synonymy); Butler 1980:231-232, fig.; Jensen 1995:46, fig. 75.

Abundant in rocky intertidal full marine areas in salinities 9-31 PSU.

Distribution – Attu, Aleutian Islands, Alaska to Bahia Magdalena, Baja California, intertidal to 128 m.

Origins: Native

PALAEEMONIDAE

Exopalaemon modestus Heller, 1862

LCRANS, LIT

Kingdom: Animalia
Phylum: Arthropoda
Subphylum: Crustacea

Leander modestus - Heller, 1862; *Leander czerniavskyi* - Brashnikov, 1907; *Leander czerniavskyi lacustris* - Brashnikov, 1907; *Palaemon leander* - modestus Gee 1925; *Leander modestus sibirica* - Brashnikov, 1907; *Exopalaemon modestus* – Holthius 1980:83.

Exopalaemon modestus is distinguished from *E. carinicauda* (introduced into San Francisco Bay Wicksten 1997) by its smaller chelae of the second pereopod and by two distal spines which extend beyond the tip of the median telson process, in contrast to the small distal spines of *E. carinicauda* which are short of the median telson process. Distribution – In Asia, siberian prawn ranges from northern Korea to southern China in freshwater lakes and rivers. *Exopalaemon modestus* is presently known in the eastern Pacific only from the Columbia River and the Willamette River (Emmett *et al.* 2002:447-450).

[Ed. Note: CDFG reports *E. modestus* from the Sacramento River see <http://www.dfg.ca.gov/cabw/camlnetste.pdf> but the author of this section is doubtful of correct identification]

Origins: Most probably introduced into the Columbia River with ballast water traffic from Asia sometime before 1995.

GRAPSIDAE

Eriocheir japonica de Haan, 1835

Japanese mitten crab

LIT

Abundant in San Francisco Bay, California and in northern Europe. One male specimen was caught on a line by a sturgeon fisherman in the Columbia River near Astoria in the summer of 1998. No other crabs of the genus *Eriocheir* have been captured since.

Origins: Introduced but not established in LCR.*

Hemigrapsus oregonensis (Dana, 1851)

LCRANS, LIT

Schmitt 1921:274-276, fig. 162 (with synonymy); Hart 1982:220-221, fig. 91; Jensen 1995:17, fig. 18.

Distribution – Resurrection Bay, Alaska to Baja California, Mexico, almost exclusively intertidal, tolerates reduced salinities and fresh water for brief periods.

Origin: Native.

Phylum: Chordata

Subphylum: Vertebrata

Superclass: Osteichthyes

* - resulted in an unsuccessful introduction

Acipenseridae

Kingdom: Animalia
 Phylum: Chordata
 Superclass: Osteichthyes

<i>Acipenser medirostris</i>	LIT	Native
<i>Acipenser transmontanus</i>	LIT	Native
<i>Acipenser</i> or <i>Scaphirhynchus</i>	LIT	Introduced*
Agonidae		
<i>Ocella verrucosa</i>	LIT	Native
<i>Pallasina barbata</i>	LIT	Native
<i>Stellerina xyosterna</i>	LIT	Native
Ammodytidae		
<i>Ammodytes hexapterus</i>	LIT	Native
Anguillidae		
<i>Anguilla</i> sp.	LIT	Introduced*
Catostomidae		
<i>Catostomus macrocheilus</i>	LIT	Native
<i>Catostomus platyrhynchus</i>	LIT	Native
Centrarchidae		
<i>Ambloplites rupestris</i>	LIT	Introduced
<i>Lepomis cyanellus</i>	LIT	Introduced
<i>Lepomis gibbosus</i>	LIT	Introduced
<i>Lepomis gulosus</i>	LIT	Introduced
<i>Lepomis macrochirus</i>	LIT	Introduced
<i>Lepomis microlophus</i>	LIT	Introduced
<i>Micropterus dolomieu</i>	LIT	Introduced
<i>Micropterus salmoides</i>	LIT	Introduced
<i>Pomoxis annularis</i>	LCRANS, LIT	Introduced
<i>Pomoxis nigromaculatus</i>	LIT	Introduced
Characidae		
<i>Piaractus brachypomus</i>	LIT	Introduced*
<i>Pygocentrus nattereri</i>	LIT	Introduced*
Clupeidae		
<i>Alosa sapidissima</i>	LIT	Introduced
Cobitidae		
<i>Misgurnus anguillicaudatus</i>	LIT	Introduced
Cottidae		
<i>Artedius fenestralis</i>	LIT	Native
<i>Cottus aleuticus</i>	LIT	Native
<i>Cottus asper</i>	LIT	Native
<i>Enophrys bison</i>	LIT	Native
<i>Hemilepidotus hemilepidotus</i>	LIT	Native
<i>Hemilepidotus spinosus</i>	LIT	Native
<i>Leptocottus armatus</i>	LIT	Native
<i>Scorpaenichthys marmoratus</i>	LIT	Native
Cyprinidae		
<i>Acrocheilus alutaceus</i>	LIT	Native
<i>Carassius auratus</i>	LIT	Introduced
<i>Ctenopharyngodon idella</i>	LIT	Introduced
<i>Cyprinus carpio</i>	LIT	Introduced

Kingdom: Animalia
 Phylum: Chordata
 Superclass: Osteichthyes

<i>Mylocheilus caurinus</i>	LIT	Native
<i>Oregonichthys crameri</i>	LIT	Native
<i>Ptychocheilus oregonensis</i>	LIT	Native
<i>Rhinichthys cataractae</i>	LIT	Native
<i>Rhinichthys falcatus</i>	LIT	Native
<i>Richardsonius balteatus</i>	LIT	Native
<i>Tinca tinca</i>	LIT	Introduced
Embiotocidae		
<i>Amphistichus rhodoterus</i>	LIT	Native
<i>Cymatogaster aggregata</i>	LIT	Native
<i>Embiotoca lateralis</i>	LIT	Native
<i>Hyperprosopon anale</i>	LIT	Native
<i>Hyperprosopon argenteum</i>	LIT	Native
<i>Hyperprosopon ellipticum</i>	LIT	Native
<i>Phanerodon furcatus</i>	LIT	Native
<i>Rhacochilus vacca</i>	LIT	Native
Engraulidae		
<i>Engraulis mordax</i>	LIT	Native
Esocidae		
<i>Esox lucius x masquinongy</i>	LIT	Introduced
Fundulidae		
<i>Fundulus diaphanus</i>	LIT	Introduced
Gadidae		
<i>Lota lota</i>	LIT	Native
<i>Microgadus proximus</i>	LIT	Native
Gasterosteidae		
<i>Gasterosteus aculeatus</i>	LIT	Cryptogenic
Gobiidae		
<i>Lepidogobius lepidus</i>	LIT	Native
Hexagrammidae		
<i>Hexagrammos decagrammus</i>	LIT	Native
<i>Ophiodon elongatus</i>	LIT	Native
Ictaluridae		
<i>Ameiurus catus</i>	LIT	Introduced
<i>Ameiurus melas</i>	LIT	Introduced
<i>Ameiurus natalis</i>	LIT	Introduced
<i>Ameiurus nebulosus</i>	LIT	Introduced
<i>Ictalurus furcatus</i>	LIT	Introduced
<i>Ictalurus punctatus</i>	LIT	Introduced
Merlucciidae		
<i>Merluccius productus</i>	LIT	Native
Moronidae		
<i>Morone chrysops</i>	LIT	Introduced
<i>Morone chrysops x saxatilis</i>	LIT	Introduced
<i>Morone saxatilis</i>	LIT	Introduced
Osmeridae		

Kingdom: Animalia
 Phylum: Chordata
 Superclass: Osteichthyes

<i>Allosmerus elongatus</i>	LIT	Native
<i>Hypomesus pretiosus</i>	LIT	Native
<i>Spirinchus starksi</i>	LIT	Native
<i>Spirinchus thaleichthys</i>	LIT	Native
<i>Thaleichthys pacificus</i>	LIT	Native
Paralichthyidae		
<i>Citharichthys sordidus</i>	LIT	Native
<i>Citharichthys stigmaeus</i>	LIT	Native
Percidae		
<i>Perca flavescens</i>	LCRANS, LIT	Introduced
<i>Stizostedion vitreum</i>	LIT	Introduced
Percopsidae		
<i>Percopsis transmontana</i>	LIT	Native
Petromyzontidae		
<i>Lampetra ayresii</i>	LIT	Native
<i>Lampetra richardsoni</i>	LIT	Native
<i>Lampetra tridentata</i>	LIT	Native
Pholidae		
<i>Pholis ornata</i>	LCRANS, LIT	Native
Pleuronectidae		
<i>Platichthys stellatus</i>	LIT	Native
<i>Pleuronichthys coenosus</i>	LIT	Native
<i>Psettichthys melanostictus</i>	LIT	Native
Pleuronectidae		
<i>Isopsetta isolepis</i>	LIT	Native
<i>Parophrys vetulus</i>	LIT	Native
Poeciliidae		
<i>Gambusia affinis</i>	LIT	Introduced
Rajidae		
<i>Raja binoculata</i>	LIT	Native
Salmonidae		
<i>Oncorhynchus clarki</i>	LIT	Native
<i>Oncorhynchus clarki x mykiss</i>	LIT	Introduced
<i>Oncorhynchus keta</i>	LIT	Native
<i>Oncorhynchus kisutch</i>	LIT	Native
<i>Oncorhynchus mykiss</i>	LIT	Native
<i>Oncorhynchus mykiss gairdneri</i>	LIT	Native
<i>Oncorhynchus nerka</i>	LIT	Native
<i>Oncorhynchus tshawytscha</i>	LIT	Native
<i>Prosopium williamsoni</i>	LIT	Native
<i>Salmo trutta</i>	LIT	Introduced
<i>Salvelinus confluentus</i>	LIT	Native
<i>Salvelinus malma</i>	LIT	Native
Scorpaenidae		
<i>Sebastes melanops</i>	LIT	Native
<i>Sebastes miniatus</i>	LIT	Native

Kingdom: Animalia
 Phylum: Chordata
 Superclass: Osteichthyes

Serrasalminae			
	<i>Piaractus brachipomus</i>	LIT	Introduced
Squalidae			
	<i>Squalus acanthias</i>	LIT	Native
Stichaeidae			
	<i>Lumpenus sagitta</i>	LIT	Native
Syngnathidae			
	<i>Syngnathus leptorhynchus</i>	LIT	Native
Trichodontidae			
	<i>Trichodon trichodon</i>	LIT	Native

ACIPENSERIDAE

Acipenser medirostris Ayres, 1854 green sturgeon
 LIT
 Origin: Native

Origin: Native distribution: Pacific Coast of North America from Alaska to Baja California. In estuaries, the lower reaches of large rivers, and in salt or brackish water off of river mouths (Froese and Pauly 2003).

Acipenser transmontanus Richardson, 1836 white sturgeon
 LIT
 Origin: Native

Native distribution: Pacific Coast of North America from Alaska to Monterey, California. Considered landlocked in parts of the Columbia River drainage. Spends most of its time in the sea, usually close to shore then enters estuaries of large rivers and moves inland to spawn (Froese and Pauly 2003).

***Acipenser* or *Scaphirhynchus* sp. - unk.** Eastern sturgeon
 LIT

At the conclusion of the 1905 Lewis and Clark Exhibition held in Portland, Oregon two specimen of Eastern sturgeon from the Atlantic coast of North America (exact species unknown) were released into Guilds Lake on the Willamette River (Lampman 1946). However, no sightings or catches of Eastern sturgeon have been reported in the lower Columbia River since then. It is likely that these fish perished naturally or were caught prior to the opening of Guild's Lake to the Willamette River in 1909.

AGONIDAE

Ocella verrucosa (Lockington, 1880) warty poacher
 Synonyms: *Brachyopsis verrucosus*
 LIT
 Origin: Native

Kingdom: Animalia
Phylum: Chordata
Superclass: Osteichthyes

Marine species, distributed throughout the Eastern Pacific from Bristol Bay, Alaska to California (Froese and Pauly 2003). Probably an infrequent visitor to the lower Columbia River estuary.

Pallasina barbata (Steindachner, 1876) tubenose poacher
Synonyms: *Siphogonus barbatus*
LIT
Origin: Native

Intertidal species often found in eelgrass or seagrass beds. Native distribution: North Pacific from the Sea of Japan to the Bering Sea and to Central California (although this may represent two subspecies) (Froese and Pauly 2003).

Stellerina xyosterna (Jordan & Gilbert, 1880) pricklebrest poacher
Synonyms: *Brachyopsis xyosternus*
LIT
Origin: Native

Demersal, marine species. Native distribution: Eastern Pacific from British Columbia to Baja California (Froese and Pauly 2003).

AMMODYTIDAE

Ammodytes hexapterus Pallas, 1814 Pacific sand lance
LIT
Origin: Native

Origin: Native distribution: Arctic and Pacific south to Southern California, and the Western Atlantic (although this may be a separate species) (Froese and Pauly 2003). Found in brackish and marine waters in schools or buried in the sand.

ANGUILLIDAE

***Anguilla* sp.** eel
LIT
Origin: Introduced

Unsuccessful introduction. Reports of *Anguilla* on the west coast of North America were reviewed by Williamson and Tabeta (1991) following the capture of several eels presumed to have escaped or released after importation as live seafood. J.L. Galbreath captured three unidentified eels of the genus *Anguilla* in the Willamette River at Portland in 1981, 1982 and 1983. Williamson and Tabeta (1991) concluded that all of the eels captured on the West Coast were the result of intentional or unintentional introductions

Kingdom: Animalia
Phylum: Chordata
Superclass: Osteichthyes

and do not represent natural spread via oceanic currents. In addition, *Anguilla* eels are catadromous, spawning in areas of the open ocean where temperatures and salinities are consistently high. Such areas are not available to eels on the west coast of North America and introduced *Anguilla* would be unable to successfully reproduce upon reaching maturity. With no further reports of eels we assume that this limited introduction has been naturally extirpated from the lower Columbia River basin.

CATOSTOMIDAE

Catostomus macrocheilus Girard, 1856 largescale sucker

LIT

Origin: Native

Native to Western North America (Froese and Pauly 2003).

Catostomus platyrhynchus (Cope, 1874) mountain sucker

Synonyms: *Minomus platyrhynchus*, *Pantosteus jordani*, *Pantosteus columbianus*

LIT

Origin: Native

Native to Western North America (Froese and Pauly 2003).

CENTRARCHIDAE

Note: Identification of specific dates and mechanisms of introductions of Centrarchidae and other spiny-rayed fishes into the lower Columbia River Basin is complicated by several poorly documented intentional fish releases. In 1893 the United States Fish Commission (USFC, predecessor of the U.S. Fish and Wildlife Service) released 50 largemouth bass, *Micropterus salmoides*, along with “various sunfish” into the Willamette River just north of Salem from a shipment of fishes captured in the Illinois River (Lampman 1946). Anecdotal information compiled by Lampman (1946) suggests that the “various sunfish” included *Pomoxis annularis*, *Pomoxis nigromaculatus*, *Lepomis gibbosus*, *Ambloplites rupestris*, and other juvenile sunfish as well as several types of catfish and channel cats. Twelve years later, at the 1905 Lewis and Clark Centennial Exhibition in Portland, the USFC displayed a tank of spiny-rayed fishes. At the conclusion of the Exhibition these fishes were reportedly released into the waters of Guild’s Lake on the Willamette River (Lampman 1946). An accurate inventory of the exhibit is unavailable but one exhibitor recalled that the collection of freshwater fishes included large- and smallmouth bass (*Micropterus salmoides*, *M. dolomieu*.), crappies (*Pomoxis annularis*, *P. nigromaculatus*), bluegill (*Lepomis macrochirus*) and two eastern sturgeon (*Acipenser* or *Scaphirhynchus* sp.) (Lampman 1946). The waters of the lake were dammed for the exhibition but the dam leading to the Willamette River was removed in 1909 after which the lake was filled in and turned into an industrial site. Between 1905 and 1909 the lake was a popular fishing hole, especially for local youth

Kingdom: Animalia
Phylum: Chordata
Superclass: Osteichthyes

(Lampman 1946). Many of these repeatedly introduced species became established and *Pomoxis annularis*, *P. nigromaculatus*, *Micropterus dolomieu*, *M. salmoides*, *Lepomis macrochirus*, *L. gibbosus*, and *L. gulosus* continue to be captured in fish surveys of the Willamette and parts of the lower Columbia (Hutchinson and Aney 1964, Farr and Ward 1993, North et al. 2002).

Ambloplites rupestris (Rafinesque, 1817) rock bass

Synonyms: *Bodianus rupestris*

LIT

Origin: Introduced

Unsuccessful introduction. Native to the Great Lakes region of North America, *Ambloplites rupestris* was first introduced unsuccessfully into the Willamette River near Salem, Oregon along with large mouth bass imported from Ohio around 1888 by Gideon Steiner, a local fish and poultry businessman (Lampman 1946). In 1893 the USFC may have also unsuccessfully introduced the rock bass into the Willamette River (see overview of the Centrarchidae above). Since that time sporadic records of *A. rupestris* in the Willamette have been attributed to misidentifications of the successfully introduced warmouth *Lepomis gulosus* (Lampman 1946). *A. rupestris* has been introduced in several lakes and rivers in Washington (Wydoski and Whitney 1979) but it is unknown if any of these introductions have been spread into the lower Columbia River basin. However, intentional stocking of *A. rupestris* for sportfishing was widespread in the late 1800s through the 1940s with successfully established populations common in the Mid-West and the Mid-Atlantic states (NAS 2003).

Lepomis cyanellus Rafinesque, 1819 green sunfish

LIT

Origin: Introduced

Native to many river basins in central and eastern North America, *Lepomis cyanellus* may have been intentionally introduced in 1893 when the USFC released “various sunfish” captured in Illinois into the Willamette River just north of Salem (Lampman 1946). *L. cyanellus* may also have been introduced into Blue Lake (Hutchinson and Aney 1964), a small lake along the bank of the Columbia River near Troutdale, Oregon that continues to be popular with sport fishermen, but the date of that introduction is not known. *L. cyanellus* has been widely introduced throughout the west and, in California, has been held partially responsible for the decline of many native amphibians and fishes (NAS 2003, Moyle 1976). In the 1960s, an attempt by the Washington Department of Fish and Wildlife to extirpate *L. cyanellus* from Satcheen Lake, Washington failed (Wydoski and Whitney 1979). The status of *L. cyanellus* as an invasive species in the lower Columbia River Basin is elusive. Lampman (1946) reported that *L. cyanellus* might occur in the river basin and noted that while previous surveys of the Willamette River failed to capture this species a suspicious hybrid *Lepomis* (green sunfish hybridizes readily with other *Lepomis* (Moyle 1976)) had been caught. In addition, *L. cyanellus* has long been documented in Washington and Oregon at locations outside of the lower Columbia River drainage basin (Chapman 1942, Wydoski and Whitney 1979, Bond 1994) and Altman et

Kingdom: Animalia
Phylum: Chordata
Superclass: Osteichthyes

al. (1997) report green sunfish present in two major rivers, the Pudding and the Tualatin, both which converge with the Willamette River

Lepomis gibbosus (Linnaeus, 1758) pumpkinseed
Synonyms: *Perca gibbosa*, *Eupomotis gibbosus*, *Pomotis vulgaris*
LIT
Origin: Introduced

Lepomis gibbosus, a sport fish native to many river basins in central and eastern North America, may also have been introduced in 1893 when the USFC released “various sunfish” captured in Illinois into the Willamette River (Lampman 1946). Although no records of this introduction exist with the USFC an editorial in *The Oregonian* makes special mention of this event reporting that *L. gibbosus* had been captured during a salvage operation on the overflowing waters of the Illinois River and released later in Oregon waters (Lampman 1946). A popular sport fish *L. gibbosus* has been widely and successfully introduced in the waters of Oregon and Washington and has been found in the Willamette and the lower Columbia River (Chapman 1942, Lampman 1946, Hutchinson and Aney 1964, Wydoski and Whitney 1979, Farr and Ward 1993, Bond 1994, Altman et al. 1997, and North et al. 2002).

Lepomis gulosus (Cuvier, 1829) warmouth
Synonyms: *Pomotis gulosus*, *Chaenobryttus gulosus*
LIT
Origin: Introduced

Easily misidentified as various other species of sunfish *Lepomis gulosus*, native to many river basins in central and eastern North America, may have been released into the Willamette in 1893 by the USFC but records of this species in the lower Columbia River basin were sporadic until the later half of the 20th Century. Chapman and DeLancy (1933) published the first report of warmouth from Washington State having capturing several fish during a survey in 1930 in a slough of the Kalama River near Kalama, Washington. It is worth noting the conspicuous absence of any mention of this species in Lampman (1946) as *The Coming of the Pond Fishes* is one of the most thorough reports of intentional and unintentional fish introductions into the Willamette and lower Columbia rivers prior to the 1950s. Current populations of warmouth may be derived from numerous plantings. Discussions of *L. gulosus* in Bond (1994) and Altman et al. (1997) indicate that the popular sport fish is widely established in the freshwaters of the lower Columbia Basin. Most recently Farr and Ward (1993) and North et al. (2002) confirmed the presence of *L. gulosus*, capturing it in fish surveys along the lower Willamette.

Lepomis macrochirus Rafinesque, 1819 bluegill, bluegill sunfish
LIT
Introduced

Kingdom: Animalia
Phylum: Chordata
Superclass: Osteichthyes

Similar in history to *Lepomis gulosus*, *L. macrochirus*, native to many river basins in central and eastern North America (Page and Burr 1991), is widely distributed throughout the Willamette and parts of the lower Columbia River Basin. *L. macrochirus* was likely to have first been intentionally introduced into the system during the 1893 USFC release of fish near Salem. Chapman and DeLancy (1933) captured two specimens of bluegill, along with *L. gulosus*, in 1930 near Kalama, Washington. Pond stocking and plantings by individuals may also have contributed to the successful establishment of the bluegill. *L. macrochirus* are regularly reported in fish surveys of the lower Willamette (North et al. 2002, Altman et al. 1997, and Farr and Ward 1993). According to Froese and Pauly (2003) several countries (South Africa, Kenya, Venezuela, Panama, Japan and Mexico) have reported adverse ecological effects after establishment of this widely introduced species sportfish.

Lepomis microlophus (Günther, 1859) redear sunfish

Synonyms: *Pomotis microlophus*

LIT

Origin: Introduced

Unlike many of the above sunfish, *L. microlophus* is native to the south-eastern United States (Page and Burr 1991). According to Mills et al. (1993), redear sunfish were first introduced into the Great Lakes Basin in 1928 and then spread into inland areas of the basin, making it less likely that the redear was introduced into the Willamette in the assortment of Illinois sunfish released by the USFC in 1893. Bond (1994) lists *L. microlophus* as found in parts of western Oregon including ponds in the Willamette Valley but includes no additional location information. Altman et al (1997) reports that *L. microlophus* has been recorded in the lower Willamette however none were captured in surveys by Farr and Ward (1993) or North et al. (2002). It is possible that reports of this species may be misidentifications of other introduced *Lepomis* or hybrids, and that introduced redear have not escaped the ponds mentioned in Bond (1994). If redear sunfish are present in the Columbia River Basin they might be considered a threat to endemic mollusks of concern as *L. microlophus* is a more voracious molluscivore than other sunfishes (NAS 2002). Ecological effects are unknown from introductions in other countries (Froese and Pauly 2003).

Micropterus dolomieu Lacepède, 1802 smallmouth bass, smallie, black bass, brown bass, white trout, green trout

Synonyms: *Centrarchus fasciatus*

smallmouth bass, smallie, black bass, brown bass, white trout, green trout

Established in the lower Columbia River basin. Native to the Midwestern United States (Scott and Crossman 1973), *Micropterus dolomieu* has been intentionally introduced throughout the world to enhance sport fishing (Froese and Pauly 2003). In 1874, Livingston Stone, inventor of the “aquarium car” used by the USFC to transport fish stock by rail across the U.S., transported 99 *M. dolomieu* from the east coast to California releasing the surviving 85 fish into tributaries of San Francisco Bay (Smith 1896). The first reported introductions of *M. dolomieu* to the lower Columbia River Basin took place

Kingdom: Animalia
Phylum: Chordata
Superclass: Osteichthyes

nearly 50 years later in the 1920s. In 1923, fish from Wisconsin were introduced by a local game warden without the approval of the USFC into Lake Oswego, Oregon (Lampman 1946). The survival of these fish is uncertain however in 1924 the same game warden imported bass from a lake in the Puget Sound region and released them into the Willamette River (Lampman 1946). In 1925 *M. dolomieu* were planted in the middle stretch of the Columbia River Basin as well. *M. dolomieu* continue to be caught by recreational fishermen and in regional fish surveys (Farr and Ward 1993). In other states introduced *M. dolomieu* have been implicated in the decline or elimination of native fishes (Minckley 1973, Jenkins and Burkhead 1994). Smallmouth bass, which have been shown to prey on smolts of Pacific salmonids under laboratory conditions, may pose a threat to declining populations of wild salmon in the lower Columbia River Basin (Dentler 1993).

Micropterus salmoides (Lacepède, 1802) largemouth bass, black bass, green trout
Synonyms: *Labrus salmoides*, *Huro salmoides*, *Aplites salmoides*, *Perca nigricans*, *Huro nigricans*, *Grystes megastoma*

LIT

Origin: Introduced

With a native range stretching from the Great Lakes to the Gulf Coast of North America (Page and Burr 1991), *M. salmoides* is a popular sport fish and has been introduced widely throughout the world (Froese and Pauly 2003). The first largemouth bass were introduced to the Willamette River in 1888 in two separate plantings. Gideon Steiner (a fish and poultry businessman), feeling that the area lacked the “splendid eastern game fish of his childhood”, imported and released a shipment of *M. salmoides* and *Ambloplites rupestris* from Toledo, Ohio into Willamette River near Salem, Oregon (Lampman 1946). The same year a prominent Portland lawyer, Edward Bingham, released 25 bass into the Willamette River, presumably near his home in Lake Oswego, Oregon (Lampman 1946). Four years later, the USFC released 500 *M. salmoides* in the Willamette River with subsequent smaller releases throughout the lower Willamette River basin in 1895 (Smith 1896). Between 1890 and 1895 the USFC also planted 5442 largemouth bass throughout the state of Washington and 1597 largemouth bass in the Boise River (a population that was subsequently boosted by a private release of 2240 bass the same year in the middle stretch of the Columbia River basin) (Smith 1896, Lampman 1946). In early August 1898, The Oregonian reported the capture of the first largemouth bass in the Columbia River just downstream of where Bonneville Dam now stands. It is not known which of the aforementioned releases led to the establishment of largemouth bass throughout the lower Columbia River basin.

Along with other introduced predatory centrarchids *M. dolomieu* may also be responsible for declines in native amphibian populations (NAS 2003). Adult fish feed on other fishes, crayfish and frogs while immature *M. dolomieu* feed on crustaceans, insects and small fishes (Page and Burr 1991). Adverse ecological effects have also been reported from France, Italy, Japan, South Africa, Cuba, Guatemala, and Mexico (Froese and Pauly 2003).

Kingdom: Animalia
Phylum: Chordata
Superclass: Osteichthyes

Pomoxis annularis Rafinesque, 1818 white crappie, calico bass

Pomoxis nigromaculatus (Lesueur, 1829) black crappie, calico bass

Synonyms: *Cantharus nigromaculatus*, *Pomoxis sparoides*

LIT

Origin: Introduced

It is largely impossible to discuss these two established species separately as they are often lumped together and referred to solely as “crappie,” as *Pomoxis* spp., and/or misidentified as a single species (Dill and Cordone 1997). *Pomoxis* spp. are native to North America spanning the Great Lakes, Hudson Bay and Mississippi River basins, Ontario, Canada west to Minnesota and South Dakota, and south to the Gulf of Mexico (Page and Burr 1991). Two prominent releases of crappie into the lower Columbia River system were made by the USFC in 1893 and in 1905. In 1893 the USFC released 50 largemouth bass, *Micropterus salmoides*, along with “various sunfish,” including crappie, into the Willamette River (Lampman 1946). At the 1905 Lewis and Clark Centennial Exhibition in Portland, the USFC displayed a tank of spiny-rayed fishes that were later released into the waters of Guild’s Lake on the Willamette River (Lampman 1946). According to a member of the Oregon Game Commission the crappie were so abundant in Guild’s Lake during their four year impoundment that small boys were catching large quantities of them using fish books baited with paraffin chewing gum (Lampman 1946). These fish are considered “harmless” by the IUCN and no ecological impacts have been reported from introduced locations around the world (Froese and Pauly 2003)

CHARACIDAE

Piaractus brachypomus (Cuvier, 1818) pirapatinga, pacu, red bellied pacu

Synonyms: *Myletes brachypomus*, *Colossoma brachypomum*

LIT

Origin: Introduced

One of several reoccurring but unsuccessful exotic aquarium species in the lower Columbia River Basin, *Piaractus brachypomus* is a tropical South American fish popular in the aquarium trade (Froese and Pauly 2003). The pacu is a member of the Serrasalminae family, a family that includes piranha, and these fish are often imported and sold under the misnomer “vegetarian piranhas.” Pacu have a rapid growth rate and voracious appetite, and may readily outgrow the hobby tank they were originally housed in. These characteristics may lead frustrated fish owners to dispose of the fish in nearby waters. Pacu have developed a reputation as the species most often found in non-native waters that creates a piranha scare in the local media especially as juveniles are readily mis-identified as the traumatogenic red-bellied piranha (NAS 2003). The first *P. brachypomus* found in Oregon waters was a specimen caught in July 1988 by a fisherman in the Willamette River near the Port of Portland’s Terminal Four (Logan et al. 1996). Since that time two additional *P. brachypomus* have been collected from the Willamette River in 1992 and 1995 respectively and two additional unverified sightings were

Kingdom: Animalia
Phylum: Chordata
Superclass: Osteichthyes

reported from Hood River and Salem, Oregon in 1990 and 1991 (Logan et al. 1996). It is highly unlikely that all of these fish were the result of one single release but rather represent five separate releases. Logan et al. (1996) tested the thermal tolerance of *Piaractus* and determined that it is not low enough to tolerate to survive normal water temperatures found in the lower Columbia River basin from November through April making it impossible for these fish to overwinter in these waters. As these fish have a primarily herbivorous and insectivorous diet no adverse effects on local fish populations have been reported (Froese and Pauly 2003).

Pygocentrus nattereri Kner, 1858 red piranha, red-belly piranha
Serrasalmus nattereri, *Pygocentrus altus*, *P. ternetzi*.

LIT

Origin: Introduced

Unsuccessful introduction. *Pygocentrus nattereri*, a common aquarium species, is notorious for its reputation as a traumatogenic species (Froese and Pauly 2003). Concern about the establishment of this species in Florida or Texas in Gulf of Mexico drainages (where water temperatures are high enough for them to overwinter) is high. While the waters of Oregon and Washington are too cold in winter for *P. nattereri* to become established, this species could pose a threat to salmon smolts and other small fishes (Quinn 2003). In August 2003, a single *P. nattereri* was caught by a teenager in Johnson Creek, Oregon (a stream that drains to the lower Willamette) (Quinn 2003).

CLUPEIDAE

Alosa sapidissima (Wilson 1811) American shad, common shad, white shad
Synonyms: *Clupea sapidissima*

LIT

Origin: Introduced

Native to the Atlantic coast of North America from Labrador to Florida (Scott and Crossman 1973), shad were planted in the Sacramento River, California in 1871 having been transported across country from the Atlantic coast (Smith 1896). Although the Columbia River was intentionally stocked several times in the late 1880s (Smith 1896, Linder 1963, Wydoski and Whitney 1979) anecdotal evidence suggests that shad began showing up in the Columbia River as early as 1876 (Smith 1896) with the first published capture of a shad made by the ichthyologist David Starr Jordan in 1880 (Jordan 1916). Due to repeated introductions into the Sacramento and Columbia Rivers, *Alosa sapidissima* is now established on the west coast from Alaska to Baja California (Froese and Pauly 2003). Large runs of shad are common in the Columbia River and the impoundment of the Columbia may have improved conditions for spawning and rearing (Sherwood and Creager 1990, Weitkamp 1994, Petersen unpublished). Returns of introduced *A. sapidissima* to the Columbia River increased significantly between 1960 and 1990, and as a result, shad have become so abundant in the fish ladders that they may interfere with the passage of other fishes (NAS 2003). Although there is a commercial

Kingdom: Animalia
Phylum: Chordata
Superclass: Osteichthyes

fishery for shad it is considered under-harvested (~ 9% of the population that passes through Bonneville Dam is caught annually by limited commercial and recreational fishing) because the timing coincides with an endangered summer Chinook salmon (Petersen et al. unpublished).

The ecological effects of *A. sapidissima* in the lower Columbia River are poorly understood. It has been speculated that juveniles could alter the zooplankton community, enhance the diet of resident predators, and/or compete with native salmon for habitat or food resources however data to support or dismiss these theories are limited (Petersen et al unpublished).

COBITIDAE

Misgurnus anguillicaudatus (Cantor, 1842) Oriental weatherfish, pond loach

Synonyms: *Cobitis anguillicaudata*

LIT

Origin: Introduced

Native to Eastern Asia, *Misgurnus anguillicaudatus* has been introduced into Hawaii, the U.S. mainland, the Philippines, Australia, Palau, Turkmenistan, and Mexico (Froese and Pauly 2003). The earliest records of *M. anguillicaudatus* in the continental United States date back to the 1930s when escapes from aquarium fish culture facilities were reported (Courtenay and Hensley 1980). While it is a popular aquarium fish and many introductions may be attributable to aquarium dumping this species has been introduced for the purposes of aquaculture in several countries where it is reared as a food fish, for bait, and for the aquarium industry (Froese and Pauly 2003, NAS 2003). *M. anguillicaudatus* has several life-history traits that may contribute to successful establishment – wide tolerance of physiological parameters, low vulnerability to predation, a flexible diet, and a high reproductive potential (Logan et al. 1996). In Oregon, an established population of *M. anguillicaudatus* has been reported from a diked secondary channel of the Clackamas River where it was discovered in the mid 1980s (Logan et al 1996). Several fish were also collected in 1997 from Multnomah Channel near the Columbia River (NAS 2003). Reports of *M. anguillicaudatus* are likely to under-represent their populations as these fish are typically found in shallow, muddy waters with dense vegetation, i.e. even when abundant they are difficult to capture with standard fish survey gear (NAS 2003). The dispersal ability of the Clackamas population of *M. anguillicaudatus* (and its relation to the population in Multnomah Channel) is unknown (Logan et al. 1996). It may be likely that these two occurrences are the result of separate aquarium releases. Adverse ecological impacts have been reported in Hawaii and Australia (Froese and Pauly 2003, NAS 2003) where these species are suspected of being carriers of fish pathogens and predators on native aquatic species.

COTTIDAE

Artedius fenestralis Jordan & Gilbert, 1883 padded sculpin

LIT

Kingdom: Animalia
Phylum: Chordata
Superclass: Osteichthyes

Origin: Native

Intertidal marine species native to the eastern Pacific: from Alaska to Southern California (Froese and Pauly 2003).

Cottus aleuticus Gilbert, 1896 coastrange sculpin

LIT

Origin: Native

Cottus aleuticus is a catadromous sculpin, native to the Pacific Coast of North America from Alaska to Northern California (Froese and Pauly 2003). It inhabits gravel and rubble riffles of medium to large rivers and rocky shores of lakes and occasionally enters estuaries (Froese and Pauly 2003).

Cottus asper Richardson, 1836 prickly sculpin

LIT

Origin: Native

Native to Pacific coast drainages of North America. The coastal form of this species is catadromous. Sometimes used as a bait species. May have been introduced east of the Rockies in Canada (Froese and Pauly 2003).

Enophrys bison (Girard, 1854) buffalo sculpin

Synonyms: *Aspicottus bison*

LIT

Origin: Native

Native to the eastern Pacific from Alaska to central California, *Enophrys bison* is a marine species commonly found in inshore rocky and sandy areas (Froese and Pauly 2003).

Hemilepidotus hemilepidotus (Tilesius, 1811) Red Irish lord

Synonyms: *Cottus hemilepidotus*

LIT

Origin: Native

A commercially and recreationally harvested marine sculpin, *Hemilepidotus hemilepidotus* is native to the North Pacific from Kamchatka, Russia to central California (Froese and Pauly 2003).

Hemilepidotus spinosus Ayres, 1854 Brown Irish lord

LIT

Origin: Native

A marine species, *Hemilepidotus spinosus* is native to the eastern Pacific from southeastern Alaska to southern California (Froese and Pauly 2003).

Kingdom: Animalia
Phylum: Chordata
Superclass: Osteichthyes

Leptocottus armatus Girard, 1854 Pacific staghorn sculpin
LIT
Origin: Native

A brackish to marine species, *Leptocottus armatus* is native to the west coast of North America from Alaska to Baja California (Froese and Pauly 2003).

Scorpaenichthys marmoratus Girard, 1854 cabezon
LIT
Origin: Native

A marine species, *Scorpaenichthys marmoratus* is native to the west coast of North America from Alaska to Baja California (Froese and Pauly 2003).

CYPRINIDAE

Acrocheilus alutaceus Agassiz & Pickering, 1855 chiselmouth
LIT
Origin: Native

Native to the Pacific Northwest, *Acrocheilus alutaceus* is a freshwater fish that inhabits flowing pools, creeks and small to medium rivers (Froese and Pauly 2003).

Carassius auratus (Linnaeus, 1758) goldfish
Synonyms: *Cyprinus auratus*, *Cyprinus langsdorfi*, *Cyprinus thoracatus*, *Carassius chinensis*, *Cyprinus maillardi*
LIT
Origin: Introduced

Origin: Introduced throughout the world, goldfish are native to central Asia. *Carassius auratus* is cultured widely by the aquarium and ornamental pond trades. It is occasionally reared for use as bait and less frequently as a food item (Froese and Pauly 2003). Goldfish have been widely and repeatedly stocked in the United States from many points of origin, including both Asia and Europe. Having been bred for a range of body forms and colors there are many varieties of goldfish in U.S. waters. It is common for goldfish to hybridize with common carp *Cyprinus carpio* (another introduced species)(NAS 2003). During the late 1800s the USFC breed goldfish and distributed them to states as fish suitable for aquaria, fountains, and ornamental ponds (McDonald 1887, 1893 in NAS 2003). Introductions in the Pacific Northwest may represent escapes from private ponds (Smith 1896) as well as from aquarium releases by individuals (Courtenay and Hensley 1979). The earliest report of goldfish in the lower Columbia River basin comes from Lampman (1946) who notes seeing goldfish feeding in the Willamette River in 1933. Chapman (1942) reports capturing goldfish in surveys at the mouth of the Columbia River and near Kalama, Washington. In the 1960s, Hutchinson

Kingdom: Animalia
Phylum: Chordata
Superclass: Osteichthyes

and Aney (1964) report goldfish scattered throughout the lower Willamette Basin. Wydoski and Whitney (1979) note that the distribution of goldfish in the northwest is “subject to constant change because people thoughtlessly discard goldfish into various waters,” and also observed that a small number of goldfish were being raised locally for bait.

Ctenopharyngodon idella (Valenciennes, 1844) grass carp, white amur
Synonyms: *Leuciscus idella*
LCRANS, LIT
Origin: Introduced

Grass carp (*Ctenopharyngodon idella*), also known as the white amur, is an herbivorous fish native to parts of eastern Asia from the Amur River of eastern Russia to southern China (NAS 2003). Grass carp have been widely introduced throughout the world although not all populations have become established (Froese and Pauly 2003). Rationalization for intentional stocking includes commercial aquaculture and exploration of aquaculture potential, research, establishment of a food resource, and biological control (Froese and Pauly 2003). First introduced from Malaysia into the U.S. by the USFWS Fish Farming Experimental Station in 1962, established populations of *C. idella* exist in parts of the Mississippi and Missouri Rivers, as well as in Alabama and Florida (Courtenay et al. 1984). Grass carp are reported to occur in 45 states (although establishment of populations is uncertain because of their primarily triploid status) where they can cause significant changes in macrophyte, phytoplankton and invertebrate communities, etc. The loss of aquatic vegetation caused by grass carp has been implicated in the decline of waterfowl habitat (NAS 2003). Stocking of triploid (functionally sterile) grass carp, both authorized and unauthorized, is a widely implemented biological control method used to reduce unwanted aquatic vegetation. According to NAS (2003) “the species has spread rapidly as a result of widely scattered research projects, stockings by federal, state, and local government agencies, legal and illegal interstate transport and release by individuals and private groups, escapes from farm ponds and aquaculture facilities; and natural dispersal from introduction sites (e.g., Pflieger 1975; Lee et al. 1980 et seq.; Dill and Cordone 1997).” However, the effectiveness of grass carp as biological control has been criticized on several levels; grass carp often consume non-target native plants as well as or in preference to unwanted weeds (Taylor et al. 1984), the reproductive potential of triploids has been questioned (as has the success of suppliers in creating truly triploid fish), and the potential for negative interactions between grass carp and both invertebrates and fishes has been raised as a unwanted cost (Courtenay et al. 1984).

Grass carp will seek out and follow flowing water, so that all inlets and outlets of the pond or lake where they have been introduced for biological control must be screened. During flood events grass carp may escape even screened ponds. Loch and Bonar (1999) observed 49 adult grass carp migrating up the Columbia River in 1996 and 1997, emphasizing the need for the carp to be truly sterile. Although they may not be established (i.e. reproducing) in the lower Columbia River a repeated pattern of escape into the river, combined with the potential for non-triploid introductions, and the

Kingdom: Animalia
Phylum: Chordata
Superclass: Osteichthyes

longevity of the species (10-40 years) have created an “artificially established” population in the lower river. Large grass carp are regularly caught in Youngs Bay and other parts of the lower Columbia River, and have been recorded passing through the fish ladders at Bonneville Dam (Jim Athern personal communication).

Cyprinus carpio Linnaeus, 1758 common carp
LIT
Origin: Introduced

Native to Eurasia (Page and Burr 1991) *Cyprinus carpio*, better known as the common carp, has been introduced into every state in the United States except Alaska (and it is believed to be established in all but Maine) (NAS 2003). Records disagree as to when and where the first carp were introduced. DeKay (1842 in NAS 2003) reported that the species was first brought into New York from France by a private citizen and released into the Hudson River a few years later but debate over the species identification exists for this and other early reports (NAS 2003). Smith (1896) reported that common carp first appeared in the United States in 1872 when several fish imported from Germany, planted in private ponds in Sonoma, California, propagated for commercial rearing, and distributed to individuals on the west coast for rearing as food fish (Lampman 1946). In 1880, one Captain Harlow of Portland, Oregon imported 35 mature German carp from San Francisco to breed and sell carp for stocking private ponds. In spring 1881, the Sandy River flooded and washed an estimated 3000 immature carp from Captian Harlow’s breeding pond into the Columbia River (Lampman 1946). This may not have been an isolated event as reports show that in 1877 the USFC imported carp from Germany and began shipping domestically breed carp to private applicants in Oregon and Washington as early as 1882 (Smith 1896, Lampman 1946). Within ten years of Captain Harlow’s carp escape *C. carpio* had established itself throughout the lower Columbia River basin and was no longer popular with the local fishermen. The Oregonian newspaper reported that locals were offering carp for sale for use as fertilizer at a price of \$5/ton (Lampman 1946). In the Columbia River *C. carpio* continue to be abundant in the sloughs and inlets of the lower river (often hybridizing with *Carassius auratus*) and populations supported a small commercial fishery in Lake Vancouver, Washington through the late 1930s (Chapman 1942). Today, *C. carpio* is regarded as a potential pest species because of its widespread introduction and establishment, and because its feeding behavior (rooting in soft sediment) often leads to the loss of vegetation and increased sediment suspension (Laird and Page 1996). Of primary concern is the destruction of submerged and emergent aquatic vegetation that provide habitat for native fish and food for waterfowl (Dentler 1993). There is also evidence that *C. carpio* will prey on fish eggs (Moyle 1976). In the Pacific Northwest, Miller and Beckman (1996) documented white sturgeon *Acipenser transmontanus* eggs in the stomachs of common carp in the Columbia River.

Mylocheilus caurinus (Richardson, 1836) peamouth
Synonyms: *Clarkina caurina*, *Cyprinus caurinus*
LIT
Origin: Native

Kingdom: Animalia
Phylum: Chordata
Superclass: Osteichthyes

Native to the Pacific Slope of North America (Froese and Pauly 2003).

Oregonichthys crameri (Snyder, 1908) Oregon chub

Synonyms: *Hybopsis crameri*

LIT

Origin: Native, endangered

Endemic to the Willamette and Umpqua River drainages in Oregon. It is rare in Willamette because of habitat alteration (Froese and Pauly 2003). Loss of habitat combined with the introduction of non-native fish species to the Willamette Valley such as largemouth bass, smallmouth bass, crappie, bluegill, and mosquitofish has resulted in a sharp decline in Oregon chub abundance. The chub was given "endangered" status under the federal Endangered Species Act in 1993.

Ptychocheilus oregonensis (Richardson, 1836) Northern pikeminnow

Synonyms: *Cyprinus oregonensis*

LIT

Origin: Native

Although native to Pacific drainages of North America (Froese and Pauly 2003), *Ptychocheilus oregonensis*, is considered a pest species because large concentrations of squawfish near hydroelectric projects are responsible for substantial salmonid predation and may further increase salmonid mortality by reducing the fish guidance efficiency of submersible traveling screens (NOAA 1994). *P. oregonensis*, a lake-adapted fish, has responded favorably the creation of reservoirs and other slow moving water habitat creation along the Columbia River. In free-flowing areas the bottom- and bank-hugging pikeminnow is not as problematic a predator for salmonid smolts.

Rhinichthys cataractae (Valenciennes, 1842) longnosed dace

Synonyms: *Gobio cataractae*, *Rhinichthys marmoratus*

Lit

Origin: Native, sensitive

The longnose dace is present on both sides of the Continental Divide and is one of the most widely distributed of the western fishes (Froese and Pauly 2003). *Rhinichthys cataractae* ssp. is listed by the ODFW as a sensitive species or "Species of Concern" in Oregon waters.

Rhinichthys falcatus (Eigenmann & Eigenmann, 1893) leopard dace

Synonyms: *Agosia falcata*

Lit

Origin: Native

Kingdom: Animalia
Phylum: Chordata
Superclass: Osteichthyes

The leopard dace is native to the Fraser and Columbia River drainages (Froese and Pauly 2003). It is primarily found in slow streams and gravel runs in the upper Columbia R. drainage.

Richardsonius balteatus (Richardson, 1836) redbside shiner

LIT

Origin: Native

Native to Pacific Slope drainage British Columbia to southern Oregon, including the Columbia Basin (Froese and Pauly 2003), the redbside shiner has been introduced (probably as bait) to drainages in Arizona, Colorado, Montana and Utah (NAS 2003).

Tinca tinca (Linnaeus 1758) tench, green tench, golden tench

Synonyms: *Cyprinus tinca*

LIT

Origin: Introduced

Native to Eurasia as well as the British Isles (Berg 1949), *Tinca tinca* was introduced to numerous locations in Africa, Australia, Japan, and North America with no known adverse ecological impacts (Froese and Pauly 2003). Prized by recreational fishermen for their tasty flesh, tench are omnivorous, feeding on benthic invertebrates, aquatic insect larvae, and algae some other invertebrates. In Great Britain tench are popular ornamental pond species but a search of ornamental pond websites in the United States did not reveal a similar opinion. Wydoski and Whitney (1979) write, in reference to the Washington tench population, “to our knowledge it has not created any particular problems.” In the late 1800s, spurred by the success in Australia with tench breeding and introduction programs, the USFC imported tench from Europe (Baughman 1947). Raised in fish ponds in Washington State, 450 *T. tinca* were introduced into several lakes and ponds in the lower Columbia River basin between 1895-1896 (Smith 1896, Baughman 1947). The current status of *T. tinca* in the lower Columbia River basin remains uncertain. Tench appear to have spread (or were transplanted) from their original introduction sites and into the lower Columbia within 40 years of the USFC planting. Chapman (1942), in a paper on introduced fishes in the Pacific Northwest, noted that tench, while found in the Columbia River, were nowhere near as abundant as were *Cyprinus carpio* (habitat requirements of tench are similar to that of *C. carpio*, and the two species are superficially similar, with tench being the smaller of the two). Hutchinson and Aney (1964) list *T. tinca* on their list of known fish species in the Willamette basin, and note their distribution as “Columbia mainstem, probably lower Willamette. Wydoski and Whitney (1979) report *T. tinca* as present in the Columbia River system, Spokane River, and Lake Washington. Bond (1961) noted tench as introduced to the Columbia River and the Willamette River but in later revisions (Bond 1973, 1994) stated that the species was in the Columbia River and was once present in lower Willamette River. No further captures of tench have been reported in the lower

Kingdom: Animalia
Phylum: Chordata
Superclass: Osteichthyes

Columbia but tench are occasionally captured in the middle Columbia River (USFWS <http://hanfordreach.fws.gov/fish.html>)

EMBIOTOCIDAE

Amphistichus rhodoterus (Agassiz, 1854) redbtail surfperch

Synonyms: *Holconotus rhodoterus*, *Cymatogaster pulchellus*, *Cymatogaster larkinsis*, *Amphistichus heermanni*

LIT

Origin: Native

Brackish, marine species native to the Eastern Pacific (Froese and Pauly 2003), popular with commercial and recreational anglers.

Cymatogaster aggregata Gibbons, 1854 shiner perch

LIT

Origin: Native

Brackish, marine species native to the Northeastern Pacific (Froese and Pauly 2003).

Embiotoca lateralis Agassiz, 1854 striped seaperch, blue seaperch

LIT

Origin: Native

Marine species found in coastal areas, native to the Eastern Pacific (Froese and Pauly 2003). Minor commercial importance, often targeted by aquarium enthusiasts.

Hyperprosopon anale Agassiz, 1861 spotfin surfperch

LIT

Origin: Native

Marine species native to the Eastern Pacific (Froese and Pauly 2003) often found in surf on sandy beaches.

Hyperprosopon argenteum Gibbons, 1854 walleye surfperch

LIT

Origin: Native

Marine gamefish native to the Eastern Pacific (Froese and Pauly 2003).

Hyperprosopon ellipticum (Gibbons, 1854) silver surfperch

Synonyms: *Cymatogaster ellipticus*

LIT

Origin: Native

Kingdom: Animalia
Phylum: Chordata
Superclass: Osteichthyes

Small marine fish native to the Eastern Pacific (Froese and Pauly 2003).

Phanerodon furcatus

LIT

Origin: Native

Marine fish native to the Eastern Pacific, usually found offshore (Froese and Pauly 2003).

Rhacochilus vacca (Girard, 1855) pile perch

Synonyms: *Damalichthys vacca*, *Ditrema vacca*, *Damalichthys argyrosomus*

LIT

Origin: Native

Marine fish native to the Eastern Pacific, usually found along the rocky shore (Froese and Pauly 2003).

ENGRAULIDAE

Engraulis mordax Girard, 1854 Northern anchovy, California anchovy

LIT

Origin: Native

Pelagic, marine species native to the Northeast Pacific (Froese and Pauly 2003).
Commercially harvested along the West Coast.

ESOCIDAE

Esox lucius x masquinongy tiger muskellunge, tiger musky

LIT

Origin: Introduced

Hybrid freshwater species *Esox lucius* x *Esox masquinongy*. Tiger muskellunge have been bred artificially and stocked by state fish and game agencies for sport fishing throughout North America. Populations are often maintained by stocking as male tiger muskellunge are always sterile, but females are often fertile (Becker 1983). This hybrid predator is probably deleterious to smaller fish. Tiger musky in the lower Columbia River basin were reported by the Warmwater Fisheries Resource Manager, Washington Department of Wildlife, Olympia, WA in 1992 (NAS 2003).

FUNDULIDAE

Fundulus diaphanous (Lesueur, 1817) banded killifish

Synonyms: *Hydrargira diaphana*, *Fundulus multifaciatius*

LIT

Kingdom: Animalia
Phylum: Chordata
Superclass: Osteichthyes

Origin: Introduced

Note: Froese and Pauly (2003) treats *F. diaphanous* and *F. diaphanous diaphanous* as valid synonyms while NAS (2003) lists *F. d. diaphanous* and *F. d. menona* as eastern and western subspecies

Fundulus diaphanous is native to Atlantic slope drainages in North America (Froese and Pauly 2003) and has been introduced to parts of Ohio, Pennsylvania, South Dakota, Oregon and Washington (NAS 2003). Banded killifish are grown commercially for aquariums and for use as bait (Froese and Pauly 2003). The ecological implications of introduced populations are not known. Banded killifish were first recorded from the upper Columbia River estuary at Jones Beach in 1971 (Misitano and Sims 1974) but were not consistently captured in fish surveys until the late 1980s (Hinton et al 1990). The source of the Columbia River introduction is unknown but thought to be the result of a bait dump (NAS 2003). Other introductions across the United States have been attributed to accidental introduction along with stocked largemouth bass (South Dakota) or as the result of an aquarium release (Ohio) (NAS 2003). The continued presence of *F. diaphanous* in the Willamette River as well as the lower Columbia River is well documented (see Misitano and Sims 1974, Hjort et al 1984, Hinton et al 1990, Hinton et al. 1992b, Farr and Ward 1993, Weitkamp 1994, Hinton and Emmett 2000, and North et al. 2002).

GADIDAE

Lota lota (Linnaeus, 1758) burbot

Synonyms: *Gadus lota*, *Gadus lacustris*, *Gadus maculosus*

LIT

Origin: Native

Lota lota is the only freshwater member of the Gadidae family. Congregate in deep pools of large rivers and lakes. Circumarctic distribution (Froese and Pauly 2003).

Microgadus proximus (Girard, 1854) Pacific tomcod

Synonyms: *Gadus proximus*, *Morrhua californica*, *Gadus californicus*

LIT

Origin: Native

Brackish and marine species native to the Eastern Pacific (Froese and Pauly 2003).

Minor commercial and recreational species.

GASTEROSTEIDAE

Gasterosteus aculeatus stickleback

LIT

Origin: Cryptogenic

Kingdom: Animalia
Phylum: Chordata
Superclass: Osteichthyes

Wide distribution may indicate that this species should be considered cryptogenic although previous studies consider this to be a native species. Large salinity tolerance can range from freshwater to marine salinities. Distributed along the West Coast of North America from Baja California to the Bering Sea (Fishbase 2004), as well as throughout the coastal regions in the North Pacific and North Atlantic.

GOBIIDAE

Lepidogobius lepidus

bay goby

LIT

Origin: Native

Intertidal marine demersal species, likely only found near the mouth of the Columbia.

HEXAGRAMMIDAE

Hexagrammos decagrammus

kelp greenling

LIT

Origin: Native

Marine species, likely not a resident of the lower Columbia River estuary.

Ophiodon elongatus

lingcod

LIT

Origin: Native

Marine species, likely not a resident of the lower Columbia River estuary.

ICTALURIDAE

Catfish, popular as both food and sport fish, were among the first fishes introduced to the West Coast (Smith 1896). In 1874, Livingston Stone and his USFC aquarium car are responsible for the first western movement of catfish and bullhead across the Rocky Mountains, their natural westernmost boundary (Smith 1896). Present on this train were three species, *Ameiurus catus*, *Ameiurus nebulosus* and *Ictalurus punctatus* (Smith 1896). It is unknown if the first catfish stocked in the Pacific Northwest were descended from this original population or the result of later importations. By the 1880s catfish (of many unreported species) had become successfully established in Silver Lake, Washington (stocked by an unknown person). Fearing that the catfish would be “another enemy to our salmon” a former Fish Commissioner of Oregon asked the Washington Commissioners of Fish for permission to rid Silver Lake (which connects to the Columbia via Cowlitz River) of its catfish population (Smith 1896). It has been theorized that fishermen’s fears were heightened by speculation that the introduced catfish population included specimen of *Ictalurus furcatus*, blue catfish, native to the

Kingdom: Animalia
Phylum: Chordata
Superclass: Osteichthyes

Mississippi, growing to over 100 lbs (the maximum recorded weight of a blue catfish is 186 lbs), and theoretically capable of consuming 20 lb salmon (Lampman 1946). However, due to importation and release of a variety of catfish species by private parties into the lower Willamette and Columbia Rivers in the 1880s, extirpation of the Silver Lake population would not have kept catfish out of the Columbia River (Lampman 1946). By 1890, The Oregonian newspaper carried an article on the newly arrived catfish stating, “The ponds and lakes of Sauvie Island are literally alive with catfish which have been carried in by the late flood waters. By every appearance our waters will soon be swarming with these fish, as they increase at an appalling rate” (Lampman 1946). By 1894 catfish were thoroughly established throughout the lower Columbia and Willamette Rivers and by the 1890s a commercial harvest of catfish had begun (Lampman 1946) however, by 1938 only 2.5 percent of the recorded catch of game fish in Washington was catfish (Chapman 1942). Due to the voracious and predatory nature of catfish most are considered ecological pests. Several species of introduced North American freshwater catfish have been implicated in the decline of native fish (Marsh and Douglas 1997, Froese and Pauly 2003) and amphibians (Rosen et al. 1995) both in the United States and elsewhere. Declared a game species by the State of Oregon in 1913, catfish are no longer commercially harvested in the Pacific Northwest. The three most common catfish species in the lower Columbia River are *A. nebulosus*, *A. natalis* and *I. punctatus*.

Ameiurus catus (Linnaeus, 1758)

ESTABLISHED

Ictalurus catus, *Silurus catus*
white catfish, white bullhead

Native to the Atlantic and Gulf slope drainages of the United States, *Ameiurus catus* were first released in California in the San Joaquin River in 1874 (Smith 1896). It is likely that they were part of the population planted in Silver Lake, Washington in the early 1880s and became distributed throughout the lower Columbia River basin by 1894 (Lampman 1946). In spite of this planting and at least one additional intentional introduction in 1930 by an Oregon hatchery superintendent (Lampman 1946) *A. catus* has never been a plentiful species in the lower Columbia River basin (Wydoski and Whitney 1979). In Bond’s (1994) revision of his key to Oregon fishes he is uncertain of their establishment, however a report by the Oregon Department of Fish and Wildlife lists a 15lb white cat was caught in Tualatin River in 1989 by Wayne Welch and setting a new state record (<http://www.dfw.state.or.us/ODFWhtml/InfoCntrFish/FishRecords.html>) indicating that their may indeed still be an established population of *A. catus* in the lower Columbia River basin.

Ameiurus melas (Rafinesque, 1820)

ESTABLISHED

Ictalurus melas, *Ictalurus melas*, *melas*, *Aneiurus melas melas*, *Silurus melas*
black bullhead, black catfish

Native to North America east of the Rockies excluding the Atlantic slope, *Ameiurus melas* has been intentionally stocked throughout the west coast and other parts Europe for sport and as a food fish (Froese and Pauly 2003, NAS 2003). Countries such as Germany, Spain and Hungary report adverse ecological impact after introduction (Froese

Kingdom: Animalia
Phylum: Chordata
Superclass: Osteichthyes

and Pauly 2003). Voracious predators, black bullhead, along with other catfish, have been implicated in the decline of native fish (Marsh and Douglas 1997) and amphibians (Rosen et al. 1995) in the United States. In addition, the black bullhead is considered a nuisance fish by anglers targeting other gamefish (Froese and Pauly 2003). The first *A. melas* in the Columbia River basin was caught in the Willamette in 1894 (Smith 1896, Lampman 1946) but it is not known when or where this species was first introduced. A 1945 Oregon State College surveyed the fishes of the Willamette River system and reported the presence of three species of bullhead catfish: *A. nebulosis*, *A. natalis* and *A. melas* (Lampman 1946). Although it is not one of the commonly captured catfish, Bond (1994) continues to list it as present in the Columbia River drainage.

Ameiurus natalis (Lesueur, 1819) ESTABLISHED
Pimelodus natalis, *Ictalurus natalis*
yellow bullhead

Native to the North America from the Mississippi basin east (Page and Burr 1991), *Ameiurus natalis* has been widely and successfully stocked throughout the western United States (NAS 2003). It has been introduced into Italy and Mexico where adverse ecological impacts have been reported from the later country where it has replaced several endemic species (Froese and Pauly 2003). Although *A. natalis* is a popular sport and food fish, predation by it and other catfish may have an impact on its introduced habitat. Lampman (1946) asserts that the first introduction of yellow bullheads in the region was probably in 1905, when tanks of warm water display fish were released following the Lewis and Clark Centennial exposition in Portland see centrarchidae discussion above for more information. *A. natalis* have been common in the Willamette Valley since then. They were captured by the 1945 Oregon State College survey of the Willamette River (Lampman 1946), as well during the Farr and Ward 1993 survey of the lower Willamette.

Ameiurus nebulosus (Lesueur, 1819) ESTABLISHED
Ictalurus nebulosus, *Pimelodus nebulosus*
brown bullhead, brown catfish

Native to Atlantic and Gulf slope drainages and parts of the Mississippi River drainage basin (NAS 2003), *Ameiurus nebulosus* is the most common catfish in the lower Columbia River basin and is especially abundant in the sloughs and slack waters of the basin (Chapman 1942, Wydoski and Whitney 1979). During the 1890s and up until catfish were declared game species by the state of Oregon (thus not open to commercial harvest) in 1913, there was a thriving commercial fishery for *A. nebulosus*, mostly in the shallow lakes of Sauvie Island. At its peak, this fishery annually produced over 100,000 pounds of catfish (Lampman 1946). Collections of *A. nebulosus* span most of the lower Columbia River basin (see Smith 1896, Chapman 1942, Lampman 1946, Bond 1973, 1994, Wydoski and Whitney 1979, Hjort 1984, Farr and Ward 1993, and USFWS 1993)

Ictalurus furcatus (Valenciennes, 1840) UNKNOWN
Pimelodus furcatus, *Ictalurus meridionalis*

Kingdom: Animalia
Phylum: Chordata
Superclass: Osteichthyes

blue catfish

Native to the Mississippi River basin, the status of *Ictalurus furcatus*, is unknown however it seems unlikely that this species of catfish is established in the lower Columbia River. Unlike many of the above catfish, *I. furcatus* prefers deeper, clear, flowing water habitats, and it is not considered a pest species by Froese and Pauly (2003). Bond (1994) describes *I. furcatus* as “introduced, Columbia River, not common” but it is unclear which portion of the Columbia River he is referring to. *I. furcatus* are present in both the Snake and the middle reach of the Columbia River but are rarely reported below the Bonneville Dam. It seems that the dreaded salmon-eating blue catfish of Silver Lake never materialized in the lower Columbia River see above discussion of Ictaluridae. It is interesting to note however that blue catfish have been intentionally stocked in parts of California for biological control of *Corbicula fluminea*, the non-native Asian clam under the hope that, even if clam populations were not controlled, the biomass of the clams would at least be significant enough to create trophy-sized catfish (Dill and Cordone 1997).

Ictalurus punctatus (Rafinesque, 1818)

ESTABLISHED

Silurus punctatus

channel catfish, graceful catfish

Ictalurus punctatus, native to the central drainages of North America from Southern Canada to Northern Mexico, is a commercially important species, is heavily aquacultured species and an Albino form is commonly encountered in the aquarium trade (Froese and Pauly 2003). In 1893, 100 channel cats were released into the Boise River in Idaho (Smith 1896). Reports of *I. punctatus* caught in the lower Columbia river were sporadic up until the 1940s, but it is suspected that channel cats were stocked in the Willamette River in the 1920s by an Oregon hatchery superintendent (Lampman 1946). Additional releases were made in ponds, lake, and rivers throughout Washington and Oregon as many species of catfish became established. Now they exist primarily in mid-Columbia and Snake River although they are established in the Willamette River as well (Hjort et al 1984, Farr and Ward 1993). Sterile populations of channel cats have been stocked in Washington lakes by the Washington Department of Fish and Wildlife, introduced to increase predation on over-abundant forage fish populations, and to add diversity to gamefish populations (WDFW 2003).

MORONIDAE

Morone chrysops (Rafinesque, 1820)

ESTABLISHED

Perca chrysops, *Roccus chrysops*

white bass

Native to the Mississippi River drainage basin, *Morone chrysops*, have been stocked legally and illegally throughout much of the United States (NAS 2003). First transported

Kingdom: Animalia
Phylum: Chordata
Superclass: Osteichthyes

west in 1895 by the USFC with a shipment of black bass from Illinois, *M. chrysops* were introduced into California waters for breeding purposes (Smith 1896). Lee et al. (1980) reports a population of *M. chrysops* with a limited range in the lower Lewis River drainage basin in Washington.

Morone chrysops x saxatilis

UNSUCCESSFUL

wiper, sunshine bass, whiterock, palmetto, Cherokee

An artificial hybrid, *Morone chrysops x saxatilis* has no native range. This cultivated sport fish has been introduced to numerous watersheds in central and eastern United States. Where one or both parent species exists with *M. chrysops x saxatilis* backcrosses are often present and are considered detrimental to the native parent population (NAS). specimen turned into ODF during Farr and Ward sampling? (Morone hybrid?). Populations of *M. chrysops x saxatilis* are artificially maintained in all locations where they have been intentionally stocked for sport. An experimental stocking program using *M. chrysops x saxatilis* exists in southwestern Oregon in Ten Mile Lakes (Farr and Ward 1993, Bond 1994). The three *M. chrysops x saxatilis* hybrids caught by anglers in the Willamette River and turned over to the fish surveys of Farr and Ward (1993) may have been migrants from this program or were illegally introduced specimen.

Morone saxatilis (Walbaum, 1792)

ESTABLISHED

Perca saxatilis, *Roccus saxatilis*

striped bass, striper, rock, rockfish

Morone saxatilis is a highly prized sportfish native to Atlantic slope drainages and the northeastern Gulf slope of the United States (Page and Burr 1991). Striped bass have been widely stocked for sportfishing in coastal waters from New York to California (landlocked stocked populations exist also) (NAS 2003). In addition, between 1886 and 1992, *M. saxatilis* has been introduced to and become established in Mexico, South Africa, Iran, Russia, Ecuador, and British Columbia (Froese and Pauly 2003). According to Chapman (1942) there are no records of striped bass introduced into the coastal waters of the Pacific Northwest however in 1879, 135 fingerlings from New Jersey were introduced into San Francisco Bay by Livingston Stone (Smith 1896). Supplemented in 1882 with 300 additional fish from New Jersey (Smith 1896), *M. saxatilis* spread up and down the West Coast and now range from British Columbia to Baja California (NAS 2003, Froese and Pauly 2003). As this is a highly valued sportfish it is interesting to note that detrimental ecological and nuisance effects of ANS on *M. saxatilis*, itself an ANS, have been described and along the West Coast (e.g. *Potamocorbula amurensis* reducing striped bass food availability, and bait theft by *Eriocheir sinensis* of anglers targeting this species). The distribution of *M. saxatilis* in the Columbia River is well documented (see Moyle 1976; Wydoski and Whitney 1979; Grabowski et al. 1984; Bond 1994) and striped bass is closed to commercial fishing throughout the state. Impacts of striped bass are unknown – however Morgan and Gerlach (1950) reported finding numerous trout and salmon fry as well as fingerlings in gut content surveys in Coos Bay, Oregon.

PERCIDAE

Kingdom: Animalia
Phylum: Chordata
Superclass: Osteichthyes

Perca flavescens (Mitchill, 1814)

ESTABLISHED

Morone flavescens

yellow perch, American perch, lake perch

Native to much of the Atlantic, Great Lakes, and Mississippi River basins (Wydoski and Whitney (1979), *Perca flavescens* is introduced or native in all but five U.S. states. The introduction of yellow perch to West Coast habitats by Livingstone Stone and his Aquarium Car was justified by Stone who wrote, “*Perca flavescens* is at all events far preferable to most of the fish at present existing in the freshwaters of California, and even if it destroyed four-fifths of the other fish there it would replace them by a better kind” (Smith 1896). Established in the Willamette and lower Columbia River (Farr and Ward 1993, NAS 2003), yellow perch are often considered a nuisance in lakes and rivers where they compete with adult trout for food resources and prey upon younger trout (Coots 1966). The Columbia River population may have been the result of several intentional introductions. In 1894, the USFC planted *Perca flavescens* in Silver Lake (on the Cowlitz River) and over the next ten years almost 1000 perch were planted in Washington lakes (Wydoski and Whitney 1976). In 1905, yellow perch were believed to be present in the aquarium exhibit released into Guild Lake (Lampman 1946). In the 1930s, *Perca flavescens* was an important species in regional lake fisheries in the 1930s but its sustainable population levels were rapidly exceeded and most fish were reported to appear “stunted” (Lampman 1946).

Sander vitreus (Mitchill, 1818)

Stizostedion vitreum, *Perca vitrea*

walleye, walleye pike

LIT

Introduced

Although *Sander vitreus* were first transplanted to the West Coast in 1874 by Livingston Stone (Smith 1896), walleye were not captured in the lower Columbia River basin until 1980 (Li et al 1979, Farr and Ward 1993, NAS 2003). It is believed that, since their introduction into the upper Columbia near Lake Roosevelt in the 1940s or 1950s, walleye have gradually spread downriver and may have established a limited population in the lower Columbia (Dentler 1993, Farr and Ward 1993, NAS 2003). Native to the Great Lakes through the Mississippi basin (Froese and Pauly 2003), *S. vitreus* is a popular gamefish that lives in aquatic habitats from ponds to large rivers. A recent literature review by McMahon and Bennett (1996) found that the effects of walleye introductions in the Pacific Northwest were complex but posed a threat to salmonids through smolt predation. Because of this *S. vitreus* is banned from introduction into Oregon waters (McMahon and Bennett 1996).

POECLIIDAE

Gambusia affinis (Baird & Girard, 1853)

Kingdom: Animalia
Phylum: Chordata
Superclass: Osteichthyes

Syn: *Heterandria affinis*, *Gambusia affinis affinis*

Mosquitofish

Introduced

LIT

Previously reported as the sub-species *Gambusia affinis affinis* but now recognized as a full species (Moyle and Davis 2000), the western mosquitofish is native to the Atlantic and Gulf Slope drainages from New Jersey to Mexico and the Mississippi River basin (Froese and Pauly 2003, NAS 2003). *Gambusia affinis* is one of the most successful introduced fish species in the world having gained a near global distribution (Welcomme 1988). Adverse ecological impacts have been reported from Europe, Asia, India, Australia, Africa and South America as well as from numerous island countries where *G. affinis* has been extensively introduced for mosquito control (see Froese and Pauly 2003). While *G. affinis* was introduced to much of the United States and to countries around the world for mosquito control starting in the 1960s, recent critical reviews of this practice suggest that this fish may not be any more successful than native minnows at consuming mosquito larvae and reducing mosquito-borne disease (Courtenay and Meffe 1989). However, adverse ecological effects have resulted from mosquitofish predation on the eggs, larvae, and juveniles of various native fishes. Although still distributed to private parties for mosquito control, in the Western United States *G. affinis* has been implicated in the extirpation and/or decline of populations of federally endangered and threatened species of minnow and chub (Courtenay and Meffe 1989). In Oregon, the sharp decline in the population of *Oregonichthys crameri*, the Oregon chub - an endangered species, has been attributed to habitat loss and predation by introduced fishes including *G. affinis* (Scheerer 1999)

SALMONIDAE

Oncorhynchus clarki x mykiss
cuttbow trout

Artificially? ESTABLISHED

The status of cuttbow trout as a nonindigenous species is non-straightforward. *Oncorhynchus clarki x mykiss*, the result of a cross between *O. clarki* x *O. mykiss*, is considered an artificial hybrid occurring in the wild where parent species come in contact with one another through stocking, and not present (or rare) where both parents occur naturally together in their native range (Sigler and Miller 1963). While both parent species are native to the lower Columbia River basin both species have been widely stocked throughout the Columbia River, the result of fish enhancement and hatchery programs (Froese and Pauly 2003). Further complicating matters, the hybrid cuttbow has also been intentionally stocked in the western U.S. as sport fish (NAS 2003). The ease of hybridization between the two parent species may be contributing to a reduction in genetic integrity of these species and the replacement of threatened cutthroat trout populations by hybridization and competition (NAS 2003).

Kingdom: Animalia
Phylum: Chordata
Superclass: Osteichthyes

Salmo trutta Linnaeus, 1758
brown trout, German brown trout

UNKNOWN

Salmo trutta is native to Europe and western Asia (Page and Burr 1991). First introduced to the inland waters of North America in 1883 by the USFC, *S. trutta* is now present throughout the U.S. (Courtenay et al. 1984, NAS 2003). Natural reproduction rates in North America are poor thus many states actively stock this popular gamefish to maintain desirable population sizes (NAS 2003). Chapman (1942) reports that while *S. trutta* was widely planted in Oregon and Washington it was successful in only a few locations.

Phylum: Chordata
Subphylum: Vertebrata
Class: **Amphibia**

Ranidae		
<i>Rana catesbeiana</i>	LCRANS, LIT	Introduced

RANIDAE

Rana catesbeiana Shaw, 1802 bullfrog, American bullfrog
LCRANS, LIT
Introduced

Native to eastern and central North America, *Rana catesbeiana*, the bullfrog, is widely introduced in the western states including Hawaii. Speculation as to the intent of early introductions includes plantings intended for food (to provide frog legs for the West Coast frog leg market which declined in the 1930s) (ODFW 2001) as well as for aesthetic purposes (i.e. for their distinctive croaking sound) (Lampman 1946). In 1914, the Oregon Fish and Game Commission granted permission to a private individual to introduce this frog into the mid-Columbia River basin below John Day (Lampman 1946). In 1924 or 1925, reports Lampman (1946), bullfrogs resulting from the above planting were shipped to Portland for further distribution in the lower Columbia River basin. Mature bullfrogs are responsible for significant levels of predation on native aquatic species, including the spotted frog (*Rana pretiosa*), the Western pond turtle (*Clemmys marmorata*) and the Oregon chub (*Oregonichthys crameri*) (ODFW 2001, Crayon 2002).

Kingdom: Animalia
Phylum: Chordata
Class: Reptilia

Phylum: Chordata
Subphylum: Vertebrata
Class: Reptilia
Order: Testudines

Reptiles were not collected or identified over the course of this study. Nonnative turtles have been introduced numerous times over the years, likely both intentionally to enhance wildlife and through aquarium/terrarium disposal. The introduced species pose a threat to native species whose populations are in decline.

Chelydridae			
	<i>Chelydra serpentina serpentina</i>	LIT	Introduced
Emydidae			
	<i>Chrysemys picta bellii</i> (Wpaint t)	LIT	Native
	<i>Clemmys marmorata</i> (WPT)	LIT	Native
	<i>Trachemys scripta elegans</i>	LIT	Introduced

CHELYDRIDAE

Chelydra serpentina serpentina Gray, 1831 Eastern snapping turtle

LIT

Introduced

Native to eastern North America several established populations of this snapping turtle have been reported from the Willamette Valley including Portland, OR (see <http://nas.er.usgs.gov/queries/SpFactSheet.asp?speciesID=1226> for more information).

EMYDIDAE

Chrysemys picta bellii (Gray, 1831) Western painted turtle

LIT

Native

This turtle is found primarily in northern Willamette Valley and ranges east through the Columbia River Gorge and Columbia Basin.

Clemmys marmorata (Baird and Girard, 1852) Western pond turtle

LIT

Native

Clemmys marmorata is considered to be rare throughout its range. It is almost extirpated in Washington State, and the current western pond turtle population in Oregon is thought to be less than 10% of its historical population.

<https://www.nwp.usace.army.mil/op/V/western.htm>

Trachemys scripta elegans (Weid-Neuwied, 1838) red-eared slider

LIT

Kingdom: Animalia
Phylum: Chordata
Class: Reptilia

Introduced

Native to the Southeastern United States and popular as an aquarium species since the 1930s, *T. scripta elegans* has been introduced throughout the western United States primarily through aquarium releases and escapes. NAS attributes part of the the turtle's recent popularity and subsequent releases/escapes tto the Teenage Mutant Ninja Turtle television cartoon craze of the late 1980s - see

<http://nas.er.usgs.gov/queries/SpFactSheet.asp?speciesID=1261>

Phylum: Chordata
Subphylum: Vertebrata
Class: Mammalia
Order: Rodentia

Echimyidae		
<i>Myocastor coypus</i>	LCRANS, LIT	Introduced

ECHIMYIDAE

Myocastor coypus Kerr, 1792 Nutria, coypu, coypu rat, swamp beaver, nutria rat
LCRANS, LIT
Introduced

Native to South America, *Myocastor coypus* - an aquatic rodent, is a textbook example of how far astray well-intentioned importation and release of nonnative species can go. Introduced numerous times into the United States, beginning as early as 1899 in California (USGS 2000), most releases (and escapes) of nutria were intended to enhance the fur trade. For example, in 1938, twenty nutria were imported from Argentina to Louisiana by Tabasco sauce tycoon E.A. McIlhenny, these nutria reportedly escaped captivity during a hurricane in the early 1940s and subsequently spread along the Gulf Coast (NAS 2003). Other introductions of nutria in North America were made for biological control of unwanted aquatic weeds such as water hyacinth (*Eichhornia crassipes*) and alligator weed (*Alternanthera philoxeroides*) (USGS 2000) a program that failed to significantly reduce the target plants. Nutria are considered an economic liability in many areas as their burrowing activity can damage earthen dams and dikes and because they often feed on the young shoots of crop plants (ODFW 2001). The burrowing activity of nutria may also contribute to streambed erosion in the lower Columbia River basin. Severe ecological impacts have been reported in the southern Atlantic states where nutria has caused extensive loss of marshland (NAS 2003). Nutria may also compete with native muskrats populations for food and habitat. ODFW (2001) reports that anecdotal evidence suggests that in locations where nutria are abundant, muskrat populations decline. Nutria were introduced into the wilds of the Pacific Northwest in 1937 when an unknown number escaped from a fur farm in

Kingdom: Animalia
Phylum: Chordata
Class: Reptilia

Tillamook Co. aided by a large flood. Today, nutria can be found throughout the lower Columbia River basin and much of western Oregon and Washington (ODFW 2001).