

2-19-2015

# Bioenergy Harvest, Climate Change, and Forest Carbon in the Oregon Coast Range

Megan K. Creutzburg  
*Portland State University, mkc3@pdx.edu*

Robert M. Scheller  
*Portland State University, rmschell@pdx.edu*

Melissa S. Lucash  
*Portland State University, lucash@pdx.edu*

Louisa B. Evers  
*Bureau of Land Management*

Stephen D. LeDuc  
*United States Environmental Protection Agency*

*See next page for additional authors*

Let us know how access to this document benefits you.

Follow this and additional works at: [http://pdxscholar.library.pdx.edu/esm\\_fac](http://pdxscholar.library.pdx.edu/esm_fac)

 Part of the [Forest Biology Commons](#)

---

## Recommended Citation

Creutzburg, M. K., Scheller, R. M., Lucash, M. S., Evers, L. B., LeDuc, S. D., & Johnson, M. G. (2015). Bioenergy harvest, climate change, and forest carbon in the Oregon Coast Range. GCB Bioenergy.

This Post-Print is brought to you for free and open access. It has been accepted for inclusion in Environmental Science and Management Faculty Publications and Presentations by an authorized administrator of PDXScholar. For more information, please contact [pdxscholar@pdx.edu](mailto:pdxscholar@pdx.edu).

---

**Authors**

Megan K. Creutzburg, Robert M. Scheller, Melissa S. Lucash, Louisa B. Evers, Stephen D. LeDuc, and Mark G. Johnson

Received Date : 01-Oct-2014

Accepted Date : 29-Jan-2015

Article type : Original Research

## **Bioenergy harvest, climate change, and forest carbon in the Oregon Coast Range**

Running head: Bioenergy harvest and climate change

Megan K. Creutzburg<sup>1</sup>, Robert M. Scheller<sup>1</sup>, Melissa S. Lucash<sup>1</sup>, Louisa B. Evers<sup>2</sup>, Stephen D. LeDuc<sup>3</sup>, Mark G. Johnson<sup>4</sup>

<sup>1</sup> Department of Environmental Science and Management, Portland State University, PO Box 751, Portland, OR 97207, USA

<sup>2</sup> Bureau of Land Management, 1220 SW 3rd Avenue, Portland, OR 97204, USA

<sup>3</sup> US Environmental Protection Agency, National Center for Environmental Assessment, 1200 Pennsylvania Ave, NW (8623P), Washington, DC 20460, USA

<sup>4</sup> US Environmental Protection Agency, National Health and Environmental Effects Research Laboratory, 200 SW 35th Street, Corvallis, OR 97333, USA

Corresponding author: telephone 971-217-7066, e-mail [mkc3@pdx.edu](mailto:mkc3@pdx.edu), fax 503-725-9960

Keywords: bioenergy, biomass energy, carbon, climate change, forest, landscape modeling, LANDIS-II, Oregon Coast Range, simulation modeling

Paper type: Original Research

## Abstract

Forests provide important ecological, economic and social services, and recent interest has emerged in the potential for using residue from timber harvest as a source of renewable woody bioenergy. The long-term consequences of such intensive harvest are unclear, particularly as forests face novel climatic conditions over the next century. We used a simulation model to project the long-term effects of management and climate change on above- and belowground carbon storage in a watershed in northwestern Oregon. The multi-ownership watershed has a diverse range of current management practices, including little-to-no harvesting on federal lands, short-rotation clear-cutting on industrial land, and a mix of practices on private non-industrial land. We simulated multiple management scenarios, varying the rate and intensity of harvest, combined with projections of climate change. Our simulations project a wide range of total ecosystem carbon storage with varying harvest rate, ranging from a 45% increase to a 16% decrease compared to current levels. Increasing the intensity of harvest for bioenergy caused an additional 2-3% decrease in ecosystem carbon relative to conventional harvest practices. Soil carbon was relatively insensitive to harvest rotation and intensity, and accumulated slowly regardless of harvest regime. Climate change reduced carbon accumulation in soil and detrital pools due to increasing heterotrophic respiration, and had variable effects on total ecosystem carbon, ranging from a 5% decrease to a 2% increase depending on management scenario.

Overall, we conclude that current levels of ecosystem carbon storage are maintained in part due to substantial portions of the landscape (federal and some private lands) remaining un-harvested or lightly managed. Increasing the intensity of harvest for bioenergy on currently harvested land,

however, led to a small reduction in the ability of forests to store carbon. Climate change is unlikely to substantially alter carbon storage in these forests, absent shifts in disturbance regimes.

## **Introduction**

Forests provide many important ecosystem services, including wildlife habitat, recreation, soil protection, clean air and water, and timber production. As we face unprecedented global challenges in the twenty first century, forests are also increasingly recognized for other services, including the ability to store carbon and mitigate the impacts of climate change (Bonan, 2008, D'Amato *et al.*, 2011, Golden *et al.*, 2011, McKinley *et al.*, 2011) and the potential to provide bioenergy from harvest residue (USDOE 2011, Malmshemer *et al.*, 2011). Bioenergy harvest involves removal of residue such as branches, tops, leaves, small trees, and/or shrubs, along with removing merchantable material as in conventional harvest practices. This harvest residue can be processed to produce electricity or other types of energy (e.g., pellets for wood stoves) from a renewable source of biomass as an alternative to energy from fossil fuels. However, concerns remain over the ability of intensively harvested forests to maintain productivity, sequester carbon, and provide ecosystem services. For example, Harmon and Marks (2002) predicted that removing residue following harvest through prescribed burning substantially lowered the ability of forests to store carbon. Soil nutrient concentrations could also decline due to the removal of additional, nutrient-rich material during bioenergy harvest, and could lead to declining productivity over time (Wall, 2012). Studies are mixed, with evidence for positive, neutral, and negative effects of bioenergy harvest on tree productivity, soil and nutrient pools (reviewed in Thiffault *et al.*, 2011).

Novel climatic conditions may interact with timber harvest in as yet unknown ways. Climate change is increasing temperatures and changing precipitation patterns in the Pacific Northwest (Stocker *et al.*, 2013), and may produce novel conditions not yet experienced by long-lived forest tree species which cannot rapidly migrate or adapt to such changes. Climate change has already caused range shifts and mortality in some forest tree species (Daniels *et al.*, 2011, Hennon *et al.*, 2012). The effects of climate change are likely to increase substantially over the next century, with expected increases in mortality due to insects and disease (Kurz *et al.*, 2008), increased frequency and severity of wildfire (Littell *et al.*, 2010, Westerling *et al.*, 2006), and shifting ranges of tree species (Bachelet *et al.*, 2001, Coops & Waring, 2011). Climate change may also reduce carbon sequestration potential and the ability of forests to mitigate climate impacts (Loudermilk *et al.*, 2013, Rogers *et al.*, 2011). If climate change causes increased stress on trees or declines in productivity, it may exacerbate any negative effects of conventional or bioenergy harvest.

US federal agencies have recently been tasked in an executive order to address potential climate change effects and promote climate resilience on federally administered lands (Executive Order 13653). Many federal lands have been managed for multiple uses, including timber production, wildlife habitat, and recreation, and the recent executive order adds carbon sequestration to the list of values. To make informed management decisions and evaluate the best options for maintaining forest productivity, carbon sequestration, and ecosystem health, an assessment of the long-term effects of management actions and climate change on ecosystem properties is required. As it is impossible to study large-scale and long-term processes experimentally, researchers and managers increasingly rely on simulation models to estimate the long-term

consequences of current practices and guide management decisions. Simulation modeling provides a framework whereby many scenarios, each containing different assumptions about future conditions or actions, can be explored to reduce some of the uncertainty about the future and help inform management (Thompson *et al.*, 2012). Several simulation models have been used in the Pacific Northwest to understand the effects of timber harvest (Harmon *et al.*, 2009, Johnson *et al.*, 2007) and climate change (Coops & Waring, 2011, Hudiburg *et al.*, 2013a, Littell *et al.*, 2010, Rogers *et al.*, 2011), but few have simulated both (Hudiburg *et al.*, 2013b).

In this study, we simulated the effects of forest management and climate change on above- and belowground carbon storage in a northwest Oregon watershed. We explored 49 combinations of management actions and climate projections to examine a wide range of possible future conditions across a multi-owner landscape. Our study questions were: How might varying rate and intensity of timber harvest affect long-term carbon storage in forest vegetation, detritus, and soils of the Oregon Coast Range? What are the likely impacts of climate change on carbon storage under a range of potential future climatic conditions? Will there be interactions between harvesting and climate change?

## **Materials and Methods**

### **Study Area**

This study focuses on the Panther Creek watershed (PCW), on the eastern slope of the Oregon Coast Range Mountains (Figure 1). Forests of the Coast Range are highly productive and are dominated by Douglas-fir (*Pseudotsuga menziesii* (Mirb.) Franco) and western hemlock (*Tsuga heterophylla* (Raf.) Sarg.), which provide high quality timber. The watershed is 7016 hectares in

size but 36% of the area is non-forested, resulting in 4520 hectares of forests simulated for this study. The land ownership and management in the watershed includes private non-industrial forest (PNIF, 44% of the watershed), private industrial forest (PIF, 39%), and public lands administered by the Bureau of Land Management (BLM, 18%). The climate is characterized by wet winters and dry summers, and soils are productive and high in carbon content. In the Coast Range, forests lie within a complex matrix of publically and privately owned lands due to the legacy of historical land development. Historically, these forests were heavily harvested, but passage of the 1994 Northwest Forest Plan dramatically reduced timber harvest and increased carbon storage on federal lands in the region (Krankina *et al.*, 2012). As a result, current stand composition and harvest practices are diverse, ranging from clear-cut harvest on industrial lands to large areas with no or limited harvest on federal lands. The current mix of stand ages range from recent clear-cuts to 300 years, with average stand ages of 46 years on PNIF, 41 years on PIF, and 62 years on BLM lands. Douglas-fir is by far the most common species in the PCW, with other dominant species including (in order of prevalence) bigleaf maple (*Acer macrophyllum* Pursh), red alder (*Alnus rubra* Bong.), Western hemlock, Western redcedar (*Thuja plicata* Donn ex D. Don), Oregon white oak (*Quercus garryana* Douglas ex Hook.), and grand fir (*Abies grandis* (Douglas ex D. Don) Lindl.). The major stand-replacing disturbance in the landscape is timber harvest, with large wildfires occurring on a long rotation interval. Wind throw and fungal diseases such as root rot (*Phellinus weirii*) and Swiss needle cast (*Phaeocryptopus gaeumannii*) can cause mortality and slow growth, but are less prevalent on the eastern slope of the Coast Range, where the PCW is located.

## Simulation Modeling Framework

We used the LANDIS-II forest simulation model (Scheller et al., 2007) to project landscape-scale forest dynamics on a 1 ha grid from 2010-2100. LANDIS-II is a process-based simulation model that represents forest communities as tree species-age cohorts within gridded cells across the landscape. LANDIS-II simulates cohort growth, mortality and regeneration, as dictated by life history and physiological attributes. Species compete for resources within each cell, and disperse spatially across cells within a simulated landscape, therefore allowing for shifts in species ranges. LANDIS-II is freely available on the web ([www.landis-ii.org](http://www.landis-ii.org)), and operates as a core module interacting with extensions, each simulating succession, disturbances, and/or management. We used two extensions for this study: the Century Succession extension and the Leaf Biomass Harvest extension.

The Century Succession extension (Scheller et al., 2011) was derived from the CENTURY Soil Organic Matter model (Parton et al., 1983). In addition to simulating growth, mortality, regeneration and competition (as described above), it estimates above- and belowground net primary production (NPP), net ecosystem exchange (NEE), multiple pools of live and dead tree biomass (including leaf, wood, fine root, coarse root, coarse woody debris) and active, passive and slow pools of soil organic matter (Parton et al., 1983). The extension incorporates monthly temperature and precipitation data that, along with other inputs (e.g., soil texture), influence soil water content and nitrogen available for tree growth. The model does not operate at the photosynthetic level but rather simulates growth and competition as dictated by limitations imposed by temperature, water, nitrogen, leaf area, and growing space. As stands age, cohorts

approach maximum biomass asymptotically. The extension does not currently include CO<sub>2</sub> fertilization effects.

We used Century extension version 3.1.1, in which we made several model adjustments to simulate forests with large trees, productive soils, and high levels of carbon storage. We increased the range of many inputs (e.g., soil organic matter) and reduced the minimum allowable leaf:wood ratio. We also modified nitrogen retranslocation for conifers so that they could utilize resorbed nitrogen throughout the year, not just during spring leaf flush. These alterations represent an improved version of the Century Succession extension that is more suitable for the Pacific Northwest and other temperate coniferous ecosystems.

We used the Leaf Biomass Harvest extension version 2.0.2 (Syphard *et al.*, 2011) to simulate conventional and bioenergy harvest. This extension is based on the Base Harvest extension (Gustafson *et al.*, 2000), simulating a wide variety of harvest prescriptions and allowing the user to specify the amount of woody and leaf material removed from a site.

#### Model Inputs

Inputs to the LANDIS-II model include initial vegetation data, ecoregion inputs, species and functional group traits, management inputs, and climate data. See Tables S1-10 for the Century extension parameter values and data sources.

### *Initial Vegetation Data*

To initialize the simulated landscape with current vegetation information, we used the gradient nearest neighbor (GNN) map for the Oregon Coast Range (map region 223) produced by the Landscape Ecology, Modeling, Mapping and Analysis group for Northwest Forest Plan Effectiveness Monitoring (Ohmann & Gregory, 2002). The GNN method imputes forest inventory plot data to every pixel in the map, characterizing tree species composition, age structure, and many other variables. Inventory plots came from a variety of sources, including the Forest Service Forest Inventory and Analysis (FIA) and Current Vegetation Survey (CVS) programs. From the supplemental TREE\_LIVE database we obtained age information for each individual tree within the imputed forest inventory plots and summarized each plot into species-age cohorts at 10-year age intervals, up to the maximum longevity age for each species.

### *Ecoregion Parameters*

LANDIS-II divides the study area into ecoregions, each of which are assumed to have homogeneous soils and climate. We defined nine ecoregions in the PCW, including three climate regions that captured the precipitation gradient from west (170 cm average annual precipitation) to east (111 cm average annual precipitation), and three soil regions, ranging from high soil organic carbon (SOC) (271 Mg C/ha) to low SOC (135 Mg C/ha). We defined climate regions based on precipitation grids from the PRISM Group (Daly *et al.*, 1997) and soil regions based on the SSURGO National Soil Survey for Yamhill County (Soil Survey Staff, accessed April 5, 2013). Ecoregion parameters included soil properties such as percent clay and sand, SOC decomposition rates, drainage class, as well as initial pools of carbon and nitrogen. We computed soil parameters as a spatially-weighted average to 1m soil depth. Percent clay, percent sand, field

capacity and drainage class were derived directly from the SSURGO database, and wilting point was calculated as field capacity minus available water content. We based initial SOC and soil organic nitrogen pools on data from soil pits collected throughout the PCW (M.G. Johnson *et al.*, manuscript in preparation). Nitrogen inputs were assumed to come from wet and dry deposition, biological fixation in lichens, soil, and decaying logs (Fenn *et al.*, 2003, Johnson *et al.*, 1982, Sollins *et al.*, 1980, Zhang *et al.*, 2012), and fertilization in managed forests. All nitrogen inputs averaged roughly 13 kg N/ha. All ecoregion parameter values and sources are listed in Tables S1-S10.

#### *Species and Functional Group Parameters*

We simulated seven tree species, listed in the Study Area section, but did not simulate any shrub or understory species. The seven simulated species were grouped into four functional groups: conifer-dry (Douglas-fir and grand fir), conifer-mesic (western hemlock and western redcedar), deciduous-dry (Oregon white oak), and deciduous-mesic (red alder and bigleaf maple). All species and functional group parameter values and sources are listed in Tables S1-S10.

#### *Management Data*

Spatial management inputs included a map of management areas (Figure 1) and a stand map. The management area map came from the Integrated Landscape Assessment Project (accessed via <http://westernlandscapesexplorer.info/AccessILAPDataMapsModelsandAnalyses#GIS>). Within the PCW, 44% was managed as PNIF, 39% as PIF, and 18% was managed as by the BLM as an Adaptive Management Area. The PNIF management area was further divided into PNIF-harvested and PNIF-reserve based on surveys by Johnson *et al.* (1999). In the survey, 75% of the

PNIF respondents indicated that timber harvest was important or very important. We randomly selected 25% of the stands (see below) within the PNIF management area, where we excluded harvest to represent PNIF landowners that do not intend to harvest timber on their lands.

Forest stand maps for BLM lands were downloaded from the BLM Oregon/Washington Data Library (accessed via <http://www.blm.gov/or/gis/data.php>) and converted to raster format. Stand maps were unavailable for private land, and therefore we developed a stand map by classifying the current vegetation map into age groups, and iteratively performing majority filter and boundary clean operations in ArcGIS 10.1 to group stands by age classes, remove very small stands and aggregate to observed stand sizes based on surveys (Briggs, 2007, Johnson *et al.*, 1999). In the final stand map, average stand size was 7 ha in PNIF, 13 ha in PIF, and 7 ha on BLM lands.

Inputs for the individual harvest prescriptions included: method for selecting stands for harvest (random for all treatments except BLM thinning, in which the oldest stands within the allowable age range were harvested first); degree of removal (total removal [clear-cut] or percentage of each species-age cohort removed for thinnings); percent harvested per 10-year time step (rotation); species selected for harvest; and species planted following harvest. For all thinning treatments we assumed that 60% of the carbon in the specified cohort age range was removed (unpublished BLM data). All species were harvested except Oregon white oak. When a stand was selected for harvest, all cells within the stand were harvested as allowed by the specific prescription parameters. See Management Scenarios section and Table 1 for information about the harvest regime in each ownership and management scenario.

### *Climate Data*

The Century Succession extension requires monthly temperature and precipitation data for model spin-up (simulating forest succession and carbon accumulation up to 2009) and future projections (years 2010-2100). We obtained climate data from the US Geological Survey GeoData Portal (<http://cida.usgs.gov/gdp/>) as an area-weighted average for each climate region. For model spin-up, we used climate data from the Parameter-elevation Relationships on Independent Slopes Model (PRISM) (Daly *et al.*, 1997) over the period from 1950-2009. See Climate Scenarios section, below, for information about future climate projections.

### *Scenarios*

We developed a suite of scenarios in a factorial design in which we ran all combinations of seven management scenarios and seven climate projections, for a total of 49 scenarios.

### *Management Scenarios*

The seven management scenarios included no harvest, three harvest rotations (current, accelerated, and industrial) and two harvest intensities (conventional harvest and bioenergy harvest). Each ownership type had an individual harvest regime under each scenario as described in Table 1. See Management Data for details about the harvest parameters.

### *Climate Scenarios*

The seven climate scenarios included continuing current climate and six models of climate change. Projections under current climate used PRISM data (Daly *et al.*, 1997) from 1950-2009. Climate change projections came from the Coupled Model Intercomparison Project Phase 5

(CMIP5) for the Continental US from the Intergovernmental Panel on Climate Change (IPCC) Fifth Assessment Report (Taylor *et al.*, 2012). We obtained 800m downscaled climate data from the NEX-DCP30 (Climate Analytics Group and NASA Ames Research Center) dataset (Thrasher *et al.*, 2013) for three global circulation models (GCMs) and two greenhouse gas forcing scenarios, called representative concentration pathways (RCPs). We selected GCMs of future climate change using two criteria: 1) GCMs that were ranked in the top 11 in an assessment of the performance of GCM historical projections compared to observed climate data for the Pacific Northwest (Rupp *et al.*, 2013); and 2) GCMs that spanned a wide range of projected future annual temperature and precipitation for the Pacific Northwest (Table 2). The GCMs chosen were: CanESM2 (Canadian Centre for Climate Modeling and Analysis), projecting hotter and wetter future conditions; CCSM4 (National Center of Atmospheric Research), projecting warmer future conditions with similar precipitation; and HadGEM2 (Met Office Hadley Center), projecting hotter and drier future conditions. For each GCM, we used two RCPs representing varying levels of greenhouse gas forcing, including a low forcing scenario (RCP 4.5) and a high forcing scenario (RCP 8.5).

#### Data Assimilation and Model Calibration

Literature and data were used to calibrate the Century extension for the PCW. We completed a literature review of Pacific Northwest forests to determine expected patterns of growth, carbon accumulation, and NEE with species composition, site type, and stand age (Acker *et al.*, 2002, Campbell *et al.*, 2004, Harcombe *et al.*, 1990, Harmon *et al.*, 2004, Hudiburg *et al.*, 2009, Humphreys *et al.*, 2006, Janisch & Harmon, 2002, Krishnan *et al.*, 2009, Raymond & McKenzie, 2013, Runyon *et al.*, 1994, Smithwick *et al.*, 2002, Sun *et al.*, 2004, Vogt, 1991). To calibrate the

Accepted Article

Century extension, we began with single-cell simulations, iteratively adjusting parameters (e.g., shape parameters for temperature response and moisture sensitivity) to match patterns of growth and NEE in literature and flux towers (Falk *et al.*, 2008, Thomas *et al.*, 2013). Then we calibrated other parameters (e.g., SOC decay rates for each soil pool) across the whole PCW to ensure that starting conditions matched input data and landscape-scale processes were adequately simulated. We used carbon estimates from the GNN maps (derived using the Component Ratio Method) to calibrate our initial aboveground carbon from model spin-up (Figure 2). The following criteria were used to ensure that the final calibration was adequate: 1) initial aboveground carbon was within 10% of GNN estimates across all ecoregions (Figure 2); 2) projected aboveground carbon and aboveground NPP was within the range of values and followed trends found in the literature at each stand age; and 3) initial SOC was within 10% of measured values and SOC accumulated 5-15% in all SOC pools over 90 years without harvest.

#### Simulation Model Runs

We simulated 10-year time steps for years 2010-2100. Each scenario was replicated five times to account for stochastic variability in climate and seedling establishment. Raw values were output by ecoregion and reported values were weighted by area. Due to the large number of scenarios, we combined the climate change projections into three categories for graphical purposes: current climate, low forcing climate change (all three GCMs under the RCP 4.5 forcing scenario) and high forcing climate change (all three GCMs under the RCP 8.5 forcing scenario).

## Results

### *Management*

At the initiation of the simulations, total ecosystem carbon was 500 Mg C/ha, with 27% in aboveground live biomass, 47% in mineral soil, and 7% in aboveground detritus (woody debris and litter). Belowground live and dead biomass encompassed 14% and 5% of total ecosystem carbon, respectively (not reported separately). Without any harvest, projected ecosystem carbon storage in forests of the PCW increased by 224 Mg C/ha (45%), storing up to 724 Mg C/ha in the PCW by the end of the century (Figure 3a). Current harvest rates slightly increased ecosystem carbon storage (10 Mg C/ha [2%]), and accelerated harvest slightly decreased ecosystem carbon storage (18 Mg C/ha [4%]). In the industrial scenario, where clear-cutting was prescribed across the entire watershed, landscape carbon declined by 80 Mg C/ha (16%). Under current harvest rates, a total of 186 Mg C/ha was removed as harvested material over the 90-year simulation, under the accelerated harvest scenario a total of 209 Mg C/ha was removed, and under the industrial scenario 265 Mg C/ha total was removed.

Most of the variation among management scenarios was due to projected differences in aboveground live carbon, which ranged from an increase of 111 Mg C/ha (82%) under no harvest to a decrease of 68 Mg C/ha (50%) under industrial harvest (Figure 3b). Soil carbon accumulated slowly (total increase of 15-26 Mg C/ha [6-11%] over 90 years) in all management scenarios, showing little response to harvest rate except for a slightly faster accumulation in the early years of the simulation followed by a leveling off late-century, under the harvested scenarios (Figure 3c). Aboveground detrital carbon increased by 9 Mg C/ha (25%) without harvest but declined under all harvest scenarios, with greater declines as harvest rate and

intensity increased (up to declines of 22 Mg C/ha [60%] under the industrial-bioenergy scenario) (Figure 3d).

Compared to conventional harvest, bioenergy harvest reduced total ecosystem carbon by 10-12 Mg C/ha (2-3%) at the end of the century (Figure 3a). Aboveground live carbon was unaffected by harvest intensity, but bioenergy harvest caused slower soil carbon accumulation (6-8 Mg C/ha [2-3%] lower levels than conventional harvest) and declines in aboveground detritus (4-6 Mg C/ha [11-17%] decrease relative to conventional harvest) (Figure 3b-d). For both of these pools, the impact of bioenergy increased with harvest rate (i.e., the difference between conventional and bioenergy harvest increased from current to accelerated to industrial harvest rates).

#### *Climate Change*

Carbon continued to accumulate under all climate projections until the end of the century without harvest, though climate change slowed ecosystem carbon accumulation by 38 Mg C/ha (8%) at the end of the century, compared to current climate (Figure 4a). Projected climate change lowered aboveground live carbon by 6-10 Mg C/ha (4-7%), lowered soil carbon by 12 Mg C/ha (5%) and reduced detrital carbon by 7-9 Mg C/ha (13-16%), relative to current climate with no harvest (Figure 4b-d). High climate forcing led to slightly greater reductions in carbon storage than low forcing, but there was high overlap and more variation among GCMs than among forcing scenarios.

In addition to annual trends, we examined the seasonal patterns of growth in response to climate change to better understand the physiological limitations experienced by trees under projected climate change. We examined monthly growth limitations from water and temperature (note that there are other growth limitations in the model not discussed here). By the end of the century, rising temperatures in both summer and winter under all climate change scenarios resulted in a lower temperature limitation and higher growth in winter, spring, and fall months (Figure 5). Conversely, in summer months, increasing limitation from high temperatures combined with water stress reduced summer growth. GCMs varied substantially (Table 2) in the degree of temperature and water limitation, ranging from relatively small changes (CCSM4) to high summer water limitation (HadGEM) and high temperature and water limitation (CanESM).

#### *Management – Climate Change Interactions*

Interactions between management and climate change indicated a lower impact of climate change under harvested scenarios (Figure S1). Although aboveground carbon decreased slightly (6-10 Mg C/ha) under climate change without harvest, it actually increased by a similar amount (6-12 Mg C/ha [6-18%]) under the six harvested scenarios, relative to current climate. The impacts of climate change on soil and detrital carbon did not change sign, but climate change appeared to have a lower impact without harvest. For instance, under the no harvest scenario, soil carbon was reduced by 12 Mg C/ha under climate change relative to current climate, whereas soil carbon declined by only 2-3 Mg C/ha (1%) with climate change under the industrial harvest scenario. Similarly for detrital carbon, climate change caused a decline of 7-9 Mg C/ha relative to no climate change without harvest, but was nearly the same under the industrial harvest scenario. Taken together, the overall decline in ecosystem carbon accumulation with climate

change projected under no harvest (Figure 4a) disappeared in the harvested scenarios, and even slightly reversed (increase of 10 Mg C/ha [2%]) in the industrial harvest scenarios.

## **Discussion**

Many questions related to sustainability in forested landscapes require information about complex, interacting processes over long time frames. Evaluating the comparative effects of multiple harvest practices has important implications for long-term forest management, particularly when anticipating climate change (Thompson *et al.*, 2012). In this study, we used a simulation model to explore 49 scenarios to bracket a large range of potential future conditions and assess their implications for forest carbon storage. We simulated varying management practices in a heterogeneous watershed, encompassing an intermixture of ownership types with very different management strategies. These variations in management practices had important implications for carbon storage, as the balance of harvested timberlands and lightly or unharvested areas determined whether carbon would accumulate, decline, or maintain current levels across the landscape.

## *Management*

The management scenarios considered in this study varied widely, ranging from managing for maximum carbon storage (no harvest) to industrial harvest across all lands (industrial scenario), with multiple scenarios in between. Ownership patterns in the Coast Range are a mosaic of federal and private land, currently managed very differently due to restrictions placed on federal lands resulting from the Northwest Forest Plan. The current and accelerated harvest scenarios reflect this heterogeneity in management across the PCW, and projections of ecosystem carbon

storage in the PCW under current and accelerated harvest rates suggest that these scenarios are likely to maintain levels of carbon storage similar to those currently found in these forests.

However, under the industrial scenario, carbon storage declined relative to current levels. In the industrial scenario, the entire watershed was harvested similar to PIF, with an additional 29% of the landscape available for timber harvest that was un-harvested or lightly harvested in the other scenarios. Our findings suggest that federal lands and other non-industrial private lands provide an important counter-balance to intensive industrial forestry in Coast Range forests. If enough lightly harvested or un-harvested land remains on the landscape, carbon storage can be maintained even with intensive private industrial management practiced on some lands.

However, if there were major changes to federal forest policy or if more private non-industrial landowners were to start harvesting for timber, forest ecosystem carbon storage may decline.

Harvest rate (current, accelerated, and industrial) had the expected impact on aboveground carbon; as more trees were removed, aboveground carbon declined. In contrast, harvest rate had little effect on soils, the largest carbon pool in the PCW and many other heavily managed forests in the Pacific Northwest. Projected soil carbon increased slowly over time under all scenarios, as organic material from plant biomass accumulated in soils (Kelly *et al.*, 1997). Soil carbon initially increased at a slightly higher rate under harvested scenarios, as roots from harvested trees began to decompose and contribute to SOC, but leveled off later in the century. Although an increase in soil carbon with harvest may be counterintuitive, it is not unexpected, as reviews have shown that harvest impacts on soil carbon can be positive, neutral, or negative (Johnson & Curtis, 2001, Nave *et al.*, 2010). This pattern is also consistent with some studies that found increases in SOC with harvest of coniferous species (Johnson & Curtis, 2001). Increasing harvest

rate and intensity also reduced aboveground detrital carbon, consistent with much of the literature (Johnson & Curtis, 2001, Nave *et al.*, 2010). Because soil carbon represents such a large carbon pool and is relatively resilient to management impacts, the stable soil carbon pool buffers the overall impact of harvest on total ecosystem carbon.

Our simulations suggest residue harvest for bioenergy (changing harvest intensity) would likely have little additional effect on total ecosystem carbon storage, although it does reduce soil and detrital carbon storage compared to conventional harvest. Also, although impacts of bioenergy were small, they appeared to increase with faster harvest rotation, indicating that there might be more concern about the sustainability of bioenergy harvest in the most frequently harvested plantations. Some studies in the Pacific Northwest have found that bioenergy harvest does not reduce forest productivity or SOC (Holub *et al.*, 2013, Knight, 2013), but others document negative impacts (Proe & Dutch, 1994, Wall, 2012, Walmsley *et al.*, 2009). Our results were not sensitive to the amount of residue harvested, as simulations varying levels of wood and leaf removal up to 100% of all plant material (data not shown) showed similar impacts. However, we did not simulate other nutrients besides nitrogen, and repeated bioenergy harvest could make other soil nutrients such as calcium, phosphorus, or potassium limiting in the long-term (Thiffault *et al.*, 2011). In reality, bioenergy harvest can vary widely in intensity, from removal of tops only with branch and leaves left on site, to whole-tree harvest, to complete removal including stumps. These various bioenergy practices can have very different ecosystem impacts. Our simulations under conventional harvest assume that most of the detrital material is left on site, even though site preparation can remove much of the detrital material prior to replanting.

Therefore, in some cases, conventional harvesting may actually have site impacts more similar to bioenergy harvest.

In this study, we only report the ecosystem impacts of harvest, and did not attempt to quantify the overall climate change feedbacks associated with conventional or bioenergy harvest. For instance, we do not quantify emissions from transportation of wood products, conversion of harvest residue to usable energy sources, ability to substitute for fossil fuels as an energy source, and many other considerations needed to determine the full implications of bioenergy harvest. It is also important to note that some of the carbon conventionally harvested in the landscape is used in long-lived structures (e.g., buildings) and can provide long-term carbon storage off-site.

Many researchers have evaluated the full carbon cycle implications of bioenergy harvest in mitigating climate change, and its impacts on forested landscapes throughout the world (e.g., de Jong *et al.*, 2007, Kaul *et al.*, 2010, Mika & Keeton, 2013, Winford & Gaither, 2012, Zanchi *et al.*, 2012). Recent studies in the Pacific Northwest indicate that bioenergy harvest is unlikely to offset greenhouse gas emissions as a climate change mitigation strategy (Hudiburg *et al.*, 2011, Hudiburg *et al.*, 2013b, Mitchell *et al.*, 2012, Schulze *et al.*, 2012), but impacts likely vary regionally (Winford & Gaither, 2012). There remains interest in forest bioenergy production as part of a climate change mitigation strategy across the United States and other parts of the world (U.S. Department of Energy, 2011, IPCC, 2014).

#### *Climate Change*

Climate change is expected to have major consequences for forested ecosystems over the next century (Bonan, 2008, IPCC, 2014, Vose *et al.*, 2012). In the Pacific Northwest, expected

changes include increased summer drought stress, shifting species ranges, and increasing disturbance frequency (Mote *et al.*, 2014). Our simulations indicate that climate change may slightly lower carbon storage potential in the PCW, mostly driven by losses of soil and detrital material to heterotrophic respiration. However, the balance of production and respiration varied among management scenarios, and scenarios with timber harvest tended to sustain lower respiration-related carbon losses. Studies have shown that surficial soil respiration increases with soil warming (Rustad *et al.*, 2001), but the responsiveness of resistant soil organic matter to temperature is still unclear (Davidson & Janssens, 2006).

All climate models projected higher annual temperatures by the end of the century, with winter minimum temperatures rising 2.3-6.1°C and summer maximum temperatures increasing by 3.4-9.7°C (Table 2). All climate models also projected greater winter precipitation in the PCW, although projections of summer precipitation varied from drier to wetter depending on the climate model. Warming temperatures resulted in a longer growing season for coniferous species that retain leaves throughout the year and are currently limited primarily by temperature in the winter, early spring and late fall. However, increasing cool-season productivity was counterbalanced by declining production in the summer due to heat and drought stress (Figure 5), as predicted in other studies of Pacific Northwest conifers (Beedlow *et al.*, 2013, Chmura *et al.*, 2011, Littell *et al.*, 2008). The climate scenarios with the greatest increases in winter production also showed the greatest productivity declines in summer months, with the result of largely canceling out variation among climate scenarios. Therefore, annual levels of aboveground live carbon and productivity were affected very little by climate change.

Climate change may affect Coast Range forests in additional ways not considered in this study.

The greatest impacts of climate change in Pacific Northwest forests may be from increasing disturbance frequency and intensity (Chmura *et al.*, 2011, Raymond & McKenzie, 2012), which we did not simulate in this study. Wildfire frequency is expected to increase with climate change (Littell *et al.*, 2010, Raymond & McKenzie, 2012, Rogers *et al.*, 2011), and may interact with other disturbances (e.g., insects and disease) to shape future forests in the region. Additionally, many Pacific Northwest conifers, including Douglas-fir and western hemlock, require winter chilling for normal bud burst and growth. Increasing winter temperatures under climate change may not provide enough cold days for continued normal growth, flowering and seed germination (Chmura *et al.*, 2011, Cumming & Burton, 1996), but we were not able to model this effect. We also did not simulate CO<sub>2</sub> fertilization, which will likely increase production under climate change due to increased photosynthetic rates and water use efficiency (Coops & Waring, 2001, Hudiburg *et al.*, 2013b, Keenan *et al.*, 2013, Norby *et al.*, 2005). Therefore, we may be underestimating production under elevated atmospheric [CO<sub>2</sub>] with climate change, although at least one experimental study found no significant effect of rising atmospheric [CO<sub>2</sub>] on Douglas-fir growth (Olszyk *et al.*, 1998).

#### *Uncertainty and Model Limitations*

There are many sources of uncertainty in simulation modeling, beginning with uncertainty in model parameters and assumptions. The Century Succession extension simulates a wide range of processes and therefore requires a large number of input parameters. Due to the rich history of ecological studies in Pacific Northwest forests, we were able to obtain field-based data for many parameters (Tables S1-S10). However, some values were not available in the literature and

others had a wide range of variability in their estimates. Additionally, projections of climate change are inherently uncertain in many ways, including uncertainty about greenhouse gas emissions levels, the climate forcing resulting from those emissions, localized climatic effects resulting from global patterns, and the impacts of changing climatic conditions on individual species and interspecific interactions (Knutti & Sedlacek, 2013). We intentionally chose several climate models to encompass much of the likely range of future climate conditions in the Coast Range, but the range of actual uncertainty is much higher than is captured in our projections.

Calibration is another major challenge with the use of simulation models, as few data sources are generally available for calibration, and variability and uncertainty in available data sets are often high. We used a set of criteria to ensure that model calibration was adequate (see Data Assimilation and Calibration), but our simulations highlighted areas for potential improvement in the LANDIS-II Century Succession extension. The model tended to underestimate summer productivity and heterotrophic respiration, appearing to be overly sensitive to the dry summers experienced in the Pacific Northwest and underestimating soil water holding capacity. This appears to be a common limitation among multiple simulation models used in the Pacific Northwest (Hudiburg *et al.*, 2013a, Schaefer *et al.*, 2012). However, our aim was to adequately simulate processes over long time frames and compare outcomes of multiple scenarios, rather than precisely predict seasonal patterns. The LANDIS-II model also simplifies the modeling of management activities to harvest practices that affect entire species-age cohorts. Although we can model a wide range of silvicultural practices and species-specific management activities, we cannot simulate more subtle changes in silviculture, such as retention of individual trees or snags, which can be important for wildlife habitat.

## **Conclusions**

In this study, we used a simulation modeling framework to explore a wide range of future management actions and climatic conditions across a multi-ownership watershed. Our results indicate that maintenance of current carbon storage levels are possible under current practices partially due to un-harvested federal and non-industrial private lands that counterbalance intensive forestry operations practiced on private industrial lands. We also find that the ecosystem impacts of bioenergy harvest are likely to be minor, suggesting that bioenergy could potentially provide a low impact, renewable energy source in the region if markets and processing facilities become available. Soils contained the largest reservoir of carbon in the PCW and were relatively resilient to the impacts of harvest, although carbon accumulation in soils slowed under climate change due to increasing heterotrophic respiration. Overall, management impacts were more influential on landscape condition than climate change, which caused relatively small declines in carbon accumulation in some pools. However, indirect effects of climate change, such as changes to disturbance regimes (e.g., increases in wildfire) warrant further study across larger landscapes.

## **Acknowledgments**

Funding was provided by the U.S. Bureau of Land Management and Environmental Protection Agency Interagency Agreement DW-14-95829801-0. Thanks to George McFadden of the Bureau of Land Management, and Greg Johnson, David Marshall, and Scott Holub of Weyerhaeuser for project input. Thanks to the Landscape Ecology, Modeling, Mapping and Analysis team for providing vegetation maps and the Ameriflux network for providing flux tower data. Special thanks to the members of the Dynamic Ecosystem and Landscapes Lab at

Portland State University for help and feedback throughout this study, with particular thanks to Alec Kretchun and Matthew Duveneck for their willingness to share their expertise in LANDIS-II modeling. Two anonymous reviewers provided comments that improved an earlier version of this manuscript. The views expressed here are those of the authors and do not necessarily represent the views or policies of the U.S. Environmental Protection Agency or Bureau of Land Management.

## References

- Acker SA, Halpern CB, Harmon ME, Dyrness CT (2002) Trends in bole biomass accumulation, net primary production and tree mortality in *Pseudotsuga menziesii* forests of contrasting age. *Tree Physiology*, **22**, 213-217.
- Bachelet D, Neilson RP, Lenihan JM, Drapek RJ (2001) Climate change effects on vegetation distribution and carbon budget in the United States. *Ecosystems*, **4**, 164-185.
- Beedlow PA, Lee EH, Tingey DT, Waschmann RS, Burdick CA (2013) The importance of seasonal temperature and moisture patterns on growth of Douglas-fir in western Oregon, USA. *Agricultural and Forest Meteorology*, **169**, 174-185.
- Bonan GB (2008) Forests and climate change: Forcings, feedbacks, and the climate benefits of forests. *Science*, **320**, 1444-1449.
- Briggs D (2007) Management practices on Pacific Northwest west-side industrial forest lands, 1991-2005: with projections to 2010. pp 84, Stand Management Cooperative Working Paper Number 6. College of Forest Resources, University of Washington.
- Campbell JL, Sun OJ, Law BE (2004) Disturbance and net ecosystem production across three climatically distinct forest landscapes. *Global Biogeochemical Cycles*, **18**.
- Chmura DJ, Anderson PD, Howe GT *et al.* (2011) Forest responses to climate change in the northwestern United States: Ecophysiological foundations for adaptive management. *Forest Ecology and Management*, **261**, 1121-1142.
- Coops NC, Waring RH (2001) Estimating forest productivity in the eastern Siskiyou Mountains of southwestern Oregon using a satellite driven process model, 3-PGS. *Canadian Journal of Forest Research*, **31**, 143-154.
- Coops NC, Waring RH (2011) Estimating the vulnerability of fifteen tree species under changing climate in Northwest North America. *Ecological Modelling*, **222**, 2119-2129.
- Cumming SG, Burton PJ (1996) Phenology-mediated effects of climatic change on some simulated British Columbia forests. *Climatic Change*, **34**, 213-222.
- D'amato AW, Bradford JB, Fraver S, Palik BJ (2011) Forest management for mitigation and adaptation to climate change: Insights from long-term silviculture experiments. *Forest Ecology and Management*, **262**, 803-816.
- Daly C, Taylor GH, Gibson WP (1997) The PRISM approach to mapping precipitation and temperature. In: *Proceedings of the 10th American Meteorological Society conference on applied climatology*. pp 10-12, Reno, Nevada, USA, American Meteorological Society.

- Daniels LD, Maertens TB, Stan AB, McCloskey SPJ, Cochrane JD, Gray RW (2011) Direct and indirect impacts of climate change on forests: three case studies from British Columbia. *Canadian Journal of Plant Pathology*, **33**, 108-116.
- Davidson EA, Janssens IA (2006) Temperature sensitivity of soil carbon decomposition and feedbacks to climate change. *Nature*, **440**, 165-173.
- De Jong BH, Masera O, Olguin M, Martinez R (2007) Greenhouse gas mitigation potential of combining forest management and bioenergy substitution: A case study from Central Highlands of Michoacan, Mexico. *Forest Ecology and Management*, **242**, 398-411.
- U.S. Department of Energy (2011) U.S. Billion-Ton Update: Biomass Supply for a Bioenergy and Bioproducts Industry. R.D. Perlack and B.J. Stokes (Leads), ORNL/TM-2011/224. pp 227, Oak Ridge, TN, Oak Ridge National Laboratory.
- Falk M, Wharton S, Schroeder M, Ustin S, U KTP (2008) Flux partitioning in an old-growth forest: seasonal and interannual dynamics. *Tree Physiology*, **28**, 509-520.
- Fenn ME, Haeuber R, Tonnesen GS *et al.* (2003) Nitrogen emissions, deposition, and monitoring in the western United States. *BioScience*, **53**, 391-403.
- Golden DM, Smith MA, Colombo SJ (2011) Forest carbon management and carbon trading: A review of Canadian forest options for climate change mitigation. *Forestry Chronicle*, **87**, 625-635.
- Gustafson EJ, Shifley SR, Mladenoff DJ, Nimerfro KK, He HS (2000) Spatial simulation of forest succession and timber harvesting using LANDIS. *Canadian Journal of Forest Research*, **30**, 32-43.
- Harcombe PA, Harmon ME, Greene SE (1990) Changes in Biomass and Production over 53 Years in a Coastal Picea-Sitichensis - Tsuga-Heterophylla Forest Approaching Maturity. *Canadian Journal of Forest Research*, **20**, 1602-1610.
- Harmon ME, Bible K, Ryan MG, Shaw DC, Chen H, Klopatek J, Li X (2004) Production, respiration, and overall carbon balance in an old-growth Pseudotsuga-tsuga forest ecosystem. *Ecosystems*, **7**, 498-512.
- Harmon ME, Marks B (2002) Effects of silvicultural practices on carbon stores in Douglas-fir-western hemlock forests in the Pacific Northwest, USA: results from a simulation model. *Canadian Journal of Forest Research*, **32**, 863-877.
- Harmon ME, Moreno A, Domingo JB (2009) Effects of Partial Harvest on the Carbon Stores in Douglas-fir/Western Hemlock Forests: A Simulation Study. *Ecosystems*, **12**, 777-791.
- Hennon PE, D'amore DV, Schaberg PG, Wittwer DT, Shanley CS (2012) Shifting Climate, Altered Niche, and a Dynamic Conservation Strategy for Yellow-Cedar in the North Pacific Coastal Rainforest. *BioScience*, **62**, 147-158.
- Holub SM, Terry TA, Harrington CA, Harrison RB, Meade R (2013) Tree growth ten years after residual biomass removal, soil compaction, tillage, and competing vegetation control in a highly-productive Douglas-fir plantation. *Forest Ecology and Management*, **305**, 60-66.
- Hudiburg T, Law B, Turner DP, Campbell J, Donato DC, Duane M (2009) Carbon dynamics of Oregon and Northern California forests and potential land-based carbon storage. *Ecological Applications*, **19**, 163-180.
- Hudiburg TW, Law BE, Thornton PE (2013a) Evaluation and improvement of the Community Land Model (CLM4) in Oregon forests. *Biogeosciences*, **10**, 453-470.
- Hudiburg TW, Law BE, Wirth C, Luyssaert S (2011) Regional carbon dioxide implications of forest bioenergy production. *Nature Climate Change*, **1**, 419-423.
- Hudiburg TW, Luyssaert S, Thornton PE, Law BE (2013b) Interactive Effects of Environmental Change and Management Strategies on Regional Forest Carbon Emissions. *Environmental Science & Technology*, **47**, 13132-13140.

- Humphreys ER, Black TA, Morgenstern K, Cai TB, Drewitt GB, Nestic Z, Trofymow JA (2006) Carbon dioxide fluxes in coastal Douglas-fir stands at different stages of development after clearcut harvesting. *Agricultural and Forest Meteorology*, **140**, 6-22.
- Intergovernmental Panel on Climate Change (IPCC) (2014) Summary for Policymakers. In: *Climate Change 2014, Mitigation of Climate Change. Contribution of Working Group III to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change* (eds Edenhofer O, Pichs-Madruga R, Sokona Y, Farahani E, Kadner S, Seyboth K, Adler A, Baum I, Brunner S, Eickemeier P, Kriemann B, Savolainen J, Schlömer S, Von Stechow C, Zwickel T, Minx JC). Cambridge, United Kingdom and New York, NY, USA, Cambridge University Press.
- Janisch JE, Harmon ME (2002) Successional changes in live and dead wood carbon stores: implications for net ecosystem productivity. *Tree Physiology*, **22**, 77-89.
- Johnson DW, Cole DW, Bledsoe CS *et al.* (1982) Chapter 7: Nutrient Cycling in Forests of the Pacific Northwest. In: *Analysis of Coniferous Forest Ecosystems in the Western United States*. (ed Edmonds RL) pp 186-232. Stroudsburg, PA, Hutchinson Ross Publishing Company.
- Johnson DW, Curtis PS (2001) Effects of forest management on soil C and N storage: meta analysis. *Forest Ecology and Management*, **140**, 227-238.
- Johnson KN, Bettinger P, Kline JD *et al.* (2007) Simulating forest structure, timber production, and socioeconomic effects in a multi-owner province. *Ecological Applications*, **17**, 34-47.
- Johnson R, Alig R, Kline J, Moulton R, Rickenbach M (1999) Management of Non-industrial Private Forest Lands: Survey Results from Western Oregon and Washington Owners. Research Contribution 28, Forest Research Laboratory, Oregon State University, Corvallis.
- Kaul M, Mohren GMJ, Dadhwal VK (2010) Carbon storage versus fossil fuel substitution: a climate change mitigation option for two different land use categories based on short and long rotation forestry in India. *Mitigation and Adaptation Strategies for Global Change*, **15**, 395-409.
- Keenan TF, Hollinger DY, Bohrer G, Dragoni D, Munger JW, Schmid HP, Richardson AD (2013) Increase in forest water-use efficiency as atmospheric carbon dioxide concentrations rise. *Nature*, **499**, 324-327.
- Kelly RH, Parton WJ, Crocker GJ *et al.* (1997) Simulating trends in soil organic carbon in long-term experiments using the century model. *Geoderma*, **81**, 75-90.
- Knight EJ (2013) Harvest intensity and competing vegetation control have little effect on soil carbon and nitrogen pools in a Pacific Northwest Douglas-fir plantation. Unpublished Master of Science University of Washington, Seattle, WA, 37 pp.
- Knutti R, Sedlacek J (2013) Robustness and uncertainties in the new CMIP5 climate model projections. *Nature Climate Change*, **3**, 369-373.
- Krankina ON, Harmon ME, Schnekenburger F, Sierra CA (2012) Carbon balance on federal forest lands of Western Oregon and Washington: The impact of the Northwest Forest Plan. *Forest Ecology and Management*, **286**, 171-182.
- Krishnan P, Black TA, Jassal RS, Chen BZ, Nestic Z (2009) Interannual variability of the carbon balance of three different-aged Douglas-fir stands in the Pacific Northwest. *Journal of Geophysical Research-Biogeosciences*, **114**.
- Kurz WA, Dymond CC, Stinson G *et al.* (2008) Mountain pine beetle and forest carbon feedback to climate change. *Nature*, **452**, 987-990.
- Littell JS, Oneil EE, Mckenzie D, Hicke JA, Lutz JA, Norheim RA, Elsner MM (2010) Forest ecosystems, disturbance, and climatic change in Washington State, USA. *Climatic Change*, **102**, 129-158.
- Littell JS, Peterson DL, Tjoelker M (2008) Douglas-fir growth in mountain ecosystems: Water limits tree growth from stand to region. *Ecological Monographs*, **78**, 349-368.

- Loudermilk EL, Scheller RM, Weisberg PJ, Yang J, Dilts TE, Karam SL, Skinner C (2013) Carbon dynamics in the future forest: the importance of long-term successional legacy and climate-fire interactions. *Global Change Biology*, **19**, 3502-3515.
- Malmsheimer RW, Bowyer JL, Fried JS *et al.* (2011) Managing Forests because Carbon Matters: Integrating Energy, Products, and Land Management Policy. *Journal of Forestry*, **109**, S7-S48.
- Mckinley DC, Ryan MG, Birdsey RA *et al.* (2011) A synthesis of current knowledge on forests and carbon storage in the United States. *Ecological Applications*, **21**, 1902-1924.
- Mika AM, Keeton WS (2013) Factors contributing to carbon fluxes from bioenergy harvests in the U.S. Northeast: an analysis using field data. *Global Change Biology Bioenergy*, **5**, 290-305.
- Mitchell SR, Harmon ME, O'connell KEB (2012) Carbon debt and carbon sequestration parity in forest bioenergy production. *Global Change Biology Bioenergy*, **4**, 818-827.
- Mote P, Snover AK, Capalbo S *et al.* (2014) Ch. 21: Northwest. In: *Climate Change Impacts in the United States: The Third National Climate Assessment*. (eds Melillo JM, Richmond TC, Yohe GW). U.S. Global Change Research Program.
- Nave LE, Vance ED, Swanston CW, Curtis PS (2010) Harvest impacts on soil carbon storage in temperate forests. *Forest Ecology and Management*, **259**, 857-866.
- Norby RJ, Delucia EH, Gielen B *et al.* (2005) Forest response to elevated CO<sub>2</sub> is conserved across a broad range of productivity. *Proceedings of the National Academy of Sciences of the United States of America*, **102**, 18052-18056.
- Ohmann JL, Gregory MJ (2002) Predictive mapping of forest composition and structure with direct gradient analysis and nearest-neighbor imputation in coastal Oregon, USA. *Canadian Journal of Forest Research*, **32**, 725-741.
- Olszyk D, Wise C, Vaness E, Tingey D (1998) Elevated temperature but not elevated CO<sub>2</sub> affects long-term patterns of stem diameter and height of Douglas-fir seedlings. *Canadian Journal of Forest Research*, **28**, 1046-1054.
- Parton WJ, Anderson DW, Cole CV, Steward JWB (1983) Simulation of soil organic matter formation and mineralization in semiarid agroecosystems. Special Publication No. 23. In: *Nutrient cycling in agricultural ecosystems*. (eds Lowrance RR, Todd RL, Asmussen LE, Leonard RA). Athens, Georgia, The University of Georgia, College of Agriculture Experiment Stations.
- Proe MF, Dutch J (1994) Impact of whole-tree harvesting on second rotation growth of sitka spruce: the first 10 years. *Forest Ecology and Management*, **66**, 39-54.
- Raymond CL, Mckenzie D (2012) Carbon dynamics of forests in Washington, USA: 21st century projections based on climate-driven changes in fire regimes. *Ecological Applications*, **22**, 1589-1611.
- Raymond CL, Mckenzie D (2013) Temporal carbon dynamics of forests in Washington, US: Implications for ecological theory and carbon management. *Forest Ecology and Management*, **310**, 796-811.
- Rogers BM, Neilson RP, Drapek R, Lenihan JM, Wells JR, Bachelet D, Law BE (2011) Impacts of climate change on fire regimes and carbon stocks of the U.S. Pacific Northwest. *Journal of Geophysical Research-Biogeosciences*, **116**, G03037.
- Runyon J, Waring RH, Goward SN, Welles JM (1994) Environmental Limits on Net Primary Production and Light-Use Efficiency across the Oregon Transect. *Ecological Applications*, **4**, 226-237.
- Rupp DE, Abatzoglou JT, Hegewisch KC, Mote PW (2013) Evaluation of CMIP5 20th century climate simulations for the Pacific Northwest USA. *Journal of Geophysical Research: Atmospheres*, **118**, 1-23.
- Rustad LE, Campbell JL, Marion GM *et al.* (2001) A meta-analysis of the response of soil respiration, net nitrogen mineralization, and aboveground plant growth to experimental ecosystem warming. *Oecologia*, **126**, 543-562.

- Schaefer K, Schwalm CR, Williams C *et al.* (2012) A model-data comparison of gross primary productivity: Results from the North American Carbon Program site synthesis. *Journal of Geophysical Research-Biogeosciences*, **117**, 1-15.
- Scheller RM, Domingo JB, Sturtevant BR, Williams JS, Rudy A, Gustafson EJ, Mladenoff DJ (2007) Design, development, and application of LANDIS-II, a spatial landscape simulation model with flexible spatial and temporal resolution. *Ecological Modelling*, **201**, 409-419.
- Scheller RM, Hua D, Bolstad PV, Birdsey RA, Mladenoff DJ (2011) The effects of forest harvest intensity in combination with wind disturbance on carbon dynamics in Lake States mesic forests. *Ecological Modelling*, **222**, 144-153.
- Schulze ED, Korner CI, Law BE, Haberl H, Luyssaert S (2012) Large-scale bioenergy from additional harvest of forest biomass is neither sustainable nor greenhouse gas neutral. *Global Change Biology Bioenergy*, **4**, 611-616.
- Smithwick EaH, Harmon ME, Remillard SM, Acker SA, Franklin JF (2002) Potential upper bounds of carbon stores in forests of the Pacific Northwest. *Ecological Applications*, **12**, 1303-1317.
- Sollins P, Grier CC, Mccorison FM, Cromack K, Fogel R (1980) The Internal Element Cycles of an Old-Growth Douglas-Fir Ecosystem in Western Oregon. *Ecological Monographs*, **50**, 261-285.
- Soil Survey Staff, Natural Resources Conservation Service, United States Department of Agriculture, Web Soil Survey. Available online at <http://websoilsurvey.nrcs.usda.gov/>. Accessed 03/20/2013.
- Stocker TF, Qin D, Plattner G-K *et al.* (2013) *The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*, Cambridge, United Kingdom and New York, NY, USA, Cambridge University Press.
- Sun OJ, Campbell J, Law BE, Wolf V (2004) Dynamics of carbon stocks in soils and detritus across chronosequences of different forest types in the Pacific Northwest, USA. *Global Change Biology*, **10**, 1470-1481.
- Syphard AD, Scheller RM, Ward BC, Spencer WD, Strittholt JR (2011) Simulating landscape-scale effects of fuels treatments in the Sierra Nevada, California. *International Journal of Fire Management*, **20**, 364-383.
- Taylor KE, Stouffer RJ, Meehl GA (2012) An overview of CMIP5 and the experiment design. *Bulletin of the American Meteorological Society*, **93**, 485-498.
- Thiffault E, Hannam KD, Pare D, Titus BD, Hazlett PW, Maynard DG, Brais S (2011) Effects of forest biomass harvesting on soil productivity in boreal and temperate forests - A review. *Environmental Reviews*, **19**, 278-309.
- Thomas CK, Martin JG, Law BE, Davis K (2013) Toward biologically meaningful net carbon exchange estimates for tall, dense canopies: Multi-level eddy covariance observations and canopy coupling regimes in a mature Douglas-fir forest in Oregon. *Agricultural and Forest Meteorology*, **173**, 14-27.
- Thompson JR, Wiek A, Swanson FJ *et al.* (2012) Scenario studies as a synthetic and integrative research activity for long-term ecological research. *BioScience*, **62**, 367-376.
- Thrasher B, Xiong J, Wang W, Melton F, Michaelis A, Nemani R (2013) Downscaled climate projections suitable for resource management. *Eos, Transactions American Geophysical Union*, **94**, 321-323.
- Vogt K (1991) Carbon Budgets of Temperate Forest Ecosystems. *Tree Physiology*, **9**, 69-86.
- Vose JM, Peterson DL, Patel-Weynand T (2012) *Effects of climatic variability and change on forest ecosystems: a comprehensive science synthesis for the U.S. forest sector. Gen. Tech. Rep. PNW-GTR-870*, U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station, Portland, OR.
- Wall A (2012) Risk analysis of effects of whole-tree harvesting on site productivity. *Forest Ecology and Management*, **282**, 175-184.

- Accepted Article
- Walmsley JD, Jones DL, Reynolds B, Price MH, Healey JR (2009) Whole tree harvesting can reduce second rotation forest productivity. *Forest Ecology and Management*, **257**, 1104-1111.
- Westerling AL, Hidalgo HG, Cayan DR, Swetnam TW (2006) Warming and earlier spring increase western U.S. forest wildfire activity. *Science*, **313**, 940-943.
- Winford EM, Gaither JC (2012) Carbon outcomes from fuels treatment and bioenergy production in a Sierra Nevada forest. *Forest Ecology and Management*, **282**, 1-9.
- Zanchi G, Pena N, Bird N (2012) Is woody bioenergy carbon neutral? A comparative assessment of emissions from consumption of woody bioenergy and fossil fuel. *Global Change Biology Bioenergy*, **4**, 761-772.
- Zhang L, Jacob DJ, Knipping EM *et al.* (2012) Nitrogen deposition to the United States: distribution, sources, and processes. *Atmospheric Chemistry and Physics*, **12**, 4539-4554.
- Zhou X, Hemstrom MA (2009) Estimating Aboveground Tree Biomass on Forest Land in the Pacific Northwest. Research Paper PNW-RP-584. pp 18, Portland, OR, U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station.

### Supporting Information Legends

Supporting Information Tables S1-S10. Model Parameters and Data Sources.

Figure S1. Interactive effects of management and climate scenarios on major ecosystem carbon pools. Projections are shown for current climate (left), and three global circulation models (GCMs) under a low forcing scenario (RCP 4.5, middle) and high forcing scenario (RCP 8.5, right).

**Tables**

<b>Scenario Name</b>	<b>Harvest Rotation</b>	<b>Harvest Intensity</b>
No Harvest	No harvest on all lands.	No harvest on all lands.
Current–Conventional (business as usual)	PIF: Clear-cut on a 50 year rotation, with planting of Douglas-fir following harvest (Briggs, 2007). PNIF: 25% of stands reserved without any harvest (Johnson <i>et al.</i> , 1999); 75% of stands thinned at 20-40 years and clear-cut on a 60 year rotation. BLM: commercial thinning at 40-80 years on a 100 year harvest schedule; no harvest in stands >80 years.	Removed 80% of wood (Zhou & Hemstrom, 2009); remaining 20% of wood and all leaves left on site.
Current–Bioenergy	Same as Current–Conventional.	Removed 96% of wood and 80% of leaves.
Accelerated–Conventional	PIF: Clear-cut on a 40 year rotation, with planting of Douglas-fir following harvest. PNIF: 25% of stands reserved without any harvest; 75% of stands thinned at 20-40 years and clear-cut on a 50 year rotation. BLM: commercial thinning at 40-100 years on a 60 year harvest schedule; no harvest in stands >160 years.	Same as Current–Conventional.
Accelerated–Bioenergy	Same as Accelerated–Conventional.	Same as Current–Bioenergy.
Industrial–Conventional	All lands (PIF, PNIF, BLM) harvested with clear-cut on a 50 year rotation, with planting of Douglas-fir following harvest.	Same as Current–Conventional.
Industrial–Bioenergy	Same as Industrial–Conventional.	Same as Current–Bioenergy.

Table 1. Description of the seven management scenarios modeled in the Panther Creek watershed by ownership category, where applicable. Ownership categories are PIF (private industrial forest), PNIF (private non-industrial forest), and BLM (Bureau of Land Management).

Table 2. Temperature and precipitation projections under current climate (years 1950-2009, top row) and six climate change scenarios for the Panther Creek watershed projected at the end of the century (2091-2100). Future climate projections are comprised of a global circulation model (CCSM4, HadGEM, and CanESM) and a representative concentration pathway (4.5 [low forcing] and 8.5 [high forcing]). Values are shown annually and for winter months (December, January and February) and summer months (June, July and August). Tmax values report maximum monthly temperature (°C) averaged across years, Tmin values are minimum monthly temperature (°C) averaged across years, and Ppt is total precipitation (cm) averaged across years.

Climate scenario	Annual			Winter months			Summer months		
	Tmax	Tmin	Ppt	Tmax	Tmin	Ppt	Tmax	Tmin	Ppt
Current	16.6	5.1	135.7	8.2	1.0	63.7	25.3	9.7	6.3
CCSM4-4.5	19.6	8.0	170.9	10.8	3.3	77.1	28.7	13.0	7.2
CCSM4-8.5	21.0	9.4	169.9	11.5	4.2	77.5	30.4	14.7	8.6
HadGEM-4.5	20.7	9.4	169.2	12.1	4.9	79.8	29.8	14.3	6.7
HadGEM-8.5	23.0	11.7	176.8	14.4	7.1	84.0	32.3	16.9	6.1
CanESM-4.5	20.2	9.0	191.6	11.0	4.2	94.1	30.5	14.5	5.9
CanESM-8.5	23.5	12.4	197.8	13.4	6.8	105.6	35.0	19.0	7.0

### Figure Legends

Figure 1. Management area (ownership type) map of the Panther Creek watershed, located in northwestern Oregon. White areas indicate lands that are nonforested and were not modeled.

Figure 2. Aboveground live carbon estimates for each ecoregion in the Panther Creek watershed from gradient nearest neighbor (GNN) imputation (year 2006) and LANDIS-II at the initiation of simulations (year 2010). Error bars show  $\pm 1SD$  across all cells within each ecoregion.

Ecoregions are defined based on annual precipitation level (high [HiPpt], medium [MidPpt] and

low [LowPpt]) and soil carbon (high [HiC], medium [MidC], and low [LowC]). See Ecoregions section for more detail.

Figure 3. Change over 90 years in ecosystem carbon pools across seven management scenarios under current climate (no climate change). Scenarios are described in Table 1. Panels depict total ecosystem carbon (a), aboveground live carbon (b), soil carbon (c), and aboveground detrital carbon (d). Note that the y-axis range varies among panels and that the soil carbon y-axis does not start at zero.

Figure 4. Change over 90 years in ecosystem carbon storage across seven climate scenarios (grouped into three categories), without harvest. The envelopes depict the mean  $\pm$  1SD under current climate (gray), three low forcing climate change scenarios (yellow), and three high forcing climate change scenarios (pink). Where envelopes overlap colors are blended (e.g., overlap between yellow and pink produces orange). Panels depict total ecosystem carbon (a), aboveground live carbon (b), soil carbon (c), and aboveground detrital carbon (d). Note that the y-axis range varies among panels and that the soil carbon y-axis does not start at zero.

Figure 5. Monthly temperature and water limitations to tree growth under current climate (left) and three climate models under the high forcing (RCP 8.5) climate change scenario (middle and right). Growth limit values closer to one indicate that a particular resource is unlimiting, and values close to zero indicate a strong limitation. Graphs depict growth limitations at year 2100 for a mixed species single-cell simulation, averaged across five replicates.





