

4-1-1986

Shell correction for the stopping power of K electrons

P.T. Leung
Portland State University

M. L. Rustgi

S. A. Long

Let us know how access to this document benefits you.

Follow this and additional works at: https://pdxscholar.library.pdx.edu/phy_fac

 Part of the [Physics Commons](#)

Citation Details

Leung, P. T., Rustgi, M. L., & Long, S. T. (1986). Shell correction for the stopping power of K electrons. *Physical Review A (General Physics)*, 33(4), 2328-2332.

This Article is brought to you for free and open access. It has been accepted for inclusion in Physics Faculty Publications and Presentations by an authorized administrator of PDXScholar. For more information, please contact pdxscholar@pdx.edu.

Shell correction for the stopping power of K electrons

P. T. Leung* and M. L. Rustgi

Physics Department, State University of New York at Buffalo, Buffalo, New York 14260

S. A. T. Long

Langley Research Center, National Aeronautics and Space Administration, Hampton, Virginia 23865

(Received 25 February 1985; revised manuscript received 12 August 1985)

In view of the inapplicability of the asymptotic expressions for the stopping number available in the literature at high energies, an alternative approach is taken to compute the shell correction to the stopping number of K electrons. Anholt's formula for the K -shell ionization has been used to calculate the excitation function for longitudinal interaction and numerical integration over energy has been carried out to evaluate the shell correction. Comparison with other theoretical calculations is made. It is proposed that, with the inclusion of relativistic effects, an asymptotic expansion of the stopping number with a leading-term logarithmic in the energy of the incident particle would be more meaningful and might enable one to extract the relativistic contribution to the shell correction from it.

INTRODUCTION

The evaluation of the shell corrections to the Bethe formula for the stopping power of matter has remained an active area of research ever since the pioneering work of Bethe and collaborators in the early 1950's.¹⁻⁵ By considering the optical oscillator strength and the mean excitation energy of the target atom, they were able to obtain some asymptotic formulas for the shellwise stopping number (B_i) for various target elements using screened hydrogenic orbitals. The shell correction (C_i) could then be extracted from a comparison made between the exact B_i obtained from numerical integration and the asymptotic B_i so obtained. The results of this approach have been reviewed in the literature by Uehling and Fano.^{6,7} Developments along this line have been carried on by Khandelwal⁸ to cover the entire Periodic Table for the K and L shells which have the most important contributions to shell corrections and, quite recently, improved asymptotic formulas have been obtained using Hartree-Slater oscillator strengths by Khandelwal.⁹ Aside from this line of development, recently there have been different approaches attempted by others such as the extraction of C_K and C_L from explicit Born-approximation calculations on subshell stopping powers by McGuire¹⁰ and the kinetic theoretical description by Sabin and Oddershede¹¹ based on Sigmund's formalism.¹² In spite of all these, Walske's results^{4,5} have many times been regarded as references for comparison between "hydrogenic calculations" with experimental or other theoretical results.¹⁰⁻¹⁴ It is the purpose of this paper to point out that due to the nonrelativistic nature, the asymptotic formulas given in the literature are not applicable for high-incident energies and hence the calculation of the shell correction by exploiting these formulas may not have general validity. Furthermore, it will be desirable to include the relativistic nature of the K electrons for heavy elements. We have therefore in this work followed an alternative approach to integrate

the shell correction numerically and the results are compared with those from other theoretical investigations. We shall limit ourselves only to the K shell contribution (C_K) in this paper and always take proton as the incident particle.

THE STOPPING NUMBERS

The asymptotic formulas of Walske⁴ and Khandelwal^{8,9} can be summarized as follows. The stopping number B_K is expanded in inverse power of the incident particle's velocity as follows:¹⁻³

$$B_K(\theta_K, \eta_K) = S_K(\theta_K) \ln \eta_K + T_K(\theta_K) - C_K(\theta_K, \eta_K) \quad (1)$$

with

$$C_K(\theta_K, \eta_K) = U_K(\theta_K) \eta_K^{-1} + V_K(\theta_K) \eta_K^{-2} + \dots \quad (2)$$

In (1) and (2), θ_K is the screening parameter which increases with the atomic number of the target atom and $\eta_K = v^2/Z_K^2$ with v being the incident particle's velocity and $Z_K = Z - 0.3$ the effective charge of the target atom. The coefficients S_K , T_K , U_K , and V_K are listed in Refs. 8 and 9 and we have used everything in atomic units in this paper. Because of the nonrelativistic nature of both the incident particle and the K electrons assumed in deriving these formulas, the results are not applicable in the limit of very large η_K 's corresponding to high incident energies or in the case with heavy target elements. For high-energy incident protons, it is more suitable to express η_K in the form

$$\left[\frac{137}{Z_K} \right]^2 \left[1 - \left[\frac{0.938}{E_p} \right]^2 \right]$$

as explained later on with E_p being the energy of the proton in GeV.

While B_K can be described by the asymptotic form using Eq. (1), it can also be obtained by exact integration. In the first Born approximation and by considering only the direct Coulomb interaction (i.e., only the longitudinal photon effect), B_K can be written as

$$B_K(\theta_K, \eta_K) = \int_{W_{\min}}^{\infty} dW \int_{Q_{\min}}^{\infty} dQ \frac{1}{Q} \phi_W(Q), \quad (3)$$

with

$$|\phi_W(Q)|^2 = \frac{2^7 d^2}{3} (3Q + W) Q \frac{\exp \left[-\frac{2}{k} \tan^{-1} [2k / (Q + 1 - k^2)] \right]}{[1 - \exp(-2\pi/k)] [(Q - k^2 + 1)^2 + 4k^2]^3}, \quad (5)$$

where

$$k^2 = W - 1, \quad (6)$$

and

$$d^2 = \left[1 + \left(\frac{Z_K \alpha}{2} \right)^2 \right]^{-1} \left[1 + (W - 1) \left(\frac{Z_K \alpha}{2} \right)^2 \right]^{-1} \quad (7)$$

arises from the normalization constants since we have used semirelativistic Darwin wave functions to describe the atomic K electron. Similar but somewhat different expressions have also been obtained by Davidović *et al.*¹⁷ but since Anholt's results agree better with experiment, we have here adopted his formalism. Previous formulations using Schrödinger wave functions would be equivalent to the above results by setting $d^2 = 1.1^{-5} \alpha$ in Eq. (7) stands for the fine-structure constant. Numerical integration for Eq. (3) has been carried out for various elements using a 256×256 points Gaussian quadrature for all energies. At low energies, our results reproduce the previous results if we set $d^2 = 1.4, 8$. At higher energies, where the asymptotic formulas of Walske and Khandelwal are not valid, our exactly integrated results for B_K with and without the effect of d^2 are shown in Fig. 1. It is clear from this figure that the use of the Darwin wave function gives very different results from those using the Schrödinger wave function for target elements with a large Z , since the K electrons of the heavy atoms are quite relativistic in nature. In fact, Anholt had pointed out earlier¹⁶ that the tables published previously^{8, 18} have to be modified due to the inclusion of the factor d^2 , although for light elements such as carbon there is essentially no difference with or without the d^2 term. However, for light elements, η_K becomes very large at large incident energies and hence Eqs. (1) and (2) cannot be applied. Taking into consideration this fact and the effect of the d^2 term, we conclude that the previous C_K 's calculated from Eqs. (1) and (2) are of questionable accuracy for high-incident energies for both light and

$$\phi_W(Q) = \frac{W |F_W(Q)|^2}{Q}, \quad (4)$$

where $F_W(Q)$ is the inelastic Coulomb form factor for the excitations and ionization of the target atom. W and Q are the energy and momentum transfers in atomic units. From kinematics and conservation of energy and momentum, one can set $W_{\min} = \theta_K$ and $Q_{\min} = W^2 / 4\eta_K$.¹⁵ For hydrogenlike wave functions, $|F_W(Q)|^2$ has the following form according to Anholt:¹⁶

heavy target elements. Furthermore, it is also worthwhile to point out that the expansion formulas for B_K given in the literature lose their asymptotic nature completely once the relativistic nature of the K electrons is taken into account (the d^2 factor) as shown in Fig. 2. In the following, therefore, we attempt to recompute C_K including the d^2 term by following an alternative method originally suggested in Walske's paper.⁴




FIG. 1. Stopping number B_K and the effect of d^2 on it. The solid line represents the exact results of this work and the dash-dotted line represents the integrated result without the factor d^2 . The curve labeled *a* is for C ($\theta_K = 0.64$) with the range of η_K being $60 < \eta_K < 600$; the integrated results with or without d^2 almost overlap with each other. The curve *b* is for Ni ($\theta_K = 0.8$) with the range of η_K being $5 < \eta_K < 25$ and the curve *c* is for Pm ($\theta_K = 0.9$) with the range of η_K being $1 < \eta_K < 5.5$.

CALCULATION OF C_K

Bethe, Brown, and Walske² have considered the following function:

$$\begin{aligned} X(\theta_K, \eta_K) &= \int_0^\infty dW \int_{1/4\eta_K}^{4\eta_K} dQ \frac{1}{Q} \phi_W(Q) - \int_{W_m}^\infty dW \int_{1/4\eta_K}^{W^2/4\eta_K} dQ \frac{1}{Q} \phi_W(0) \\ &\quad - \lim_{Q_1 \rightarrow 0} \int_0^{W_m} dW \int_{1/4\eta_K}^{Q_1} dQ \frac{1}{Q} \phi_W(0) - \lim_{Q_1 \rightarrow 0} \int_0^{W_m} dW \int_{Q_1}^\infty dQ \frac{1}{Q} \phi_W(Q) \end{aligned} \quad (8)$$

and have shown that $X \sim A \ln_K + B$ for all η_K and fixed θ_K . Hence by comparing (1) and (8), one can find C_K as

$$-C_K = B_K - X. \quad (9)$$

In what follows, we shall first rederive Eq. (8) of Ref. 4 in a bit simpler way than the original one and then apply numerical integration techniques to compute C_K from Eq. (9).

The first integral of Eq. (8) can be rewritten as

$$\begin{aligned} \int_0^\infty dW \int_{1/4\eta_K}^{4\eta_K} dQ \frac{1}{Q} \phi_W(Q) &= \int_{W_m}^\infty dW \int_{1/4\eta_K}^{4\eta_K} dQ \frac{1}{Q} \phi_W(Q) \\ &\quad + \int_0^{W_m} dW \int_{1/4\eta_K}^0 dQ \frac{1}{Q} \phi_W(Q) + \int_0^{W_m} dW \int_0^{4\eta_K} dQ \frac{1}{Q} \phi_W(Q). \end{aligned} \quad (10)$$

Substitute (10) into (8), recombine terms and we can write

$$\begin{aligned} X(\theta_K, \eta_K) &= \int_{W_m}^\infty dW \int_{1/4\eta_K}^{4\eta_K} dQ \frac{1}{Q} \phi_W(Q) - \int_{W_m}^0 dW \int_{1/4\eta_K}^{W^2/4\eta_K} dQ \frac{1}{Q} \phi_W(0) + \int_0^{W_m} dW \int_{1/4\eta_K}^0 dQ \frac{1}{Q} [\phi_W(Q) - \phi_W(0)] \\ &\quad - \int_0^{W_m} dW \int_{4\eta_K}^\infty dQ \frac{1}{Q} \phi_W(Q). \end{aligned} \quad (11)$$

Substituting (3) and (11) into (9), we get

$$\begin{aligned} -C_K &= \int_{W_m}^\infty dW \int_{W^2/4\eta_K}^\infty dQ \frac{1}{Q} \phi_W(Q) - \int_{W_m}^\infty dW \int_{1/4\eta_K}^{4\eta_K} dQ \frac{1}{Q} \phi_W(Q) + \int_{W_m}^\infty dW \int_{1/4\eta_K}^{W^2/4\eta_K} dQ \frac{1}{Q} \phi_W(0) \\ &\quad + \int_0^{W_m} dW \int_{4\eta_K}^\infty dQ \frac{1}{Q} \phi_W(Q) + \int_0^{W_m} dW \int_0^{1/4\eta_K} dQ \frac{1}{Q} [\phi_W(Q) - \phi_W(0)]. \end{aligned} \quad (12)$$

Let us denote the last integral by C' and rewrite it as follows:

$$C' = \int_0^\infty dW \int_0^{1/4\eta_K} dQ \frac{1}{Q} [\phi_W(Q) - \phi_W(0)] - \int_{W_m}^\infty dW \int_0^{1/4\eta_K} dQ \frac{1}{Q} [\phi_W(Q) - \phi_W(0)]. \quad (13)$$

Interchanging the order of integration in the first term in (13) and applying the Bethe sum rule¹⁹ in the form

$$\int_0^\infty dW \phi_W(Q) = \int_0^\infty dW \phi_W(0), \quad (14)$$

we get only the second term left for C' . Substituting back in (12) and neglecting the fourth term in (12) because it becomes less than 10^{-2} for $\eta_K \geq 0.5$, one obtains

$$\begin{aligned} -C_K &= \int_{W_m}^\infty dW \int_{W^2/4\eta_K}^\infty dQ \frac{1}{Q} \phi_W(Q) - \int_{W_m}^\infty dW \int_{1/4\eta_K}^{4\eta_K} dQ \frac{1}{Q} \phi_W(Q) + \int_{W_m}^\infty dW \int_{1/4\eta_K}^{W^2/4\eta_K} dW \frac{1}{Q} \phi_W(0) \\ &\quad - \int_{W_m}^\infty dW \int_0^{1/4\eta_K} dQ \frac{1}{Q} [\phi_W(Q) - \phi_W(0)]. \end{aligned} \quad (15)$$

If we rewrite the last term in (15) as

$$- \int_{W_m}^\infty dW \int_0^{W^2/4\eta_K} dQ \frac{1}{Q} [\phi_W(Q) - \phi_W(0)] - \int_{W_m}^\infty dW \int_{W^2/4\eta_K}^{1/4\eta_K} dQ \frac{1}{Q} \phi_W(Q) + \int_{W_m}^\infty dW \int_{W^2/4\eta_K}^{1/4\eta_K} dQ \frac{1}{Q} \phi_W(0), \quad (16)$$




FIG. 2. Comparison of the integrated results for B_K with the asymptotic formulas given in Ref. 8. The solid line represents the exact result with the factor d^2 . The dash-dotted line represents the integrated result without the factor d^2 and the dotted line represents the results of the asymptotic formulas.

substitute (16) back in (15), simplify and regroup terms, we arrive finally at Eq. (8) of Ref. (4),

$$\begin{aligned}
 -C_K = & \int_{w_m}^{\infty} dW \int_{4\eta_K}^{\infty} dQ \frac{1}{Q} \phi_W(Q) \\
 & - \int_{w_m}^{\infty} dW \int_0^{w^2/4\eta_K} dQ \frac{1}{Q} [\phi_W(Q) - \phi_W(0)].
 \end{aligned}
 \tag{17}$$

Note that we have only applied the sum rule once in our derivation. With the $\phi_W(Q)$ being given above [Eqs. (4)–(7)], we have computed C_K via numerical integration for various elements. Incidentally, we found that it is more convenient to manipulate with the four integrals in Eq. (15) than the two in Eq. (17) because of faster convergence and above all, the first term in Eq. (15) is just the B_K which we have computed before and the third term is independent of η_K . In Fig. 3 we show a comparison of the results obtained in this work and Walske's original results for two elements: Ni ($\theta_K=0.8$) and Pm ($\theta_K=0.9$). Figure 4 shows a comparison between the present result with the results from other calculations.^{4,10,11}

DISCUSSION AND CONCLUSION

It is seen in Fig. 3 that the results obtained in this work are in general smaller than those of Walske's original calculations. Furthermore, the results here seem to show a reverse trend as far as the screening parameter (θ_K) is con-




FIG. 3. Comparison between this work and that of Walske (Ref. 4) for C_K . The solid line represents the results of the present work and the dashed line represents Walske's results.

cerned. In fact, it will be very interesting if one can argue physically what the trend of the C_K should be with respect to the variation of atomic number of the target element. It should also be remarked that the curves in Fig. 3 are trustworthy only for not too high values of η_K since the expansion of C_K in $O(1/\eta_K)$ will never vanish since η_K approaches a maximum value as the velocity of the incident particle approaches that of light, whereas we expect that no shell correction is necessary for incident particles with infinitely high energies.²⁰ We suggest the lower limit $0.2 \leq 1/\eta_K$ for the C_K curves to be trustworthy. Furthermore, since the Bethe sum rule [Eq. (14)] holds exactly only for nonrelativistic atomic wave functions while we have been using Darwin wave functions in our formalism, the results for heavy target element (e.g., for $\theta_K=0.9$) will not be very accurate. However, since the term d^2 does not play an important role for light atoms (see Fig. 1), we expect that our results for Ni ($\theta_K=0.8$) and Al ($\theta_K=0.7$) would still be quite accurate.

The comparison of C_K for Al from different calculations is shown in Fig. 4. It is interesting to note that




FIG. 4. Comparison of C_K for Al from different theoretical calculations. *a* is the result of this work, *b* is Walske's results (Ref. 4), *c* is the result of McGuire (Ref. 10), and *d* is the result of Sabin and Oddershede (Ref. 11).




FIG. 5. Stopping number for Ni. *a* includes only longitudinal effect and *b* includes the sum of longitudinal, transverse, and spin-flip effect.

while the present result is quite close to Walske's original result, both results are consistently smaller than the result of McGuire.¹⁰ It may be more interesting to note that the result of Sabin and Oddershede¹¹ lies just between the two results mentioned above, being very close to the "hydrogenic results" for high-incident energies and very close to McGuire's result for low-incident energies. It would be of great interest if we can understand more physically the behaviors of these various results. Finally, it might also be of interest if one can find new asymptotic formulas for B_K which better fit the exact integrated results in Fig. 1 and reproduce the C_K curves in Fig. 4 at the same time.

However, from previous investigations²¹ the "saturation characteristics" of the B_K 's will disappear if we also include the relativistic effects, namely, the transverse and spin-flip effects.^{16,17} Figures 5 and 6 show the results for Ni and U from which we can see the importance of the spin-flip effect for heavy elements. Furthermore, it is found that with the inclusion of relativistic effects, B_K becomes almost linear in $\ln(E_p)$ and hence we propose that it might be more meaningful to seek asymptotic expansion




FIG. 6. Stopping number for U. *a* includes only longitudinal effect, *b* includes the sum of longitudinal and transverse effects, and *c* includes the sum of longitudinal, transverse, and spin-flip effects.

of B_K with a leading logarithmic term in energy (rather than in velocity) of the incident particle. In other words, η_K should be defined to be proportional to energy only if all these other relativistic effects are included. In that case, we might be able to extract the shell correction accurately for all energies with the relativistic effects (mainly the spin-flip effect for heavy target atoms) on C_K being included. Of course, one is always left with the possibility that other processes (e.g., pair creation in the atomic field of the target) may come in which suppress the B_K curve again towards saturation at high-incident energies so that an expansion in velocity ($\sim \eta_K$) with modified coefficients may still be possible. All these will be left as a future study in the relativistic effects on the shell corrections of Bethe's theory of stopping power of matter.

ACKNOWLEDGMENT

This work was partially supported by the U.S. National Aeronautics and Space Administration under Grant No. NAG1442. The authors are grateful to Dr. L. N. Pandey for his help with the numerical work.

*Permanent address: Physics Department, Tamkang University, Tansui, Taiwan 251, Republic of China.

¹L. M. Brown, Phys. Rev. **79**, 297 (1950).

²M. A. Bethe, L. M. Brown, and M. C. Walski, Phys. Rev. **79**, 413 (1950).

³M. C. Walske and H. A. Bethe, Phys. Rev. **83**, 457 (1951).

⁴M. C. Walske, Phys. Rev. **88**, 1283 (1952).

⁵M. C. Walske, Phys. Rev. **101**, 940 (1956).

⁶E. A. Uehling, Ann. Rev. Nucl. Sci. **4**, 315 (1954).

⁷U. Fano, Ann. Rev. Nucl. Sci. **13**, 1 (1963).

⁸G. S. Khandelwal, Nucl. Phys. **A116**, 97 (1968).

⁹G. S. Khandelwal, Phys. Rev. A **26**, 2983 (1982).

¹⁰E. J. McGuire, Phys. Rev. A **28**, 49 (1983); **28**, 53 (1983); **28**, 57 (1983).

¹¹J. R. Sabin and J. Oddershede, Phys. Rev. A **26**, 3209 (1982); **29**, 1757 (1984).

¹²P. Sigmund, Phys. Rev. A **26**, 2497 (1982).

¹³H. Bichsel and L. E. Porter, Phys. Rev. A **25**, 2499 (1982).

¹⁴H. H. Andersen *et al.*, Phys. Rev. A **16**, 1929 (1977).

¹⁵E. Merzbacher and H. W. Lewis, *Encyclopedia of Physics*, edited by S. Flügge (Springer-Verlag, Berlin, 1958), Vol. 34, p. 166.

¹⁶R. Anholt, Phys. Rev. A **19**, 1004 (1979).

¹⁷D. M. Davidović, B. L. Moiseiwitsch, and P. H. Norrington, J. Phys. B **11**, 947 (1978).

¹⁸G. S. Khandelwal, B. H. Choi, and E. Merzbacher, At. Data **1**, 103 (1969).

¹⁹See, e.g., H. A. Bethe and R. W. Jackiw, *Intermediate Quantum Mechanics* (Benjamin/Cummings, Reading, 1968), pp. 303–306.

²⁰See the remark by Fano, Ref. 7, Sec. 4.4.

²¹P. T. Leung and M. L. Rustgi, Phys. Rev. A **27**, 540 (1983).